



wwPDB EM Validation Summary Report ⓘ

Mar 10, 2026 – 01:39 AM UTC

PDB ID : 9CVT / pdb_00009cvt
EMDB ID : EMD-45966
Title : Melbournevirus Mini variant Nucleosome
Authors : Villalta, A.; Luger, K.
Deposited on : 2024-07-29
Resolution : 4.41 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 4.41 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	3133 (3.91 - 4.91)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	168	<div> <div>5%</div> <div>57%</div> <div>33%</div> <div>10%</div> </div>
1	B	168	<div> <div>55%</div> <div>42%</div> <div>..</div> </div>
2	C	216	<div> <div>54%</div> <div>26%</div> <div>20%</div> </div>
2	D	216	<div> <div>50%</div> <div>28%</div> <div>22%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	I	147	<div><div></div><div>14%</div><div>48%</div><div>38%</div></div>
4	J	147	<div><div></div><div>19%</div><div>43%</div><div>38%</div></div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8897 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone doublet miniH2B-H2A.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	151	Total	C	N	O	S	0	0
			1213	773	212	222	6		
1	B	164	Total	C	N	O	S	0	0
			1308	831	230	241	6		

- Molecule 2 is a protein called Histone doublet H4-H3.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	C	172	Total	C	N	O	S	0	0
			1337	847	241	239	10		
2	D	168	Total	C	N	O	S	0	0
			1308	828	235	235	10		

- Molecule 3 is a DNA chain called Widom 601 Strand 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	I	91	Total	C	N	O	P	0	0
			1851	879	333	548	91		

- Molecule 4 is a DNA chain called Widom 601 Strand 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	J	91	Total	C	N	O	P	0	0
			1880	888	357	544	91		

E178	T181	V192	R193	L194	L195	E196	K197	I191	S194	R197	T198	R199	S202	K203	D204	I205	E206	M210	L211	E212	HIS	GLY	PRO	LEU	T100	LVS	ASP	SER	GLY	ALA	LVS	GLY	VAL	SER	VAL	ALA	PHE	LEU	SER	CYS	ARG	GLN	LVS	GLY	ALA	GLY	ALA	SER	ALA	GLY	THR	ASN	SER	GLN	E131	Q135	C140	L141	I142	I143	E146	R147	F148	R149	T150	M151	I155	E159	G160	H161	I165	A166	L170	Q174	V175	Met	SER	LYS	ALA	GLY	LVS	VAL	LVS	ALA	GLN	GLN	HIS	GLY	T25	Q26	I27	P28	K29	T32	Q33	H34	R37	A44	A45	T48	E49	V50	P51	I52	R53	K56	L59	A70	M71	R76	K77	T78	I79	M80	K81	K86	L90	M91	H92	L93	I96
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	-----	-----	-----	-----	------	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

Chain I: 14% 48% 38%

DC	DC	C-12	DA
DG	DT	G-11	DT
DG	DG	C-8	DG
DT	DT	G-7	DG
DC	DC	T-6	DA
DA	DA	A-5	DG
DG	DG	C-4	DA
DA	DA	G-3	DA
DT	DT	C-2	DT
DA	DA	G-1	DC
DT	DT	G-0	DC
DA	DA	T1	DG
DT	DT	G2	DG
DA	DA	T3	DT
DC	DC	C4	DG
DA	DA	C5	DC
DT	DT	C6	DC
DC	DC	C7	DG
DC	DC	C8	DA
DA	DA	G9	DG
DA	DA	C10	DG
DT	DT	G11	DC
		T12	DC
		T13	DG
		T14	DC
		T15	DT
		A16	DC
		A17	DA
		C18	DA
		C19	DT
		G20	T-42
			G-41
		A24	G-40
		G25	T-39
		G26	C-38
		G27	G-37
		G28	T-36
		A29	A-35
		T30	G-34
		T31	A-33
		A32	C-32
		C33	A-31
		T34	G-30
		C35	C-29
		C36	T-28
		C37	C-27
		T38	T-26
		A39	A-25
		G40	G-24
		T41	C-23
		C42	A-22
		T43	
		C44	G-19
		C45	T-17
		A46	T-16
		G47	A-15
		G48	A-14
		DC	

Chain J:

[illegible]

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	36051	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	20000	Depositor
Magnification	Not provided	
Image detector	FEI FALCON IV (4k x 4k)	Depositor
Maximum map value	0.098	Depositor
Minimum map value	-0.050	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.005	Depositor
Recommended contour level	0.0222	Depositor
Map size (Å)	217.28, 217.28, 217.28	wwPDB
Map dimensions	448, 448, 448	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.48499998, 0.48499998, 0.48499998	Depositor

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.18	0/1228	0.42	0/1643
1	B	0.19	0/1326	0.45	0/1775
2	C	0.19	0/1355	0.43	0/1818
2	D	0.18	0/1324	0.41	0/1775
3	I	0.25	0/2072	0.48	0/3192
4	J	0.24	0/2112	0.42	0/3261
All	All	0.21	0/9417	0.44	0/13464

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1213	0	1268	40	0
1	B	1308	0	1362	50	0
2	C	1337	0	1426	40	0
2	D	1308	0	1398	54	0
3	I	1851	0	1022	56	0
4	J	1880	0	1020	55	0
All	All	8897	0	7496	276	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 276 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:165:ILE:HD12	2:D:170:LEU:HD21	1.57	0.83
2:C:32:THR:HB	2:C:52:ILE:HD11	1.61	0.83
1:A:43:GLN:HE22	2:D:90:LEU:HB2	1.48	0.77
2:D:135:GLN:OE1	2:D:135:GLN:N	2.20	0.74
1:B:43:GLN:HE22	2:C:90:LEU:HD11	1.51	0.74

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	147/168 (88%)	139 (95%)	8 (5%)	0	100	100
1	B	162/168 (96%)	143 (88%)	17 (10%)	2 (1%)	10	42
2	C	168/216 (78%)	160 (95%)	8 (5%)	0	100	100
2	D	164/216 (76%)	160 (98%)	4 (2%)	0	100	100
All	All	641/768 (84%)	602 (94%)	37 (6%)	2 (0%)	37	71

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	162	LYS
1	B	18	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	135/148 (91%)	135 (100%)	0	100	100
1	B	144/148 (97%)	144 (100%)	0	100	100
2	C	149/180 (83%)	149 (100%)	0	100	100
2	D	146/180 (81%)	146 (100%)	0	100	100
All	All	574/656 (88%)	574 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	157	HIS
2	D	34	HIS
1	A	101	ASN
1	B	74	GLN
1	B	124	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-45966. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 224




Y Index: 224




Z Index: 224


6.2.2 Raw map



X Index: 224



Y Index: 224




Z Index: 224


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 224




Y Index: 207




Z Index: 210


6.3.2 Raw map



X Index: 0



Y Index: 0




Z Index: 0


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)

6.4.1 Primary map



6.4.2 Raw map



The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

This section was not generated.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 98 nm³; this corresponds to an approximate mass of 88 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.227 \AA^{-1}

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.227 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	4.41	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	6.35	7.60	6.54

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 6.35 differs from the reported value 4.41 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-45966 and PDB model 9CVT. Per-residue inclusion information can be found in [section 3](#) on [page 5](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.0222 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0222).

9.4 Atom inclusion [i](#)



At the recommended contour level, 97% of all backbone atoms, 85% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.0222) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div><div></div>0.8520</div>	<div><div></div>0.3210</div>
A	<div><div></div>0.7300</div>	<div><div></div>0.3110</div>
B	<div><div></div>0.7650</div>	<div><div></div>0.3290</div>
C	<div><div></div>0.8450</div>	<div><div></div>0.3650</div>
D	<div><div></div>0.8120</div>	<div><div></div>0.3480</div>
I	<div><div></div>0.9310</div>	<div><div></div>0.2950</div>
J	<div><div></div>0.9430</div>	<div><div></div>0.2970</div>

1.0

0.0

<0.0