Full wwPDB X-ray Structure Validation Report (i)

Sep 26, 2023 – 03:45 PM EDT

PDB ID	:	6CYT
Title	:	HIV-1 TAR loop in complex with Tat:AFF4:P-TEFb
Authors	:	Schulze Gahmen, U.; Hurley, J.H.
Deposited on	:	2018-04-06
Resolution	:	3.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1659 (3.60-3.40)
Clashscore	141614	1036 (3.58-3.42)
Ramachandran outliers	138981	1005 (3.58-3.42)
Sidechain outliers	138945	1006 (3.58-3.42)
RSRZ outliers	127900	1559 (3.60-3.40)
RNA backbone	3102	1002 (4.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	332	2%, 92%, ·, ·
2	B	264	%, 90%, 6%, ·
3	C	36	3%, 64%, 14%, 8%, 14%
4	D	58	2%, 78%, 7%, 16%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	N	20	20% 50% 15% 15%

2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 5621 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cyclin-dependent kinase 9.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	319	2561	1640	442	463	1	15	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP P50750
A	0	HIS	-	expression tag	UNP P50750

- Molecule 2 is a protein called Cyclin-T1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	255	2075	1327	358	380	10	0	0	0

- Molecule 3 is a protein called AF4/FMR2 family member 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	31	242	152	38	50	2	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	47	TYR	LYS	conflict	UNP Q9UHB7

- Molecule 4 is a protein called Protein Tat.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	49	377	240	64	65	8	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	0	ACE	-	acetylation	UNP A0A0C5HAL9

- Molecule 5 is a RNA chain called RNA (5'-R(P*AP*UP*CP*UP*GP*AP*GP*CP*CP*UP *GP*GP*GP*AP*GP*CP*U)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
5	N	17	364	162	65	120	17	0	0	0

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
6	D	2	2	2	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Cyclin-dependent kinase 9

Chain A:





• Molecule 2: Cyclin-T1

Chain B:



• Molecule 3: AF4/FMR2 family member 4

Chain C:



• Molecule 4: Protein Tat

Chain D:



• Molecule 5: RNA (5'-R(P*AP*UP*CP*UP*GP*AP*GP*CP*CP*UP*GP*GP*GP*AP*GP*CP*U)-3')

Chain N:

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α, β, γ	148.58Å 148.58Å 104.63Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	128.67 – 3.50 128.67 – 3.45	Depositor EDS
% Data completeness (in resolution range)	99.5 (128.67-3.50) 84.6 (128.67-3.45)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	0.36 (at 3.49Å)	Xtriage
Refinement program	PHENIX 1.13_2998	Depositor
R, R_{free}	0.247 , 0.283 0.245 , 0.277	Depositor DCC
R_{free} test set	1788 reflections (10.01%)	wwPDB-VP
Wilson B-factor (Å^2)	106.5	Xtriage
Anisotropy	0.325	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.30 , 82.9	EDS
L-test for twinning[2]	$< \|L\| > = 0.48$, $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	0.038 for -h,-k,l	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	5621	wwPDB-VP
Average B, all atoms (Å^2)	154.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.01% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, ACE, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.26	0/2600	0.46	0/3510
2	B	0.25	0/2129	0.42	0/2904
3	C	0.24	0/245	0.36	0/329
4	D	0.26	0/387	0.45	0/524
5	N	0.25	0/406	0.83	0/631
All	All	0.26	0/5767	0.48	0/7898

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2561	0	2600	8	0
2	B	2075	0	2039	12	0
3	C	242	0	220	4	0
4	D	377	0	364	4	0
5	N	364	0	184	7	0
6	D	2	0	0	0	0
All	All	5621	0	5407	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 3.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:N:33:G:O2'	5:N:34:G:O4'	2.13	0.62
2:B:259:ARG:HH11	4:D:26:TYR:HB2	1.67	0.60
1:A:261:LEU:H	1:A:261:LEU:HD23	1.67	0.58
5:N:25:U:H2'	5:N:26:G:C8	2.40	0.57
1:A:261:LEU:HD23	1:A:261:LEU:N	2.20	0.56
3:C:51:ARG:NH2	3:C:55:MET:SD	2.79	0.56
1:A:119:VAL:O	1:A:225:ARG:NH2	2.38	0.54
2:B:209:ASN:OD1	3:C:53:GLN:NE2	2.41	0.51
5:N:29:C:N4	5:N:30:C:H41	2.09	0.50
1:A:177:ALA:HB1	1:A:178:LYS:CG	2.42	0.50
2:B:249:PRO:HG2	4:D:43:LEU:O	2.11	0.49
1:A:177:ALA:HB1	1:A:178:LYS:HG2	1.94	0.49
3:C:61:GLU:O	4:D:28:LYS:NZ	2.46	0.49
2:B:252:LEU:O	2:B:256:TRP:HB2	2.13	0.49
2:B:251:ARG:HG2	2:B:255:ILE:HG23	1.96	0.48
2:B:251:ARG:NE	2:B:255:ILE:HG21	2.28	0.48
5:N:27:A:H2'	5:N:28:G:C8	2.49	0.47
1:A:30:PHE:O	1:A:51:LEU:HB3	2.16	0.46
5:N:32:G:H4'	5:N:33:G:OP2	2.16	0.46
2:B:251:ARG:HH11	2:B:251:ARG:HG3	1.82	0.44
1:A:178:LYS:O	1:A:179:ASN:HB2	2.18	0.44
5:N:32:G:H21	5:N:32:G:P	2.41	0.43
2:B:251:ARG:HG2	2:B:255:ILE:CG2	2.48	0.43
2:B:251:ARG:O	2:B:255:ILE:HG12	2.19	0.43
1:A:64:LEU:HD22	2:B:148:LEU:HD13	2.02	0.42
5:N:30:C:N4	5:N:36:G:O6	2.53	0.41
3:C:62:MET:SD	3:C:62:MET:N	2.94	0.41
2:B:249:PRO:O	2:B:251:ARG:N	2.47	0.41
2:B:158:VAL:HG11	4:D:6:PRO:HD3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	314/332 (95%)	295 (94%)	16 (5%)	3 (1%)	15	54
2	B	253/264 (96%)	241 (95%)	12 (5%)	0	100	100
3	C	29/36 (81%)	20 (69%)	7 (24%)	2 (7%)	1	12
4	D	47/58 (81%)	46 (98%)	1 (2%)	0	100	100
All	All	643/690 (93%)	602 (94%)	36 (6%)	5 (1%)	19	58

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	190	VAL
3	C	58	ASN
1	A	266	GLU
1	A	325	LYS
3	C	57	GLY

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	277/292 (95%)	274 (99%)	3 (1%)	73	88
2	B	231/239 (97%)	222 (96%)	9 (4%)	32	64
3	C	25/33 (76%)	21 (84%)	4 (16%)	2	14
4	D	43/52 (83%)	43 (100%)	0	100	100
All	All	576/616 (94%)	560 (97%)	16 (3%)	43	72

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	62	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	243	GLN
1	A	261	LEU
2	B	8	ASN
2	B	25	ARG
2	B	35	LEU
2	B	116	GLU
2	B	136	LEU
2	B	241	PHE
2	B	248	THR
2	B	252	LEU
2	B	261	CYS
3	C	35	PHE
3	C	51	ARG
3	C	55	MET
3	C	62	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
5	N	16/20 (80%)	6 (37%)	0

All (6) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
5	N	31	U
5	N	32	G
5	N	33	G
5	N	35	A
5	N	36	G
5	N	37	C

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul

statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
1	TPO	A	186	1	8,10,11	1.56	1 (12%)	10,14,16	1.69	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	A	186	1	-	1/9/11/13	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	186	TPO	P-O1P	3.27	1.61	1.50

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	186	TPO	P-OG1-CB	-4.67	109.12	123.21

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	186	TPO	O-C-CA-CB

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	318/332 (95%)	0.17	5 (1%)	72	66	90, 121, 175, 289	0
2	B	255/264 (96%)	0.09	2 (0%)	86	81	103, 157, 207, 223	0
3	C	31/36 (86%)	-0.05	1 (3%)	47	42	173, 213, 224, 228	0
4	D	48/58 (82%)	0.02	1 (2%)	63	58	134, 165, 192, 199	0
5	N	17/20 (85%)	-0.53	0	100	100	215, 256, 285, 286	0
All	All	669/710 (94%)	0.10	9 (1%)	77	71	90, 145, 218, 289	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	30	PHE	5.6
1	A	29	THR	5.1
3	C	54	SER	3.1
1	A	330	THR	2.8
2	B	7	ASN	2.6
4	D	29	LYS	2.3
1	A	179	ASN	2.0
2	B	118	LEU	2.0
1	A	241	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
1	TPO	A	186	11/12	0.92	0.21	104,109,115,116	0

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
6	ZN	D	102	1/1	0.94	0.10	192,192,192,192	1
6	ZN	D	101	1/1	0.95	0.20	167,167,167,167	0

6.5 Other polymers (i)

There are no such residues in this entry.