# wwPDB NMR Structure Validation Summary Report (i)

Dec 11, 2022 – 12:24 AM EST

| | | |
|---|---|---|
| PDB ID | : | 1D1D |
| Title | : | NMR SOLUTION STRUCTURE OF THE CAPSID PROTEIN FROM ROUS SARCOMA VIRUS |
| Authors | : | Campos-Olivas, R.; Newman, J.L.; Summers, M.F. |
| Deposited on | : | 1999-09-15 |

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

| | | |
|---|---|---|
| MolProbity | : | 4.02b-467 |
| Percentile statistics | : | 20191225.v01 (using entries in the PDB archive December 25th 2019) |
| RCI | : | v_1n_11_5_13_A (Berjanski et al., 2005) |
| PANAV | : | Wang et al. (2010) |
| ShiftChecker | : | 2.31.2 |
| Ideal geometry (proteins) | : | Engh & Huber (2001) |
| Ideal geometry (DNA, RNA) | : | Parkinson et al. (1996) |
| Validation Pipeline (wwPDB-VP) | : | 2.31.2 |

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
*SOLUTION NMR*

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




| Metric | Whole archive (#Entries) | NMR archive (#Entries) |
|---|---|---|
| Clashscore | 158937 | 12864 |
| Ramachandran outliers | 154571 | 11451 |
| Sidechain outliers | 154315 | 11428 |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

| Mol | Chain | Length | Quality of chain |
|---|---|---|---|
| 1 | A | 262 | 30% 39% 7% 8% 16% |

# 2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 8 is the overall representative, medoid model (most similar to other models). The authors have identified model 7 as representative.

The following residues are included in the computation of the global validation metrics.

| Well-defined (core) protein residues | | | |
|---|---|---|---|
| **Well-defined core** | **Residue range (total)** | **Backbone RMSD (Å)** | **Medoid model** |
| 1 | A:15-A:93, A:100-A:145 (125) | 0.48 | 8 |
| 2 | A:152-A:226 (75) | 0.28 | 19 |

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 1 single-model cluster was found.

| Cluster number | Models |
|---|---|
| 1 | 2, 3, 6, 11, 14, 16, 17 |
| 2 | 5, 8, 9, 10, 12, 19 |
| 3 | 7, 15, 18, 20 |
| 4 | 1, 13 |
| Single-model clusters | 4 |

# 3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 3395 atoms, of which 1721 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called PROTEIN (CAPSID PROTEIN).

| Mol | Chain | Residues | Atoms | | | | | | Trace |
|---|---|---|---|---|---|---|---|---|---|
| | | | Total | C | H | N | O | S | |
| 1 | A | 220 | 3395 | 1058 | 1721 | 303 | 307 | 6 | 0 |

There are 21 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|---|---|---|---|---|---|
| A | -22 | GLY | - | expression tag | UNP O92954 |
| A | -21 | HIS | - | expression tag | UNP O92954 |
| A | -20 | HIS | - | expression tag | UNP O92954 |
| A | -19 | HIS | - | expression tag | UNP O92954 |
| A | -18 | HIS | - | expression tag | UNP O92954 |
| A | -17 | HIS | - | expression tag | UNP O92954 |
| A | -16 | HIS | - | expression tag | UNP O92954 |
| A | -15 | HIS | - | expression tag | UNP O92954 |
| A | -14 | HIS | - | expression tag | UNP O92954 |
| A | -13 | HIS | - | expression tag | UNP O92954 |
| A | -12 | HIS | - | expression tag | UNP O92954 |
| A | -11 | SER | - | expression tag | UNP O92954 |
| A | -10 | SER | - | expression tag | UNP O92954 |
| A | -9 | GLY | - | expression tag | UNP O92954 |
| A | -8 | HIS | - | expression tag | UNP O92954 |
| A | -7 | ILE | - | expression tag | UNP O92954 |
| A | -6 | GLU | - | expression tag | UNP O92954 |
| A | -5 | GLY | - | expression tag | UNP O92954 |
| A | -4 | ARG | - | expression tag | UNP O92954 |
| A | -3 | HIS | - | expression tag | UNP O92954 |
| A | -2 | ASN | - | expression tag | UNP O92954 |

# 4 Residue-property plots (i)

## 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: PROTEIN (CAPSID PROTEIN)

Chain A: 30% 39% 7% 8% 16%




## 4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 8. Colouring as in section 4.1 above.

• Molecule 1: PROTEIN (CAPSID PROTEIN)

Chain A: 29% 39% 8% 8% 16%



R185
A186
P187
V188
I189
I190
D191
R194
Q195
K196
S197
Q198
P199
D200
I201
Q202
Q203
L204
I205
R206
A207
A208
P209
S210
T211
L212
T213
T214
P215
G216
E217
T218
I219
K220
Y221
Y222
L223
D224
R225
Q226
K227
I228
A229
P230
LEU
THR
ASP
GLN
GLY
ILE
ALA
ALA
ALA
MET

# 5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *TORSION ANGLE DYNAMICS SIMULATED ANNEALING*.

Of the 20 calculated structures, 20 were deposited, based on the following criterion: *target function*.

The following table shows the software used for structure solution, optimisation and refinement.

| Software name | Classification | Version |
|---|---|---|
| DYANA | refinement | 1.5 |
| XwinNMR | structure solution | 2.5 |
| NMRPipe | structure solution | 1999 |
| NMRView | structure solution | 3.0 |
| TALOS | structure solution | 1999.019.15.47 |

No chemical shift data was provided.

# 6 Model quality (i)

## 6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

## 6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes |
|---|---|---|---|---|---|
| 1 | A | 1530 | 1577 | 1575 | 114±10 |
| All | All | 30600 | 31540 | 31500 | 2282 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 37.

5 of 687 unique clashes are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Clash(Å) | Distance(Å) | Models | |
|---|---|---|---|---|---|
| | | | | Worst | Total |
| 1:A:43:LEU:CD1 | 1:A:56:LEU:HD21 | 0.95 | 1.90 | 11 | 16 |
| 1:A:57:MET:O | 1:A:61:LEU:HD12 | 0.92 | 1.65 | 3 | 5 |
| 1:A:113:MET:SD | 1:A:124:LEU:HD21 | 0.91 | 2.05 | 15 | 6 |
| 1:A:44:MET:SD | 1:A:48:LEU:HD21 | 0.90 | 2.05 | 13 | 1 |
| 1:A:81:ILE:HD13 | 1:A:103:LEU:HD12 | 0.90 | 1.41 | 13 | 3 |

## 6.3 Torsion angles (i)

### 6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles |
|---|---|---|---|---|---|---|
| 1 | A | 200/262 (76%) | 184±2 (92±1%) | 9±2 (4±1%) | 7±1 (3±1%) | 6 35 |
| All | All | 4000/5240 (76%) | 3690 (92%) | 171 (4%) | 139 (3%) | 6 35 |

5 of 14 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

| Mol | Chain | Res | Type | Models (Total) |
|---|---|---|---|---|
| 1 | A | 186 | ALA | 19 |
| 1 | A | 216 | GLY | 19 |
| 1 | A | 183 | SER | 18 |
| 1 | A | 45 | SER | 14 |
| 1 | A | 117 | PRO | 13 |

### 6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles |
|---|---|---|---|---|---|
| 1 | A | 164/210 (78%) | 121±4 (74±2%) | 43±4 (26±2%) | 2 23 |
| All | All | 3280/4200 (78%) | 2426 (74%) | 854 (26%) | 2 23 |

5 of 101 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

| Mol | Chain | Res | Type | Models (Total) |
|---|---|---|---|---|
| 1 | A | 35 | ILE | 20 |
| 1 | A | 48 | LEU | 20 |
| 1 | A | 90 | HIS | 20 |
| 1 | A | 125 | ARG | 20 |
| 1 | A | 137 | LEU | 20 |

### 6.3.3 RNA (i)

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

## 6.6 Ligand geometry (i)

There are no ligands in this entry.

## 6.7 Other polymers (i)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

# 7 Chemical shift validation (i)

No chemical shift data were provided