# wwPDB X-ray Structure Validation Summary Report (i)

Nov 20, 2023 – 07:21 PM JST

| | | |
|---|---|---|
| PDB ID | : | 7D92 |
| Title | : | Crystal Structure of the Na+,K+-ATPase in the E2P state with bound Mg2+ and anthroylouabain (P4(3)2(1)2 symmetry) |
| Authors | : | Kanai, R.; Cornelius, F.; Ogawa, H.; Motoyama, K.; Vilsen, B.; Toyoshima, C. |
| Deposited on | : | 2020-10-12 |
| Resolution | : | 3.90 Å(reported) |

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

---

The following versions of software and data (see references (i)) were used in the production of this report:

| | | |
|---|---|---|
| MolProbity | : | 4.02b-467 |
| Mogul | : | 1.8.5 (274361), CSD as541be (2020) |
| Xtriage (Phenix) | : | 1.13 |
| EDS | : | 2.36 |
| buster-report | : | 1.1.7 (2018) |
| Percentile statistics | : | 20191225.v01 (using entries in the PDB archive December 25th 2019) |
| Refmac | : | 5.8.0158 |
| CCP4 | : | 7.0.044 (Gargrove) |
| Ideal geometry (proteins) | : | Engh & Huber (2001) |
| Ideal geometry (DNA, RNA) | : | Parkinson et al. (1996) |
| Validation Pipeline (wwPDB-VP) | : | 2.36 |

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




| Metric | Whole archive (#Entries) | Similar resolution (#Entries, resolution range(Å)) |
|---|---|---|
| $R_{free}$ | 130704 | 1002 (4.14-3.66) |
| Clashscore | 141614 | 1004 (4.12-3.68) |
| Ramachandran outliers | 138981 | 1021 (4.14-3.66) |
| Sidechain outliers | 138945 | 1014 (4.14-3.66) |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

| Mol | Chain | Length | Quality of chain |
|---|---|---|---|
| 1 | A | 1016 | 74% 23% • • |
| 2 | B | 303 | 66% 27% • • |
| 3 | G | 65 | 42% 8% 51% |
| 4 | C | 2 | 100% |

# 2 Entry composition (i)

There are 10 unique types of molecules in this entry. The entry contains 10740 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sodium/potassium-transporting ATPase subunit alpha-1.

| Mol | Chain | Residues | Atoms | | | | | | | ZeroOcc | AltConf | Trace |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | | Total | Be | C | F | N | O | S | | | |
| 1 | A | 996 | 7730 | 1 | 4922 | 3 | 1301 | 1456 | 47 | 0 | 0 | 0 |

- Molecule 2 is a protein called Sodium/potassium-transporting ATPase subunit beta-1.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | Total | C | N | O | S | | | |
| 2 | B | 291 | 2386 | 1546 | 390 | 437 | 13 | 0 | 0 | 0 |

- Molecule 3 is a protein called FXYD domain-containing ion transport regulator.

| Mol | Chain | Residues | Atoms | | | | ZeroOcc | AltConf | Trace |
|---|---|---|---|---|---|---|---|---|---|
| | | | Total | C | N | O | | | |
| 3 | G | 32 | 255 | 174 | 37 | 44 | 0 | 0 | 0 |

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.

| Mol | Chain | Residues | Atoms | | | | ZeroOcc | AltConf | Trace |
|---|---|---|---|---|---|---|---|---|---|
| | | | Total | C | N | O | | | |
| 4 | C | 2 | 28 | 16 | 2 | 10 | 0 | 0 | 0 |

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

| Mol | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|---|---|---|---|---|---|---|
| | | | Total | Mg | | |
| 5 | A | 3 | 3 | 3 | 0 | 0 |

- Molecule 6 is CHOLESTEROL (three-letter code: CLR) (formula: $C_{27}H_{46}O$).




| Mol | Chain | Residues | Atoms | | | ZeroOcc | AltConf |
|---|---|---|---|---|---|---|---|
| | | | Total | C | O | | |
| 6 | A | 1 | 28 | 27 | 1 | 0 | 0 |
| | | | Total | C | O | | |
| 6 | B | 1 | 28 | 27 | 1 | 0 | 0 |
| | | | Total | C | O | | |
| 6 | G | 1 | 28 | 27 | 1 | 0 | 0 |

- Molecule 7 is 1,2-DIOLEOYL-SN-GLYCERO-3-PHOSPHOCHOLINE (three-letter code: PCW) (formula: $C_{44}H_{85}NO_8P$).



| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf |
|---|---|---|---|---|---|---|---|---|---|
| | | | Total | C | N | O | P | | |
| 7 | A | 1 | 22 | 12 | 1 | 8 | 1 | 0 | 0 |
| 7 | A | 1 | 22 | 12 | 1 | 8 | 1 | 0 | 0 |
| 7 | A | 1 | 22 | 12 | 1 | 8 | 1 | 0 | 0 |
| 7 | A | 1 | 22 | 12 | 1 | 8 | 1 | 0 | 0 |
| 7 | A | 1 | 22 | 12 | 1 | 8 | 1 | 0 | 0 |
| 7 | A | 1 | 22 | 12 | 1 | 8 | 1 | 0 | 0 |
| 7 | A | 1 | 22 | 12 | 1 | 8 | 1 | 0 | 0 |
| 7 | A | 1 | 22 | 12 | 1 | 8 | 1 | 0 | 0 |

- Molecule 8 is [(2 {R},3 {R},4 {R},5 {S},6 {S})-2-[[(1 {R},3 {S},5 {S},8 {R},9 {S},10 {R},11 {R},13 {R},14 {S},17 {R})-10-(hydroxymethyl)-13-methyl-1,5,11,14-tetrakis(oxidanyl)-17-(5-oxidanylidene-2 {H}-furan-3-yl)-2,3,4,6,7,8,9,11,12,15,16,17-dodecahydro-1 {H}-cyclopenta[a]phenanthren-3-yl]oxy]-6-methyl-3,5-bis(oxidanyl)oxan-4-yl] anthracene-9-carboxylate (three-letter code: H0C) (formula: $C_{44}H_{52}O_{13}$) (labeled as "Ligand of Interest" by depositor).




| Mol | Chain | Residues | Atoms | | | ZeroOcc | AltConf |
|---|---|---|---|---|---|---|---|
| | | | Total | C | O | | |
| 8 | A | 1 | 57 | 44 | 13 | 0 | 0 |

- Molecule 9 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).




| Mol | Chain | Residues | Atoms | | | | ZeroOcc | AltConf |
|---|---|---|---|---|---|---|---|---|
| | | | Total | C | N | O | | |
| 9 | B | 1 | 14 | 8 | 1 | 5 | 0 | 0 |

- Molecule 10 is water.

| Mol | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|---|---|---|---|---|---|---|
| | | | Total | O | | |
| 10 | A | 7 | 7 | 7 | 0 | 0 |

# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Sodium/potassium-transporting ATPase subunit alpha-1

Chain A:




- Molecule 2: Sodium/potassium-transporting ATPase subunit beta-1

Chain B:






• Molecule 3: FXYD domain-containing ion transport regulator

Chain G: 42% 8% 51%




• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C: 100%

# 4 Data and refinement statistics (i)

| Property | Value | Source |
|---|---|---|
| Space group | P 43 21 2 | Depositor |
| Cell constants<br>a, b, c, $\alpha$, $\beta$, $\gamma$ | 84.23Å 84.23Å 646.26Å<br>90.00° 90.00° 90.00° | Depositor |
| Resolution (Å) | 14.94 – 3.90<br>48.19 – 3.90 | Depositor<br>EDS |
| % Data completeness<br>(in resolution range) | 41.4 (14.94-3.90)<br>42.5 (48.19-3.90) | Depositor<br>EDS |
| $R_{merge}$ | 0.22 | Depositor |
| $R_{sym}$ | (Not available) | Depositor |
| $< I/\sigma(I) >$ [1] | 1.21 (at 3.88Å) | Xtriage |
| Refinement program | PHENIX 1.17.1_3660 | Depositor |
| R, $R_{free}$ | 0.209 , 0.257<br>0.249 , 0.293 | Depositor<br>DCC |
| $R_{free}$ test set | 484 reflections (5.01%) | wwPDB-VP |
| Wilson B-factor (Å$^2$) | 112.6 | Xtriage |
| Anisotropy | 0.653 | Xtriage |
| Bulk solvent $k_{sol}$(e/Å$^3$), $B_{sol}$(Å$^2$) | 0.28 , 64.9 | EDS |
| L-test for twinning[2] | $< \|L\| > = 0.42$, $< L^2 > = 0.25$ | Xtriage |
| Estimated twinning fraction | No twinning to report. | Xtriage |
| $F_o$,$F_c$ correlation | 0.84 | EDS |
| Total number of atoms | 10740 | wwPDB-VP |
| Average B, all atoms (Å$^2$) | 149.0 | wwPDB-VP |

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.05% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, PCW, BFD, NAG, CLR, H0C

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|---|---|---|---|---|---|
| | | RMSZ | $\#\|Z\| >5$ | RMSZ | $\#\|Z\| >5$ |
| 1 | A | 0.29 | 0/7867 | 0.50 | 1/10674 (0.0%) |
| 2 | B | 0.30 | 0/2449 | 0.54 | 0/3301 |
| 3 | G | 0.29 | 0/261 | 0.42 | 0/354 |
| All | All | 0.29 | 0/10577 | 0.51 | 1/14329 (0.0%) |

There are no bond length outliers.

All (1) bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed($^o$) | Ideal($^o$) |
|---|---|---|---|---|---|---|---|
| 1 | A | 492 | ALA | C-N-CA | 6.27 | 137.38 | 121.70 |

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|---|---|---|---|---|---|---|
| 1 | A | 7730 | 0 | 7777 | 134 | 0 |
| 2 | B | 2386 | 0 | 2362 | 57 | 0 |
| 3 | G | 255 | 0 | 259 | 6 | 0 |
| 4 | C | 28 | 0 | 25 | 0 | 0 |
| 5 | A | 3 | 0 | 0 | 0 | 0 |
| 6 | A | 28 | 0 | 46 | 2 | 0 |
| 6 | B | 28 | 0 | 46 | 2 | 0 |

*Continued on next page...*

*Continued from previous page...*

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|---|---|---|---|---|---|---|
| 6 | G | 28 | 0 | 46 | 6 | 0 |
| 7 | A | 176 | 0 | 144 | 1 | 0 |
| 8 | A | 57 | 0 | 0 | 0 | 0 |
| 9 | B | 14 | 0 | 13 | 0 | 0 |
| 10 | A | 7 | 0 | 0 | 1 | 0 |
| All | All | 10740 | 0 | 10718 | 194 | 0 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 194 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|---|---|---|---|
| 1:A:864:VAL:HG22 | 2:B:57:MET:HG3 | 1.59 | 0.83 |
| 1:A:807:THR:HB | 1:A:954:GLU:HG3 | 1.64 | 0.78 |
| 2:B:80:ILE:HG12 | 2:B:177:ILE:HG12 | 1.63 | 0.77 |
| 1:A:399:GLN:HE21 | 1:A:436:LEU:HD11 | 1.50 | 0.76 |
| 2:B:113:LYS:HA | 2:B:153:LEU:HD11 | 1.72 | 0.71 |

There are no symmetry-related clashes.

## 5.3 Torsion angles (i)

### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles |
|---|---|---|---|---|---|---|
| 1 | A | 993/1016 (98%) | 931 (94%) | 60 (6%) | 2 (0%) | 47 79 |
| 2 | B | 289/303 (95%) | 257 (89%) | 32 (11%) | 0 | 100 100 |
| 3 | G | 30/65 (46%) | 28 (93%) | 2 (7%) | 0 | 100 100 |
| All | All | 1312/1384 (95%) | 1216 (93%) | 94 (7%) | 2 (0%) | 47 79 |

All (2) Ramachandran outliers are listed below:

| Mol | Chain | Res | Type |
|---|---|---|---|
| 1 | A | 116 | GLU |
| 1 | A | 489 | PRO |

### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|---|---|---|---|---|---|---|
| 1 | A | 846/861 (98%) | 804 (95%) | 42 (5%) | 24 | 53 |
| 2 | B | 261/269 (97%) | 242 (93%) | 19 (7%) | 14 | 43 |
| 3 | G | 26/52 (50%) | 24 (92%) | 2 (8%) | 13 | 42 |
| All | All | 1133/1182 (96%) | 1070 (94%) | 63 (6%) | 21 | 51 |

5 of 63 residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|---|---|---|---|
| 1 | A | 537 | GLU |
| 2 | B | 173 | LYS |
| 1 | A | 763 | ASP |
| 2 | B | 165 | GLU |
| 2 | B | 232 | LEU |

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

| Mol | Chain | Res | Type |
|---|---|---|---|
| 1 | A | 849 | GLN |
| 1 | A | 898 | GLN |
| 2 | B | 193 | ASN |
| 2 | B | 82 | GLN |
| 1 | A | 422 | ASN |

### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | | | Counts | RMSZ | $\#\|Z\| > 2$ | Counts | RMSZ | $\#\|Z\| > 2$ |
| 1 | BFD | A | 369 | 5,1 | 8,11,12 | 0.97 | 0 | 3,15,17 | 1.51 | 0 |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|---|---|---|---|---|---|---|---|
| 1 | BFD | A | 369 | 5,1 | - | 0/5/11/13 | - |

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

| Mol | Chain | Res | Type | Clashes | Symm-Clashes |
|---|---|---|---|---|---|
| 1 | A | 369 | BFD | 1 | 0 |

## 5.5 Carbohydrates (i)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | | | Counts | RMSZ | $\#\|Z\| > 2$ | Counts | RMSZ | $\#\|Z\| > 2$ |
| 4 | NAG | C | 1 | 2,4 | 14,14,15 | 0.55 | 0 | 17,19,21 | 1.40 | 3 (17%) |
| 4 | NAG | C | 2 | 4 | 14,14,15 | 0.82 | 1 (7%) | 17,19,21 | 0.45 | 0 |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|---|---|---|---|---|---|---|---|
| 4 | NAG | C | 1 | 2,4 | - | 3/6/23/26 | 0/1/1/1 |
| 4 | NAG | C | 2 | 4 | - | 0/6/23/26 | 0/1/1/1 |

All (1) bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|---|---|---|---|---|---|---|---|
| 4 | C | 2 | NAG | C1-C2 | 2.29 | 1.55 | 1.52 |

All (3) bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed($^o$) | Ideal($^o$) |
|---|---|---|---|---|---|---|---|
| 4 | C | 1 | NAG | C1-O5-C5 | 3.17 | 116.49 | 112.19 |
| 4 | C | 1 | NAG | O3-C3-C4 | 2.21 | 115.45 | 110.35 |
| 4 | C | 1 | NAG | O4-C4-C3 | 2.07 | 115.13 | 110.35 |

There are no chirality outliers.

All (3) torsion outliers are listed below:

| Mol | Chain | Res | Type | Atoms |
|---|---|---|---|---|
| 4 | C | 1 | NAG | C4-C5-C6-O6 |
| 4 | C | 1 | NAG | O5-C5-C6-O6 |
| 4 | C | 1 | NAG | C3-C2-N2-C7 |

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

**Oligosaccharide Chain C**




## 5.6 Ligand geometry (i)

Of 16 ligands modelled in this entry, 3 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | | | Counts | RMSZ | $\#\|Z\| > 2$ | Counts | RMSZ | $\#\|Z\| > 2$ |
| 7 | PCW | A | 1106 | - | 21,21,53 | 0.84 | 0 | 27,29,61 | 1.16 | 3 (11%) |
| 7 | PCW | A | 1111 | - | 21,21,53 | 0.85 | 0 | 27,29,61 | 0.90 | 1 (3%) |
| 8 | H0C | A | 1121 | - | 63,65,65 | 1.13 | 4 (6%) | 93,104,104 | 1.25 | 6 (6%) |

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | | | Counts | RMSZ | $\#\|Z\| > 2$ | Counts | RMSZ | $\#\|Z\| > 2$ |
| 9 | NAG | B | 421 | 2 | 14,14,15 | 0.44 | 0 | 17,19,21 | 0.54 | 0 |
| 6 | CLR | A | 1104 | - | 31,31,31 | 1.48 | 9 (29%) | 48,48,48 | 1.44 | 12 (25%) |
| 7 | PCW | A | 1105 | - | 21,21,53 | 0.84 | 0 | 27,29,61 | 0.90 | 1 (3%) |
| 7 | PCW | A | 1108 | - | 21,21,53 | 0.84 | 0 | 27,29,61 | 1.11 | 3 (11%) |
| 7 | PCW | A | 1110 | - | 21,21,53 | 0.84 | 0 | 27,29,61 | 1.13 | 3 (11%) |
| 6 | CLR | G | 101 | - | 31,31,31 | 1.44 | 4 (12%) | 48,48,48 | 1.37 | 6 (12%) |
| 7 | PCW | A | 1107 | - | 21,21,53 | 0.85 | 0 | 27,29,61 | 1.25 | 3 (11%) |
| 7 | PCW | A | 1112 | - | 21,21,53 | 0.85 | 0 | 27,29,61 | 0.88 | 1 (3%) |
| 6 | CLR | B | 501 | - | 31,31,31 | 1.54 | 7 (22%) | 48,48,48 | 1.44 | 8 (16%) |
| 7 | PCW | A | 1109 | - | 21,21,53 | 0.86 | 0 | 27,29,61 | 1.25 | 3 (11%) |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|---|---|---|---|---|---|---|---|
| 7 | PCW | A | 1106 | - | - | 13/23/23/57 | - |
| 7 | PCW | A | 1111 | - | - | 5/23/23/57 | - |
| 8 | H0C | A | 1121 | - | - | 5/19/124/124 | 0/9/9/9 |
| 9 | NAG | B | 421 | 2 | - | 1/6/23/26 | 0/1/1/1 |
| 6 | CLR | A | 1104 | - | - | 5/10/68/68 | 0/4/4/4 |
| 7 | PCW | A | 1105 | - | - | 8/23/23/57 | - |
| 7 | PCW | A | 1108 | - | - | 7/23/23/57 | - |
| 7 | PCW | A | 1110 | - | - | 11/23/23/57 | - |
| 6 | CLR | G | 101 | - | - | 4/10/68/68 | 0/4/4/4 |
| 7 | PCW | A | 1107 | - | - | 10/23/23/57 | - |
| 7 | PCW | A | 1112 | - | - | 7/23/23/57 | - |
| 6 | CLR | B | 501 | - | - | 4/10/68/68 | 0/4/4/4 |
| 7 | PCW | A | 1109 | - | - | 10/23/23/57 | - |

The worst 5 of 24 bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|---|---|---|---|---|---|---|---|
| 6 | B | 501 | CLR | C10-C5 | 3.68 | 1.60 | 1.52 |
| 6 | G | 101 | CLR | C10-C5 | 3.54 | 1.59 | 1.52 |
| 6 | A | 1104 | CLR | C10-C5 | 2.89 | 1.58 | 1.52 |

*Continued on next page...*

*Continued from previous page...*

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|---|---|---|---|---|---|---|---|
| 6 | B | 501 | CLR | C10-C9 | 2.85 | 1.60 | 1.56 |
| 8 | A | 1121 | H0C | C26-C27 | 2.84 | 1.58 | 1.54 |

The worst 5 of 50 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed($^o$) | Ideal($^o$) |
|---|---|---|---|---|---|---|---|
| 8 | A | 1121 | H0C | O8-C16-C17 | 5.62 | 116.01 | 108.67 |
| 7 | A | 1109 | PCW | C2-O2-C31 | -4.36 | 109.78 | 117.90 |
| 8 | A | 1121 | H0C | O9-C16-C17 | -4.24 | 122.22 | 130.81 |
| 7 | A | 1106 | PCW | C2-O2-C31 | -3.83 | 110.76 | 117.90 |
| 7 | A | 1107 | PCW | C3-O3-C11 | -3.38 | 108.60 | 117.10 |

There are no chirality outliers.

5 of 90 torsion outliers are listed below:

| Mol | Chain | Res | Type | Atoms |
|---|---|---|---|---|
| 7 | A | 1105 | PCW | C1-O3P-P-O1P |
| 7 | A | 1105 | PCW | C1-O3P-P-O2P |
| 7 | A | 1105 | PCW | C4-O4P-P-O1P |
| 7 | A | 1106 | PCW | C32-C31-O2-C2 |
| 7 | A | 1106 | PCW | O31-C31-O2-C2 |

There are no ring outliers.

4 monomers are involved in 10 short contacts:

| Mol | Chain | Res | Type | Clashes | Symm-Clashes |
|---|---|---|---|---|---|
| 6 | A | 1104 | CLR | 2 | 0 |
| 6 | G | 101 | CLR | 6 | 0 |
| 7 | A | 1107 | PCW | 1 | 0 |
| 6 | B | 501 | CLR | 2 | 0 |

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.

**Ligand PCW A 1106**



**Bond lengths**



**Bond angles**



**Torsions**



**Rings**

**Ligand PCW A 1111**



**Bond lengths**



**Bond angles**



**Torsions**



**Rings**

**Ligand H0C A 1121**




**Ligand CLR A 1104**



**Ligand PCW A 1105**



Bond lengths



Bond angles



Torsions



Rings

**Ligand PCW A 1108**



Bond lengths



Bond angles



Torsions



Rings

## Ligand PCW A 1110



Bond lengths



Bond angles



Torsions



Rings

## Ligand CLR G 101



Bond lengths



Bond angles



Torsions



Rings

**Ligand PCW A 1107**



Bond lengths



Bond angles



Torsions



Rings

**Ligand PCW A 1112**



Bond lengths



Bond angles



Torsions



Rings

**Ligand CLR B 501**




**Ligand PCW A 1109**




## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

# 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

## 6.3 Carbohydrates (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around Chain C:**

$2mF_o$-$DF_c$ (at 0.7 rmsd) in gray
$mF_o$-$DF_c$ (at 3 rmsd) in purple (negative) and green (positive)



## 6.4 Ligands (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

| | |
|---|---|
| **Electron density around CLR A 1104:**<br><br>$2mF_o$-$DF_c$ (at 0.7 rmsd) in gray<br>$mF_o$-$DF_c$ (at 3 rmsd) in purple (negative)<br>and green (positive) |  |
|  |  |

**Electron density around CLR B 501:**

$2mF_o$-$DF_c$ (at 0.7 rmsd) in gray
$mF_o$-$DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



**Electron density around CLR G 101:**

$2mF_o$-$DF_c$ (at 0.7 rmsd) in gray
$mF_o$-$DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



**Electron density around PCW A 1105:**

$2mF_o$-$DF_c$ (at 0.7 rmsd) in gray
$mF_o$-$DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)







**Electron density around PCW A 1106:**

$2mF_o$-$DF_c$ (at 0.7 rmsd) in gray
$mF_o$-$DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)







**Electron density around PCW A 1107:**

$2mF_o$-$DF_c$ (at 0.7 rmsd) in gray
$mF_o$-$DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)







**Electron density around PCW A 1108:**

$2mF_o$-$DF_c$ (at 0.7 rmsd) in gray
$mF_o$-$DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)







**Electron density around PCW A 1109:**

$2mF_o$-$DF_c$ (at 0.7 rmsd) in gray
$mF_o$-$DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)







**Electron density around PCW A 1110:**

$2mF_o$-$DF_c$ (at 0.7 rmsd) in gray
$mF_o$-$DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)







**Electron density around PCW A 1111:**

$2mF_o$-$DF_c$ (at 0.7 rmsd) in gray
$mF_o$-$DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



**Electron density around PCW A 1112:**

$2mF_o$-$DF_c$ (at 0.7 rmsd) in gray
$mF_o$-$DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



**Electron density around H0C A 1121:**

$2mF_o$-$DF_c$ (at 0.7 rmsd) in gray
$mF_o$-$DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



## 6.5 Other polymers (i)

Unable to reproduce the depositors R factor - this section is therefore empty.