



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:37 PM UTC

PDB ID : 9DDD / pdb\_00009ddd  
Title : SufS-SufU complex from Mycobacterium Tuberculosis  
Authors : Stuteley, S.M.; Bashiri, G.  
Deposited on : 2024-08-27  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	417	<div> <div>2%</div> <div>91%</div> <div>6%</div> <div>.</div> </div>
1	B	417	<div> <div>2%</div> <div>93%</div> <div>5%</div> <div>.</div> </div>
2	C	162	<div> <div>7%</div> <div>85%</div> <div>12%</div> <div>.</div> </div>
2	D	162	<div> <div>10%</div> <div>84%</div> <div>13%</div> <div>.</div> </div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8668 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable cysteine desulfurase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	406	Total	C	N	O	P	S	0	2	0
			3071	1928	552	578	1	12			
1	B	408	Total	C	N	O	P	S	0	1	0
			3091	1935	559	583	1	13			


- Molecule 2 is a protein called Possible nitrogen fixation related protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	143	Total	C	N	O	S	0	0	0
			1089	682	192	211	4			
2	D	141	Total	C	N	O	S	0	0	0
			1067	671	184	208	4			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	1	VAL	-	expression tag	UNP O53156
D	1	VAL	-	expression tag	UNP O53156

- Molecule 3 is ALANINE (CCD ID: ALA) (formula: C<sub>3</sub>H<sub>7</sub>NO<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			6	3	1	2		
3	B	1	Total	C	N	O	0	0
			6	3	1	2		

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	1	Total	Zn	0	0
			1	1		
4	D	1	Total	Zn	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	162	Total	O	0	0
			162	162		
5	B	166	Total	O	0	0
			166	166		
5	C	4	Total	O	0	0
			4	4		
5	D	4	Total	O	0	0
			4	4		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Probable cysteine desulfurase




- Molecule 1: Probable cysteine desulfurase



- Molecule 2: Possible nitrogen fixation related protein



- Molecule 2: Possible nitrogen fixation related protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	66.80Å 73.62Å 76.35Å 91.03° 100.22° 110.02°	Depositor
Resolution (Å)	48.68 – 2.00 48.68 – 2.00	Depositor EDS
% Data completeness (in resolution range)	97.9 (48.68-2.00) 97.9 (48.68-2.00)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.14 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.8.0349	Depositor
R, $R_{free}$	0.179 , 0.203 0.188 , 0.210	Depositor DCC
$R_{free}$ test set	4434 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	30.0	Xtriage
Anisotropy	0.011	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 28.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	8668	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.13% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: LLP, ZN, CSS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.58	0/3106	0.82	0/4224
1	B	0.58	0/3122	0.84	1/4243 (0.0%)
2	C	0.54	0/1108	0.84	1/1504 (0.1%)
2	D	0.54	0/1086	0.83	0/1475
All	All	0.57	0/8422	0.83	2/11446 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	1
2	C	0	1
2	D	0	1
All	All	0	6

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	42	ASP	CA-CB-CG	8.57	121.17	112.60
1	B	137	ILE	N-CA-CB	5.80	114.15	110.50

There are no chirality outliers.

All (6) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	117	ARG	Sidechain
1	A	411[A]	ARG	Sidechain
1	A	411[B]	ARG	Sidechain
1	B	411	ARG	Sidechain
2	C	56	ARG	Sidechain
2	D	56	ARG	Sidechain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3071	0	3002	17	0
1	B	3091	0	3032	15	0
2	C	1089	0	1051	2	0
2	D	1067	0	1030	2	0
3	A	6	0	4	1	0
3	B	6	0	4	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	A	162	0	0	0	0
5	B	166	0	0	0	0
5	C	4	0	0	0	0
5	D	4	0	0	0	0
All	All	8668	0	8123	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:62:ARG:O	1:B:68[A]:MET:SD	2.49	0.70
2:C:77:LEU:HD21	2:C:139:PHE:HB2	1.75	0.68
2:D:77:LEU:HD21	2:D:139:PHE:HB2	1.81	0.63
1:B:65:HIS:NE2	1:B:68[A]:MET:HE3	2.16	0.60
1:A:313:ARG:NH2	1:A:337:MET:O	2.39	0.56
1:A:65:HIS:NE2	1:A:68:MET:HE3	2.21	0.55

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:196:ARG:HA	1:A:199:GLN:HE21	1.71	0.55
1:A:129:GLU:HB3	1:A:155:VAL:HG23	1.92	0.51
1:A:207:ASP:OD2	1:A:233:LLP:N1	2.45	0.49
1:B:207:ASP:OD2	1:B:233:LLP:N1	2.48	0.47
1:B:360:LEU:HD23	1:B:406:GLY:HA3	1.97	0.47
1:A:34:ASP:OD2	1:B:68[A]:MET:HE3	2.15	0.46
3:A:501:ALA:HB3	1:B:62:ARG:HH22	1.80	0.46
1:B:107:SER:O	1:B:142:LEU:HD13	2.16	0.45
1:A:368:ARG:NH2	1:B:62:ARG:HG2	2.31	0.45
1:A:65:HIS:CE1	1:A:68:MET:HG3	2.52	0.44
1:A:105:LEU:C	1:A:105:LEU:HD23	2.43	0.44
2:C:99:MET:HE1	2:C:116:GLY:O	2.17	0.43
1:B:212:VAL:N	1:B:213:PRO:CD	2.82	0.43
1:A:349:GLU:O	1:A:411[B]:ARG:NH2	2.51	0.43
1:B:65:HIS:CE1	1:B:68[A]:MET:HG3	2.53	0.43
1:A:155:VAL:HG12	1:A:156:THR:O	2.17	0.43
1:B:343:PRO:HB3	1:B:390:SER:HB3	2.01	0.43
1:B:105:LEU:C	1:B:105:LEU:HD23	2.44	0.42
1:B:63:GLY:C	1:B:68[A]:MET:SD	3.03	0.42
1:A:49:GLU:HG3	1:A:288:SER:HB2	2.02	0.42
1:A:210:GLN:HE21	1:A:233:LLP:HO3	1.66	0.42
1:A:107:SER:O	1:A:142:LEU:HD13	2.20	0.41
1:A:110:LEU:HD13	1:A:174:VAL:HG21	2.02	0.41
1:A:212:VAL:N	1:A:213:PRO:CD	2.84	0.41
1:A:68:MET:HE3	1:B:34:ASP:OD2	2.21	0.41
1:B:286:MET:O	1:B:286:MET:HG2	2.21	0.41
2:D:102:SER:O	2:D:103:ARG:HB2	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	404/417 (97%)	393 (97%)	11 (3%)	0	100	100
1	B	405/417 (97%)	394 (97%)	11 (3%)	0	100	100
2	C	139/162 (86%)	139 (100%)	0	0	100	100
2	D	137/162 (85%)	137 (100%)	0	0	100	100
All	All	1085/1158 (94%)	1063 (98%)	22 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	307/319 (96%)	306 (100%)	1 (0%)	86	91
1	B	312/319 (98%)	309 (99%)	3 (1%)	68	75
2	C	113/132 (86%)	113 (100%)	0	100	100
2	D	111/132 (84%)	111 (100%)	0	100	100
All	All	843/902 (94%)	839 (100%)	4 (0%)	81	87

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	128	THR
1	B	128	THR
1	B	200	SER
1	B	248	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	199	GLN
1	A	210	GLN
1	A	278	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	199	GLN
2	C	80	GLN
2	C	84	GLN
2	C	145	GLN
2	D	80	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	LLP	A	233	1	23,24,25	0.51	0	25,32,34	0.65	0
1	CSS	A	373	1	4,6,7	0.58	0	2,6,8	0.95	0
1	CSS	B	373	1	4,6,7	0.54	0	2,6,8	0.73	0
1	LLP	B	233	1	23,24,25	0.46	0	25,32,34	0.66	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	A	233	1	-	1/16/17/19	0/1/1/1
1	CSS	A	373	1	-	0/1/5/7	-
1	CSS	B	373	1	-	0/1/5/7	-
1	LLP	B	233	1	-	3/16/17/19	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	233	LLP	C4-C4'-NZ-CE
1	A	233	LLP	C4-C4'-NZ-CE
1	B	233	LLP	CG-CD-CE-NZ
1	B	233	LLP	C3-C4-C4'-NZ

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	233	LLP	2	0
1	B	233	LLP	1	0

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	ALA	B	501	-	5,5,5	0.98	0	6,6,6	0.89	0
3	ALA	A	501	-	5,5,5	1.05	0	6,6,6	0.85	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ALA	B	501	-	-	2/4/4/4	-
3	ALA	A	501	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	501	ALA	O-C-CA-N
3	A	501	ALA	OXT-C-CA-N
3	B	501	ALA	O-C-CA-N
3	B	501	ALA	OXT-C-CA-N


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	501	ALA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	404/417 (96%)	-0.05	8 (1%) 65 64	17, 29, 42, 68	2 (0%)
1	B	406/417 (97%)	-0.06	10 (2%) 58 58	16, 28, 45, 65	1 (0%)
2	C	143/162 (88%)	1.02	11 (7%) 19 18	33, 48, 67, 78	0
2	D	141/162 (87%)	1.08	17 (12%) 8 8	35, 50, 67, 74	0
All	All	1094/1158 (94%)	0.23	46 (4%) 40 39	16, 31, 60, 78	3 (0%)

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	3	LEU	5.6
2	D	5	LEU	4.5
1	B	63	GLY	4.3
2	C	150	PHE	4.3
2	C	36	VAL	4.2
1	A	62	ARG	4.0
2	C	6	GLU	4.0
2	D	36	VAL	3.9
1	A	167[A]	TYR	3.7
1	A	63	GLY	3.7
2	D	34	TYR	3.6
2	D	150	PHE	3.5
2	D	149	ALA	3.4
1	A	412	HIS	3.2
1	B	384	ALA	3.1
1	B	415	GLY	3.0
1	B	62	ARG	3.0
2	C	18	LYS	2.8
2	C	17	TYR	2.8
2	D	19	HIS	2.7
2	D	12	VAL	2.7

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	167	TYR	2.7
2	D	106	VAL	2.6
2	D	25	LEU	2.6
1	A	384	ALA	2.6
2	D	33	VAL	2.5
1	B	412	HIS	2.5
2	C	149	ALA	2.4
2	D	17	TYR	2.3
1	B	64	ALA	2.3
2	D	24	GLY	2.3
2	D	107	PRO	2.3
1	B	8	LEU	2.3
2	D	90	LEU	2.3
2	C	106	VAL	2.2
2	C	19	HIS	2.2
2	C	34	TYR	2.2
2	D	144	ALA	2.1
2	D	124	LYS	2.1
1	A	64	ALA	2.1
1	A	264	GLU	2.1
2	D	65	GLN	2.1
2	C	4	ARG	2.1
1	B	413	PHE	2.0
1	B	362	ASP	2.0
1	A	350	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
1	CSS	B	373	7/8	0.95	0.07	22,23,30,36	0
1	CSS	A	373	7/8	0.96	0.07	23,24,29,35	0
1	LLP	B	233	24/25	0.97	0.07	19,26,29,32	0
1	LLP	A	233	24/25	0.97	0.07	21,25,28,30	0

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	ALA	B	501	6/6	0.87	0.15	35,44,47,48	0
3	ALA	A	501	6/6	0.88	0.15	33,42,46,47	0
4	ZN	C	201	1/1	0.98	0.03	33,33,33,33	0
4	ZN	D	201	1/1	0.98	0.04	37,37,37,37	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around ALA B 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around ALA A 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around ZN C 201:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





**Electron density around ZN D 201:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**6.5 Other polymers** [i](#)

There are no such residues in this entry.