Full wwPDB X-ray Structure Validation Report (i)

Aug 22, 2023 – 04:58 PM EDT

PDB ID	:	3DHD
Title	:	Crystal structure of human NAMPT complexed with nicotinamide mononucleotide and pyrophosphate
Authors	:	Ho, M.; Burgos, E.S.; Almo, S.C.; Schramm, V.L.
Deposited on	:	2008-06-17
Resolution	:	2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	484	
1	B	484	

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 7890 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nicotinamide phosphoribosyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	464	3733	2404	615	707	7	0	4	0
1	B	464	3732	2403	614	708	7	0	4	0

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
2	A	2	2	2	0	0
2	B	2	2	2	0	0

- Molecule 3 is BETA-NICOTINAMIDE RIBOSE MONOPHOSPHATE (three-letter code: NMN) (formula: $C_{11}H_{16}N_2O_8P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	22	11	2	8	1	0	0
3	B	1	22	11	2	8	1	0	0

- Molecule 4 is PYROPHOSPHATE 2- (three-letter code: POP) (formula: $H_2O_7P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	O	P				
4	A	1	9	7	2			0	0
4	B	1	9	7	2			0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	180	180	180	0	0
5	B	179	179	179	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Nicotinamide phosphoribosyltransferase

Chain A:



• Molecule 1: Nicotinamide phosphoribosyltransferase

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	61.17Å 107.07Å 83.34Å 90.00° 97.02° 90.00°	Depositor
Resolution (Å)	19.89 – 2.00 19.89 – 2.00	Depositor EDS
% Data completeness (in resolution range)	96.8 (19.89-2.00) 96.7 (19.89-2.00)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.72 (at 2.01Å)	Xtriage
Refinement program	REFMAC 5.4.0069	Depositor
R, R_{free}	0.180 , 0.221 0.179 , 0.220	Depositor DCC
R_{free} test set	3460 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å^2)	23.5	Xtriage
Anisotropy	0.055	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.37 , 42.3	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	7890	wwPDB-VP
Average B, all atoms (Å^2)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.36% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: POP, MG, NMN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.86	2/3834 (0.1%)	0.78	1/5197 (0.0%)
1	B	0.88	0/3832	0.77	1/5194 (0.0%)
All	All	0.87	2/7666 (0.0%)	0.78	2/10391 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	82	GLU	CG-CD	5.90	1.60	1.51
1	A	60	TYR	CD1-CE1	-5.44	1.31	1.39

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	95	VAL	CB-CA-C	-6.22	99.58	111.40
1	A	318	LEU	CA-CB-CG	5.39	127.69	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3733	0	3712	33	0
1	B	3732	0	3713	23	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	22	0	13	0	0
3	B	22	0	13	0	0
4	A	9	0	0	0	0
4	B	9	0	0	0	0
5	A	180	0	0	2	0
5	B	179	0	0	0	0
All	All	7890	0	7451	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:63:GLN:HE22	1:A:471:TYR:H	1.21	0.88
1:A:316:ASN:HD22	1:A:319:ASP:H	1.31	0.78
1:A:245:ALA:HA	1:B:25:GLN:HE22	1.55	0.71
1:A:88:LYS:HD2	1:A:88:LYS:O	1.98	0.63
1:A:32:LYS:HA	1:A:136:ASN:HD21	1.64	0.62
1:A:431:SER:OG	1:A:433:HIS:HE1	1.82	0.62
1:A:136:ASN:HD22	1:A:136:ASN:H	1.46	0.61
1:B:21:THR:HG22	1:B:95:VAL:HG11	1.83	0.61
1:B:420:ASP:OD1	1:B:422[A]:ASN:HB2	2.02	0.60
1:B:318:LEU:HD13	1:B:364:ILE:HA	1.85	0.58
1:A:321:VAL:HG23	1:A:352:GLN:HE21	1.69	0.58
1:A:183:LEU:HD12	1:A:484:ILE:HD12	1.85	0.57
1:B:224:LEU:HD22	1:B:238:PRO:HD2	1.85	0.57
1:A:40:ARG:HD3	1:A:422:ASN:O	2.04	0.57
1:A:113:PRO:HA	1:A:464:ASN:HD22	1.70	0.55
1:B:404[A]:VAL:CG1	1:B:411:ILE:HD11	2.37	0.55
1:A:40:ARG:NH1	1:A:396:ASN:ND2	2.56	0.54
1:B:420:ASP:OD1	1:B:422[B]:ASN:HB3	2.05	0.54
1:B:107:LYS:HG2	1:B:108:TYR:CE1	2.42	0.54
1:B:178:GLU:OE2	1:B:369:LYS:NZ	2.28	0.53
1:A:149:GLU:O	1:A:153:VAL:HG23	2.10	0.51
1:B:411:ILE:HD12	1:B:413:VAL:CG1	2.41	0.51
1:A:107:LYS:HG2	1:A:108:TYR:CE1	2.47	0.49
1:B:40:ARG:HH21	1:B:398:SER:HB2	1.78	0.48
1:A:318:LEU:HD21	1:A:367:GLY:HA3	1.95	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:31:SER:HB2	1:A:139:PRO:HB3	1.95	0.47
1:A:201:GLN:NE2	1:B:151:ILE:HD12	2.30	0.47
1:A:183:LEU:HD12	1:A:484:ILE:CD1	2.45	0.47
1:A:393:ASP:CG	5:A:569:HOH:O	2.53	0.46
1:B:291:TRP:CE3	1:B:299:ILE:HD11	2.50	0.46
1:B:484:ILE:O	1:B:484:ILE:HG13	2.15	0.45
1:A:136:ASN:H	1:A:136:ASN:ND2	2.12	0.45
1:A:259:LYS:HD2	1:A:294:ASP:HB3	1.96	0.45
1:B:113:PRO:HD2	1:B:144:LEU:CD2	2.46	0.45
1:A:482:LEU:O	1:A:484:ILE:HG22	2.17	0.45
1:B:404[A]:VAL:HG12	1:B:411:ILE:HG13	1.99	0.45
1:A:175:TYR:HB3	1:A:375:ILE:HG13	1.99	0.45
1:B:113:PRO:HA	1:B:464:ASN:HD22	1.82	0.44
1:A:169:LYS:HB2	1:A:215:PHE:CZ	2.53	0.44
1:A:361:LEU:C	1:A:361:LEU:HD23	2.38	0.44
1:B:32:LYS:O	1:B:404[A]:VAL:HG23	2.18	0.43
1:B:27:PRO:HA	1:B:28:PRO:HD3	1.89	0.43
1:A:91:PHE:O	1:A:92:GLN:HB2	2.18	0.43
1:A:178[A]:GLU:HG3	5:A:537:HOH:O	2.19	0.43
1:A:420:ASP:OD1	1:A:422:ASN:HB2	2.19	0.43
1:B:55:GLU:H	1:B:55:GLU:HG2	1.66	0.43
1:A:40:ARG:HH21	1:A:398:SER:CB	2.32	0.43
1:A:478:LYS:HA	1:A:481:GLN:HE21	1.84	0.43
1:B:375:ILE:HG22	1:B:375:ILE:O	2.19	0.42
1:A:120:PRO:O	1:A:123:PHE:HB2	2.20	0.42
1:B:291:TRP:HE3	1:B:299:ILE:HD11	1.85	0.41
1:A:40:ARG:NH1	1:A:396:ASN:HD22	2.18	0.40
1:A:430:LEU:HD23	1:A:444:GLU:HA	2.03	0.40
1:B:40:ARG:NH1	1:B:423:LYS:HG2	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	464/484 (96%)	449 (97%)	15 (3%)	0	100	100
1	B	464/484 (96%)	453 (98%)	11 (2%)	0	100	100
All	All	928/968 (96%)	902 (97%)	26 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	412/425 (97%)	400 (97%)	12 (3%)	42	43
1	B	412/425 (97%)	398 (97%)	14 (3%)	37	36
All	All	824/850 (97%)	798 (97%)	26 (3%)	39	38

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	TYR
1	A	32	LYS
1	A	41	GLU
1	A	102	ASN
1	A	107	LYS
1	A	136	ASN
1	A	151	ILE
1	A	195	TYR
1	A	318	LEU
1	A	396	ASN
1	A	409	LEU
1	A	434	ARG
1	B	18	TYR
1	B	55	GLU
1	B	94	ASP
1	B	107	LYS
1	B	151	ILE
1	B	183	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	195	TYR
1	B	303	SER
1	B	305	GLN
1	B	370	GLN
1	B	409	LEU
1	B	434	ARG
1	B	469	LYS
1	B	484	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	63	GLN
1	A	136	ASN
1	A	164	ASN
1	A	305	GLN
1	A	316	ASN
1	A	362	GLN
1	A	396	ASN
1	A	433	HIS
1	A	455	GLN
1	A	464	ASN
1	A	481	GLN
1	B	25	GLN
1	B	164	ASN
1	B	362	GLN
1	B	396	ASN
1	B	464	ASN
1	B	481	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	NMN	B	503	2	22,23,23	0.88	1 (4%)	30,34,34	1.47	6 (20%)
3	NMN	A	503	2	22,23,23	1.02	1 (4%)	30,34,34	1.94	8 (26%)
4	POP	A	504	2	6,8,8	0.67	0	13,13,13	1.82	4 (30%)
4	POP	B	504	2	6,8,8	0.78	0	13,13,13	1.46	3 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NMN	B	503	2	-	6/14/30/30	0/2/2/2
3	NMN	A	503	2	-	4/14/30/30	0/2/2/2
4	POP	A	504	2	-	0/6/6/6	-
4	POP	B	504	2	-	0/6/6/6	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	503	NMN	O4R-C1R	2.72	1.44	1.41
3	B	503	NMN	O2R-C2R	2.19	1.48	1.43

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	A	503	NMN	O7-C7-N7	-5.02	115.45	122.58
3	B	503	NMN	C3R-C2R-C1R	3.89	106.83	100.98
3	A	503	NMN	C3R-C2R-C1R	3.86	106.79	100.98
4	A	504	POP	P2-O-P1	-3.75	119.95	132.83
3	A	503	NMN	C3-C7-N7	3.64	122.12	117.75
3	A	503	NMN	O3R-C3R-C4R	-3.58	100.68	111.05
4	A	504	POP	O6-P2-O	2.93	114.47	104.64
4	A	504	POP	O3-P1-O	2.77	113.91	104.64
4	B	504	POP	O2-P1-O	2.75	113.87	104.64
4	B	504	POP	P2-O-P1	-2.73	123.46	132.83
3	B	503	NMN	C6-N1-C2	-2.60	119.60	121.97
3	A	503	NMN	C3-C2-N1	2.59	122.96	120.43
3	A	503	NMN	O2P-P-O5R	-2.54	99.97	106.73
3	B	503	NMN	C3-C2-N1	2.52	122.89	120.43
4	B	504	POP	O5-P2-O	2.42	112.77	104.64
3	A	503	NMN	O7-C7-C3	2.32	122.41	119.63
3	B	503	NMN	C5R-C4R-C3R	-2.29	106.60	115.18
3	A	503	NMN	C5R-C4R-C3R	-2.22	106.87	115.18
4	A	504	POP	O2-P1-O	2.14	111.81	104.64
3	B	503	NMN	O1P-P-O5R	-2.12	101.10	106.73
3	B	503	NMN	O2P-P-O5R	2.01	112.08	106.73

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	503	NMN	C5R-O5R-P-O1P
3	A	503	NMN	C5R-O5R-P-O2P
3	A	503	NMN	C2R-C1R-N1-C2
3	A	503	NMN	C2R-C1R-N1-C6
3	B	503	NMN	C5R-O5R-P-O1P
3	B	503	NMN	C5R-O5R-P-O2P
3	B	503	NMN	C2R-C1R-N1-C2
3	B	503	NMN	C2R-C1R-N1-C6
3	B	503	NMN	C5R-O5R-P-O3P
3	B	503	NMN	C3R-C4R-C5R-O5R

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand NMN B 503



Ligand NMN A 503



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	464/484 (95%)	-0.27	6 (1%)	77	76	13, 21, 32, 48	0
1	B	464/484 (95%)	-0.28	4 (0%)	84	83	13, 21, 33, 46	0
All	All	928/968 (95%)	-0.28	10 (1%)	80	79	13, 21, 33, 48	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	484	ILE	4.7
1	B	484	ILE	4.6
1	A	409	LEU	3.4
1	A	483	ASN	3.2
1	B	483	ASN	2.8
1	B	409	LEU	2.7
1	B	451	GLU	2.6
1	A	450	LEU	2.4
1	A	234	LYS	2.2
1	A	181	GLY	2.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	MG	B	502	1/1	0.93	0.07	29,29,29,29	0
3	NMN	B	503	22/22	0.95	0.09	14,23,28,28	0
2	MG	A	502	1/1	0.96	0.06	31,31,31,31	0
3	NMN	A	503	22/22	0.96	0.09	17,23,27,29	0
2	MG	A	501	1/1	0.96	0.05	30,30,30,30	0
4	POP	A	504	9/9	0.96	0.11	22,27,29,30	0
2	MG	B	501	1/1	0.97	0.03	30,30,30,30	0
4	POP	B	504	9/9	0.97	0.08	21,25,28,29	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around NMN B 503:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)





Electron density around NMN A 503:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.