



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 23, 2023 – 04:09 AM EDT

PDB ID : 3E0I  
Title : Cu<sup>2+</sup> substituted Aquifex aeolicus KDO8PS in complex with PEP  
Authors : Gatti, D.L.  
Deposited on : 2008-07-31  
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	267	 4% 88% 9% ..
1	B	267	 3% 81% 13% . .

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4819 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 2-dehydro-3-deoxyphosphooctonate aldolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	263	Total 2128	C 1380	N 350	O 392	S 6	0	10	0
1	B	255	Total 2078	C 1356	N 339	O 376	S 7	0	12	0

- Molecule 2 is COPPER (II) ION (three-letter code: CU) (formula: Cu).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Cu		
2	A	1	Total 1	Cu 1	0	0
2	B	1	Total 1	Cu 1	0	0

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O P 5 4 1	0	0
3	B	1	Total O P 5 4 1	0	0

- Molecule 4 is PHOSPHOENOLPYRUVATE (three-letter code: PEP) (formula:  $C_3H_5O_6P$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O P 10 3 6 1	0	0
4	B	1	Total C O P 10 3 6 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	307	Total O 307 307	0	0
5	B	274	Total O 274 274	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 2-dehydro-3-deoxyphosphooctonate aldolase



- Molecule 1: 2-dehydro-3-deoxyphosphooctonate aldolase



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	84.79Å 84.79Å 159.90Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	21.20 – 1.70 21.20 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.0 (21.20-1.70) 99.1 (21.20-1.70)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.81 (at 1.70Å)	Xtrriage
Refinement program	REFMAC 5.4	Depositor
R, $R_{free}$	0.135 , 0.166 0.133 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.6	Xtrriage
Anisotropy	0.130	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 54.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.024 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	4819	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.98% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, CU, PEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.03	0/2195	0.89	6/2961 (0.2%)
1	B	1.02	0/2153	0.93	6/2907 (0.2%)
All	All	1.02	0/4348	0.91	12/5868 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	B	2106	ARG	NE-CZ-NH2	-12.95	113.82	120.30
1	B	2106	ARG	NE-CZ-NH1	11.72	126.16	120.30
1	A	1201	ARG	NE-CZ-NH1	-7.94	116.33	120.30
1	A	1133	ASP	CB-CG-OD1	7.62	125.16	118.30
1	B	2201	ARG	NE-CZ-NH1	-6.61	117.00	120.30
1	A	1201	ARG	NE-CZ-NH2	5.79	123.19	120.30
1	B	2081	ASP	CB-CG-OD1	5.75	123.47	118.30
1	A	1081	ASP	CB-CG-OD2	-5.64	113.22	118.30
1	A	1208	ILE	CG1-CB-CG2	-5.36	99.61	111.40
1	A	1133	ASP	CB-CG-OD2	-5.33	113.51	118.30
1	B	2106	ARG	CD-NE-CZ	5.27	130.98	123.60
1	B	2029	LEU	CB-CG-CD2	-5.06	102.40	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1002	GLU	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2128	0	2186	24	0
1	B	2078	0	2155	47	1
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
4	A	10	0	2	0	0
4	B	10	0	2	0	0
5	A	307	0	0	9	2
5	B	274	0	0	7	2
All	All	4819	0	4345	69	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:2221[B]:MET:SD	1:B:2238[B]:LEU:CD1	2.02	1.47
1:B:2221[B]:MET:SD	1:B:2238[B]:LEU:HD11	1.62	1.34
1:B:2208[A]:ILE:HD11	1:B:2219[A]:VAL:HG11	1.22	1.12
1:B:2221[B]:MET:CG	1:B:2238[B]:LEU:HD12	1.80	1.10
1:B:2208[A]:ILE:CD1	1:B:2219[A]:VAL:HG11	1.84	1.08
1:A:1201:ARG:HD3	5:A:5342:HOH:O	1.53	1.08
1:B:2221[B]:MET:SD	1:B:2238[B]:LEU:HD12	1.84	1.04
1:A:1174[B]:LYS:NZ	1:A:1217:ASP:OD2	2.02	0.92
1:B:2201:ARG:HD3	5:B:5414:HOH:O	1.74	0.87
1:B:2188[B]:GLN:NE2	1:B:2236:THR:OG1	2.07	0.86
1:B:2178:LYS:HE2	5:B:5358:HOH:O	1.76	0.86

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1002:GLU:O	1:A:1002:GLU:HG2	1.76	0.85
1:B:2069:LYS:HE3	1:B:2073:GLU:OE1	1.84	0.76
1:A:1193:LEU:HB2	1:A:1196[B]:LYS:HG3	1.68	0.75
1:B:2208[B]:ILE:CG2	1:B:2250[B]:ILE:CG2	2.64	0.75
1:B:2208[B]:ILE:CG2	1:B:2250[B]:ILE:HG21	2.18	0.74
1:B:2221[B]:MET:HG2	1:B:2238[B]:LEU:HD12	1.69	0.74
1:A:1189:LEU:HD11	1:A:1200:MET:HE2	1.70	0.73
1:B:2133:ASP:OD1	5:B:5453:HOH:O	2.07	0.71
1:A:1238[A]:LEU:HD21	1:A:1246:ILE:HD12	1.82	0.62
1:A:1196[A]:LYS:HE3	5:A:5420:HOH:O	2.01	0.60
1:A:1048:ASN:H	1:A:1048:ASN:HD22	1.47	0.60
1:B:2208[A]:ILE:HD12	1:B:2219[A]:VAL:HG11	1.81	0.60
1:A:1212[A]:VAL:HG11	1:A:1250:ILE:HG23	1.84	0.59
1:B:2014:GLU:O	1:B:2058:HIS:HE1	1.85	0.59
1:B:2184:THR:HB	5:B:5423:HOH:O	2.03	0.58
1:B:2208[B]:ILE:HG22	1:B:2250[B]:ILE:HG22	1.85	0.58
1:B:2208[A]:ILE:HD11	1:B:2219[A]:VAL:CG1	2.15	0.58
1:B:2208[B]:ILE:HG22	1:B:2250[B]:ILE:CG2	2.34	0.58
1:A:1201:ARG:HG3	1:A:1204:ILE:HD12	1.87	0.57
5:A:5306:HOH:O	1:B:2053:HIS:HE1	1.88	0.57
1:B:2055:PHE:CZ	1:B:2057:GLY:HA2	2.40	0.57
1:B:2048:ASN:H	1:B:2048:ASN:HD22	1.52	0.56
1:B:2188[B]:GLN:HE22	1:B:2233:ASP:CG	2.10	0.55
1:A:1119:ARG:HD3	5:A:5273:HOH:O	2.07	0.54
1:A:1002:GLU:N	5:A:5655:HOH:O	2.40	0.53
1:B:2188[B]:GLN:HG2	1:B:2236:THR:HG21	1.90	0.53
1:A:1174[B]:LYS:NZ	1:A:1217:ASP:CG	2.64	0.51
1:A:1204:ILE:O	1:A:1208:ILE:HG13	2.10	0.51
1:B:2212[B]:VAL:HG21	1:B:2250[B]:ILE:HB	1.92	0.51
1:B:2208[B]:ILE:HG21	1:B:2250[B]:ILE:HG21	1.92	0.51
1:A:1183:ALA:HB2	1:A:1208:ILE:HG12	1.92	0.50
1:B:2093:GLU:HG3	5:B:5132:HOH:O	2.12	0.49
5:A:5194:HOH:O	1:B:2083:HIS:HD2	1.97	0.48
1:B:2064:VAL:HG13	1:B:2094:VAL:HG21	1.95	0.48
1:A:1204:ILE:HG21	1:A:1238[B]:LEU:HD21	1.96	0.47
1:B:2201:ARG:HG3	1:B:2204:ILE:HD12	1.96	0.47
1:A:1183:ALA:CB	1:A:1208:ILE:HG12	2.45	0.47
1:B:2208[B]:ILE:HG21	1:B:2250[B]:ILE:CG2	2.41	0.47
1:A:1020:LEU:HD21	1:A:1069:LYS:HG3	1.99	0.45
1:B:2064:VAL:HG13	1:B:2094:VAL:CG2	2.46	0.45
1:B:2228:GLU:HG2	1:B:2229:LYS:HG3	1.98	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:2188[B]:GLN:HG2	1:B:2236:THR:CG2	2.48	0.44
1:A:1136:ASN:HB3	1:B:2050[A]:SER:O	2.17	0.44
1:B:2184:THR:CB	5:B:5423:HOH:O	2.64	0.44
1:A:1055:PHE:CZ	1:A:1057:GLY:HA2	2.53	0.43
1:A:1195:ASP:HA	1:B:2136:ASN:OD1	2.18	0.43
1:B:2247:ILE:O	1:B:2250[B]:ILE:HG13	2.19	0.43
1:B:2188[B]:GLN:CG	1:B:2236:THR:HG21	2.49	0.43
1:B:2048:ASN:HD22	1:B:2048:ASN:N	2.13	0.42
1:A:1002:GLU:HG3	5:A:5572:HOH:O	2.18	0.42
1:B:2077:LYS:HE3	5:B:5359:HOH:O	2.18	0.42
1:B:2205:PHE:O	1:B:2208[A]:ILE:HG22	2.21	0.41
1:A:1200:MET:HG2	5:A:5402:HOH:O	2.20	0.41
5:A:5194:HOH:O	1:B:2083:HIS:CD2	2.72	0.41
1:B:2188[B]:GLN:HE21	1:B:2236:THR:HG1	1.52	0.40
1:A:1238[A]:LEU:HD11	1:A:1246:ILE:CD1	2.52	0.40
1:B:2014:GLU:O	1:B:2058:HIS:CE1	2.70	0.40
1:B:2020:LEU:HD21	1:B:2069:LYS:HG3	2.04	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:2035:GLU:OE1	5:A:5220:HOH:O[3_664]	2.05	0.15
5:A:5278:HOH:O	5:B:5415:HOH:O[2_655]	2.12	0.08
5:B:5408:HOH:O	5:B:5408:HOH:O[4_555]	2.16	0.04

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	271/267 (102%)	268 (99%)	3 (1%)	0	<b>100</b> <b>100</b>

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	263/267 (98%)	260 (99%)	3 (1%)	0	100	100
All	All	534/534 (100%)	528 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	230/224 (103%)	226 (98%)	4 (2%)	60	46
1	B	227/224 (101%)	220 (97%)	7 (3%)	40	21
All	All	457/448 (102%)	446 (98%)	11 (2%)	50	31

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1048	ASN
1	A	1154	ARG
1	A	1162	ASN
1	A	1188	GLN
1	B	2029	LEU
1	B	2048	ASN
1	B	2106	ARG
1	B	2154	ARG
1	B	2162	ASN
1	B	2221[A]	MET
1	B	2221[B]	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1048	ASN
1	A	1162	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	2048	ASN
1	B	2053	HIS
1	B	2058	HIS
1	B	2083	HIS
1	B	2122	ASN
1	B	2162	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	PO4	B	4269	-	4,4,4	2.49	2 (50%)	6,6,6	0.97	0
4	PEP	B	4268	-	9,9,9	2.62	5 (55%)	11,13,13	1.54	3 (27%)
4	PEP	A	3268	-	9,9,9	3.00	5 (55%)	11,13,13	1.28	0
3	PO4	A	3269	-	4,4,4	2.30	2 (50%)	6,6,6	0.93	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEP	B	4268	-	-	0/9/9/9	-
4	PEP	A	3268	-	-	0/9/9/9	-

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	3268	PEP	C2-C1	5.60	1.54	1.49
4	B	4268	PEP	P-O2	4.46	1.66	1.59
3	B	4269	PO4	P-O2	3.80	1.66	1.54
3	A	3269	PO4	P-O3	3.74	1.65	1.54
4	B	4268	PEP	C2-C1	3.66	1.52	1.49
4	A	3268	PEP	C3-C2	3.39	1.41	1.31
4	A	3268	PEP	P-O2	3.33	1.64	1.59
4	A	3268	PEP	P-O3P	3.28	1.67	1.54
3	B	4269	PO4	P-O3	3.11	1.63	1.54
4	B	4268	PEP	O2-C2	-3.09	1.30	1.39
4	A	3268	PEP	P-O2P	2.57	1.64	1.54
4	B	4268	PEP	P-O3P	2.27	1.63	1.54
3	A	3269	PO4	P-O4	-2.11	1.48	1.54
4	B	4268	PEP	P-O2P	2.05	1.62	1.54

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	4268	PEP	O1-C1-C2	-3.55	116.44	121.79
4	B	4268	PEP	O2P-P-O2	2.38	112.50	105.25
4	B	4268	PEP	O2'-C1-O1	2.02	128.22	123.61

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	263/267 (98%)	-0.37	10 (3%) 40 45	11, 17, 30, 47	0
1	B	255/267 (95%)	-0.27	7 (2%) 54 58	11, 19, 33, 39	0
All	All	518/534 (97%)	-0.32	17 (3%) 46 51	11, 18, 33, 47	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1194	GLY	6.0
1	A	1193	LEU	5.6
1	A	1197	SER	5.0
1	A	1198	GLY	4.4
1	B	2264	ILE	4.4
1	A	1264	ILE	3.8
1	A	1195	ASP	3.6
1	B	2199	GLY	3.3
1	A	1192	GLY	3.0
1	B	2147	LYS	2.8
1	B	2191	GLY	2.6
1	A	1263	THR	2.6
1	A	1199	GLY	2.6
1	A	1196[A]	LYS	2.5
1	B	2263	THR	2.5
1	B	2228	GLU	2.2
1	B	2035	GLU	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	PO4	B	4269	5/5	0.98	0.11	22,26,28,33	0
3	PO4	A	3269	5/5	0.99	0.04	23,23,26,31	0
4	PEP	A	3268	10/10	0.99	0.05	12,14,16,17	0
4	PEP	B	4268	10/10	0.99	0.05	13,15,18,19	0
2	CU	A	3270	1/1	1.00	0.04	18,18,18,18	1
2	CU	B	4270	1/1	1.00	0.02	19,19,19,19	1

### 6.5 Other polymers [i](#)

There are no such residues in this entry.