



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 04:11 AM UTC

PDB ID : 9E5P / pdb\_00009e5p  
Title : env2 cobalamin riboswitch aptamer domain in complex with ethynyl-benzofuran-cobalamin  
Authors : Olenginski, L.T.; Batey, R.T.  
Deposited on : 2024-10-28  
Resolution : 2.36 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 2.36 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1596 (2.36-2.36)
Clashscore	190562	1663 (2.36-2.36)
RSRZ outliers	180081	1598 (2.36-2.36)
RNA backbone	3983	1027 (2.66-2.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	76	 58% 30% 9% .
1	B	76	 50% 36% 14%

## 2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 3799 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called RNA (76-MER).


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	76	Total	C	N	O	P	0	0	0
			1636	730	306	523	77			
1	B	76	Total	C	N	O	P	0	0	0
			1636	730	306	523	77			

- Molecule 2 is ethynyl-benzofuran-cobalamin (CCD ID: A1BEX) (formula:  $C_{72}H_{94}CoN_{13}O_{16}P$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	Co	N	O	P	0	0
			103	72	1	13	16	1		
2	B	1	Total	C	Co	N	O	P	0	0
			103	72	1	13	16	1		

- Molecule 3 is N-methylpropane-1,3-diamine (CCD ID: N3D) (formula:  $C_4H_{12}N_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	N	0	0
			6	4	2		
3	B	1	Total	C	N	0	0
			6	4	2		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	9	Total	Mg	0	0
			9	9		
4	B	9	Total	Mg	0	0
			9	9		

- Molecule 5 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	K	0	0
			1	1		
5	B	1	Total	K	0	0
			1	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	138	Total	O	0	0
			138	138		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	151	Total 151	O 151	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: RNA (76-MER)

Chain A: 



#### • Molecule 1: RNA (76-MER)

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	41.11Å 78.06Å 80.11Å 90.00° 90.07° 90.00°	Depositor
Resolution (Å)	20.00 – 2.36 20.00 – 2.36	Depositor EDS
% Data completeness (in resolution range)	91.1 (20.00-2.36) 91.2 (20.00-2.36)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.34 (at 2.35Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, $R_{free}$	0.199 , 0.280 0.200 , 0.280	Depositor DCC
$R_{free}$ test set	1919 reflections (9.10%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	38.4	Xtriage
Anisotropy	0.161	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 35.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.000 for -h,l,k 0.017 for -h,-l,-k 0.036 for h,-k,-l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3799	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.19% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A1BEX, GDP, K, N3D, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.52	0/1802	0.75	2/2807 (0.1%)
1	B	0.50	0/1802	0.72	0/2807
All	All	0.51	0/3604	0.73	2/5614 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	5	A	C2'-C3'-O3'	5.53	121.99	113.70
1	A	66	G	C5'-C4'-C3'	-5.10	107.55	115.20

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1636	0	823	20	0
1	B	1636	0	823	29	0
2	A	103	0	0	6	0
2	B	103	0	0	7	0
3	A	6	0	12	1	0
3	B	6	0	12	1	0
4	A	9	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	9	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	138	0	0	2	0
6	B	151	0	0	6	0
All	All	3799	0	1670	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:101:A1BEX:CPR3	2:A:101:A1BEX:CPR2	1.52	1.51
2:B:101:A1BEX:CPR2	2:B:101:A1BEX:CPR1	1.52	1.51
2:A:101:A1BEX:CPR2	2:A:101:A1BEX:CPR1	1.52	1.49
2:B:101:A1BEX:CPR2	2:B:101:A1BEX:CPR3	1.50	1.49
2:B:101:A1BEX:CPR1	2:B:101:A1BEX:CPR3	2.50	1.09
2:A:101:A1BEX:CPR3	2:A:101:A1BEX:CPR1	2.50	1.08
2:B:101:A1BEX:CPR2	2:B:101:A1BEX:N59	2.47	0.88
2:A:101:A1BEX:CPR1	2:A:101:A1BEX:OP3	2.34	0.84
2:B:101:A1BEX:CPR1	2:B:101:A1BEX:OP3	2.38	0.81
2:B:101:A1BEX:CPR3	2:B:101:A1BEX:OP3	2.37	0.80
2:A:101:A1BEX:CPR3	2:A:101:A1BEX:OP3	2.40	0.79
2:A:101:A1BEX:CPR2	2:A:101:A1BEX:N59	2.54	0.78
1:A:5:A:H2'	1:A:6:A:C8	2.17	0.77
1:B:30:A:N1	6:B:204:HOH:O	2.20	0.74
1:B:11:U:OP1	6:B:201:HOH:O	2.11	0.69
1:A:22:U:H4'	1:A:32:U:O4	1.94	0.68
1:A:45:A:H5''	1:A:45:A:H8	1.60	0.67
1:A:34:C:H2'	1:A:35:G:H8	1.60	0.66
1:B:38:C:OP2	6:B:202:HOH:O	2.14	0.65
1:A:25:C:OP1	6:A:201:HOH:O	2.13	0.65
1:B:31:A:H4'	1:B:32:U:OP1	2.03	0.58
1:B:76:A:OP1	6:B:203:HOH:O	2.17	0.58
1:A:72:U:H2'	1:A:73:A:C8	2.39	0.57
1:A:72:U:H2'	1:A:73:A:H8	1.70	0.57
1:B:28:G:H22	1:B:31:A:H5''	1.70	0.56
1:B:35:G:OP2	3:B:102:N3D:H3	2.06	0.56
1:A:34:C:H2'	1:A:35:G:C8	2.40	0.55
1:A:45:A:H2'	1:A:46:A:C8	2.42	0.54

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:42:C:O3'	2:B:101:A1BEX:N34	2.41	0.53
1:B:45:A:H2'	1:B:46:A:C8	2.44	0.53
1:B:27:U:H2'	1:B:28:G:O4'	2.09	0.53
1:B:29:A:O2'	1:B:30:A:OP1	2.22	0.52
1:B:11:U:O2	1:B:11:U:H2'	2.11	0.50
1:B:1:GDP:H2'	1:B:2:G:H8	1.75	0.50
1:B:1:GDP:H2'	1:B:2:G:C8	2.47	0.49
1:A:27:U:H2'	1:A:28:G:O4'	2.14	0.48
1:A:45:A:H8	1:A:45:A:C5'	2.27	0.47
1:B:34:C:H2'	1:B:35:G:H8	1.79	0.47
1:A:69:A:O2'	1:A:70:C:H5'	2.15	0.47
1:B:22:U:H4'	1:B:32:U:O4	2.15	0.46
1:A:21:G:H3'	1:A:22:U:H5''	1.98	0.45
1:B:5:A:H2'	1:B:6:A:C8	2.52	0.45
1:A:23:G:H2'	1:A:24:A:C8	2.52	0.45
1:A:35:G:OP2	3:A:102:N3D:N5	2.47	0.44
1:A:70:C:O2'	6:A:203:HOH:O	2.21	0.44
1:B:18:A:H4'	1:B:19:A:OP2	2.18	0.43
1:A:73:A:H2'	1:A:74:C:C6	2.52	0.43
1:B:70:C:H2'	1:B:71:C:O4'	2.18	0.43
1:B:54:C:H2'	6:B:317:HOH:O	2.17	0.43
1:B:61:C:H2'	1:B:62:G:O4'	2.20	0.42
1:A:26:G:O2'	1:A:27:U:H5'	2.20	0.42
1:A:66:G:H3'	1:A:67:C:C6	2.55	0.42
1:B:73:A:H2'	1:B:74:C:C6	2.55	0.42
1:B:35:G:H2'	1:B:36:U:H6	1.85	0.41
1:B:38:C:H5''	6:B:242:HOH:O	2.19	0.41
1:B:39:A:H2'	1:B:40:C:O4'	2.20	0.41
1:B:34:C:H2'	1:B:35:G:C8	2.56	0.41
1:B:14:U:O5'	1:B:14:U:H6	2.04	0.41
1:B:19:A:H8	1:B:19:A:O5'	2.04	0.41
1:A:70:C:H2'	1:A:71:C:O4'	2.22	0.40
1:B:45:A:O2'	1:B:46:A:H5'	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

### 5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

### 5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	75/76 (98%)	17 (22%)	3 (4%)
1	B	75/76 (98%)	15 (20%)	3 (4%)
All	All	150/152 (98%)	32 (21%)	6 (4%)

All (32) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	A	5	A
1	A	6	A
1	A	11	U
1	A	12	A
1	A	18	A
1	A	21	G
1	A	22	U
1	A	23	G
1	A	32	U
1	A	38	C
1	A	45	A
1	A	46	A
1	A	54	C
1	A	55	A
1	A	56	U
1	A	65	U
1	A	66	G
1	B	5	A
1	B	11	U
1	B	12	A
1	B	18	A
1	B	21	G
1	B	22	U
1	B	23	G
1	B	29	A
1	B	30	A
1	B	32	U
1	B	54	C
1	B	56	U

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	65	U
1	B	66	G
1	B	76	A

All (6) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	A	5	A
1	A	11	U
1	A	45	A
1	B	11	U
1	B	29	A
1	B	31	A

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
1	GDP	A	1	4,1	29,30,30	1.68	5 (17%)	45,47,47	2.37	16 (35%)
1	GDP	B	1	1	29,30,30	1.52	8 (27%)	45,47,47	2.45	26 (57%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	GDP	A	1	4,1	-	2/16/32/32	0/3/3/3
1	GDP	B	1	1	-	5/16/32/32	0/3/3/3

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1	GDP	PA-O3A	4.70	1.64	1.59
1	A	1	GDP	O3'-C3'	3.23	1.50	1.43
1	A	1	GDP	PB-O1B	3.09	1.60	1.50
1	B	1	GDP	O3'-C3'	2.85	1.50	1.43
1	B	1	GDP	PA-O3A	2.82	1.62	1.59
1	B	1	GDP	PB-O1B	2.60	1.58	1.50
1	B	1	GDP	C4-N9	-2.42	1.31	1.38
1	A	1	GDP	C4-N9	-2.40	1.32	1.38
1	B	1	GDP	C5-C4	2.36	1.45	1.38
1	B	1	GDP	O6-C6	2.34	1.28	1.23
1	B	1	GDP	C8-N7	2.28	1.38	1.32
1	A	1	GDP	C8-N7	2.21	1.38	1.32
1	B	1	GDP	O4'-C4'	2.10	1.49	1.45

All (42) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1	GDP	C5-C4-N3	-5.65	119.40	128.39
1	A	1	GDP	O6-C6-C5	-5.43	112.20	126.53
1	B	1	GDP	O3A-PA-O1A	-5.25	94.90	110.70
1	A	1	GDP	C2-N3-C4	5.21	121.27	112.30
1	B	1	GDP	C5-C4-N3	-4.75	120.84	128.39
1	A	1	GDP	N9-C4-N3	4.55	135.04	125.95
1	B	1	GDP	O3B-PB-O3A	4.11	118.42	104.64
1	B	1	GDP	O6-C6-C5	-4.10	115.70	126.53
1	B	1	GDP	C2-N3-C4	4.08	119.33	112.30
1	B	1	GDP	O4'-C1'-N9	-3.99	99.32	108.36
1	A	1	GDP	C2-N1-C6	-3.98	117.90	125.11
1	B	1	GDP	N9-C4-N3	3.71	133.38	125.95
1	A	1	GDP	O4'-C4'-C3'	-3.36	98.49	105.15
1	A	1	GDP	C5-C6-N1	3.26	121.56	113.25
1	A	1	GDP	O6-C6-N1	3.26	126.24	120.11
1	B	1	GDP	N2-C2-N3	-3.05	113.73	119.67
1	B	1	GDP	C8-N9-C4	3.03	111.70	106.03
1	B	1	GDP	C2-N1-C6	-3.00	119.67	125.11
1	A	1	GDP	O3A-PA-O1A	-2.99	101.72	110.70
1	B	1	GDP	C2'-C3'-C4'	-2.90	97.01	102.61
1	B	1	GDP	O2'-C2'-C1'	-2.79	100.50	110.10
1	A	1	GDP	O4'-C1'-N9	-2.71	102.21	108.36
1	B	1	GDP	O3'-C3'-C4'	2.67	118.74	111.08
1	B	1	GDP	C6-C5-N7	2.66	135.13	130.29
1	B	1	GDP	O6-C6-N1	2.63	125.06	120.11
1	B	1	GDP	N2-C2-N1	2.58	122.20	116.76

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1	GDP	N9-C8-N7	-2.50	108.77	113.40
1	A	1	GDP	C8-N9-C4	2.43	110.58	106.03
1	A	1	GDP	O3'-C3'-C4'	2.40	117.97	111.08
1	B	1	GDP	O4'-C4'-C3'	-2.39	100.40	105.15
1	B	1	GDP	C4'-O4'-C1'	-2.38	104.21	109.47
1	B	1	GDP	O2A-PA-O1A	2.37	123.46	112.44
1	B	1	GDP	C5-C6-N1	2.35	119.25	113.25
1	B	1	GDP	O5'-PA-O1A	2.35	118.25	108.94
1	A	1	GDP	O2A-PA-O1A	2.33	123.26	112.44
1	B	1	GDP	O5'-C5'-C4'	-2.30	101.15	108.99
1	B	1	GDP	O3'-C3'-C2'	-2.29	104.48	111.82
1	A	1	GDP	O5'-C5'-C4'	-2.23	101.40	108.99
1	B	1	GDP	O2'-C2'-C3'	-2.19	104.80	111.82
1	B	1	GDP	C3'-C2'-C1'	2.17	105.56	101.46
1	A	1	GDP	N9-C8-N7	-2.05	109.59	113.40
1	A	1	GDP	O5'-PA-O1A	2.05	117.06	108.94

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	1	GDP	O4'-C4'-C5'-O5'
1	A	1	GDP	C3'-C4'-C5'-O5'
1	B	1	GDP	C5'-O5'-PA-O3A
1	B	1	GDP	C5'-O5'-PA-O1A
1	B	1	GDP	C5'-O5'-PA-O2A
1	B	1	GDP	O4'-C4'-C5'-O5'
1	B	1	GDP	C3'-C4'-C5'-O5'

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	1	GDP	2	0

## 5.5 Carbohydrates

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry

Of 24 ligands modelled in this entry, 20 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	A1BEX	A	101	-	103,115,115	0.70	1 (0%)	157,189,189	1.44	20 (12%)
3	N3D	B	102	-	5,5,5	0.60	0	4,4,4	1.10	0
3	N3D	A	102	-	5,5,5	0.52	0	4,4,4	0.79	0
2	A1BEX	B	101	-	103,115,115	0.64	0	157,189,189	1.40	17 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1BEX	A	101	-	-	8/58/247/247	0/5/13/13
3	N3D	B	102	-	-	0/3/3/3	-
3	N3D	A	102	-	-	1/3/3/3	-
2	A1BEX	B	101	-	-	8/58/247/247	0/5/13/13

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	101	A1BEX	C6-C5	-2.38	1.37	1.44

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	101	A1BEX	O01-C07-C01	-6.86	104.61	108.22
2	A	101	A1BEX	O01-C07-C01	-6.69	104.71	108.22
2	B	101	A1BEX	CB9-CB8-NB1	5.10	107.57	105.30
2	A	101	A1BEX	CB9-CB8-NB1	4.94	107.49	105.30
2	B	101	A1BEX	O01-C08-C06	-4.69	101.32	104.51
2	B	101	A1BEX	C14-N23-C11	4.15	111.10	105.44

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	101	A1BEX	O01-C07-O02	4.00	125.31	121.09
2	B	101	A1BEX	C03-C04-C2L	-3.52	114.55	120.78
2	A	101	A1BEX	C14-N23-C11	3.49	110.20	105.44
2	A	101	A1BEX	CPR2-CPR1-N59	3.49	118.06	112.92
2	B	101	A1BEX	C04-C05-C06	-3.24	116.73	120.77
2	A	101	A1BEX	C03-C04-C2L	-3.19	115.13	120.78
2	A	101	A1BEX	C31-C30-C3	-3.19	105.61	114.65
2	A	101	A1BEX	C06-C01-C07	3.18	109.75	108.38
2	A	101	A1BEX	O62-C61-C60	3.12	127.38	120.87
2	B	101	A1BEX	C03-C02-C01	-2.87	116.13	120.86
2	A	101	A1BEX	C20-C1-C2	-2.79	110.49	113.66
2	A	101	A1BEX	C05-C04-C2L	2.76	125.19	120.22
2	A	101	A1BEX	C18-C17-C16	-2.73	97.40	100.69
2	A	101	A1BEX	C5-C6-N22	-2.59	121.11	126.52
2	B	101	A1BEX	C3-C4-N21	2.49	112.95	109.21
2	B	101	A1BEX	C30-C3-C4	2.48	114.66	108.51
2	A	101	A1BEX	C36-C7-C8	-2.45	107.53	112.05
2	B	101	A1BEX	C02-C01-C06	2.45	123.93	120.88
2	B	101	A1BEX	C35-C5-C4	-2.44	113.57	118.42
2	B	101	A1BEX	C7-C6-N22	-2.39	106.65	110.96
2	A	101	A1BEX	C36-C7-C6	2.31	124.39	112.41
2	A	101	A1BEX	OR6-CR1-NB1	2.31	112.52	108.09
2	A	101	A1BEX	C3-C4-C5	-2.31	120.81	124.32
2	B	101	A1BEX	OP5-P1-OP4	2.26	122.96	112.44
2	B	101	A1BEX	C42-C41-C8	2.23	120.96	114.65
2	A	101	A1BEX	C47-C12-C46	2.20	113.05	109.41
2	A	101	A1BEX	C7-C6-C5	2.14	129.65	126.26
2	A	101	A1BEX	C55-C17-C18	-2.12	107.07	111.12
2	B	101	A1BEX	C55-C17-C16	2.11	120.72	116.59
2	B	101	A1BEX	C2-C3-C4	-2.03	99.22	101.68
2	A	101	A1BEX	C60-C61-N63	-2.01	111.38	116.19

There are no chirality outliers.

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	101	A1BEX	C13-C48-C49-C50
2	B	101	A1BEX	C42-C41-C8-C9
2	B	101	A1BEX	OR6-CR4-CR5-OR8
2	B	101	A1BEX	C42-C41-C8-C7
2	B	101	A1BEX	CR3-CR4-CR5-OR8
2	A	101	A1BEX	C2-C26-C27-N29

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
2	A	101	A1BEX	C2-C26-C27-O28
2	B	101	A1BEX	C41-C42-C43-O44
2	A	101	A1BEX	C41-C42-C43-O44
3	A	102	N3D	N1-C2-C3-C4
2	A	101	A1BEX	C41-C42-C43-N45
2	B	101	A1BEX	C18-C60-C61-O62
2	B	101	A1BEX	C41-C42-C43-N45
2	A	101	A1BEX	C25-C2-C26-C27
2	B	101	A1BEX	CPR2-OP3-P1-OP5
2	A	101	A1BEX	C17-C18-C60-C61
2	A	101	A1BEX	C19-C18-C60-C61


There are no ring outliers.

4 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	101	A1BEX	6	0
3	B	102	N3D	1	0
3	A	102	N3D	1	0
2	B	101	A1BEX	7	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	75/76 (98%)	-0.44	0 100 100	25, 34, 44, 59	0
1	B	75/76 (98%)	-0.38	0 100 100	27, 35, 45, 70	0
All	All	150/152 (98%)	-0.41	0 100 100	25, 35, 45, 70	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	GDP	A	1	28/28	0.81	0.11	26,42,73,83	0
1	GDP	B	1	28/28	0.86	0.12	26,38,83,92	0

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	MG	B	103	1/1	0.60	0.18	57,57,57,57	0
3	N3D	B	102	6/6	0.71	0.20	18,26,34,34	0
4	MG	A	103	1/1	0.81	0.18	65,65,65,65	0
4	MG	A	107	1/1	0.82	0.15	35,35,35,35	0
3	N3D	A	102	6/6	0.86	0.10	22,28,29,30	0
5	K	B	112	1/1	0.88	0.10	52,52,52,52	0
4	MG	B	110	1/1	0.92	0.08	32,32,32,32	0
4	MG	B	107	1/1	0.92	0.06	38,38,38,38	0
4	MG	A	104	1/1	0.93	0.06	26,26,26,26	0
2	A1BEX	B	101	103/103	0.93	0.09	27,32,42,48	0
2	A1BEX	A	101	103/103	0.94	0.09	19,30,36,45	0
4	MG	B	108	1/1	0.95	0.06	40,40,40,40	0
4	MG	A	111	1/1	0.96	0.09	32,32,32,32	0
4	MG	A	108	1/1	0.96	0.04	27,27,27,27	0
4	MG	A	109	1/1	0.97	0.04	29,29,29,29	0
4	MG	A	110	1/1	0.97	0.04	31,31,31,31	0
4	MG	B	109	1/1	0.97	0.04	34,34,34,34	0
4	MG	B	104	1/1	0.97	0.05	36,36,36,36	0
5	K	A	112	1/1	0.97	0.04	41,41,41,41	0
4	MG	B	105	1/1	0.97	0.04	30,30,30,30	0
4	MG	A	105	1/1	0.98	0.03	30,30,30,30	0
4	MG	A	106	1/1	0.98	0.04	29,29,29,29	0
4	MG	B	106	1/1	0.98	0.03	30,30,30,30	0
4	MG	B	111	1/1	0.99	0.02	26,26,26,26	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around A1BEX B 101:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around A1BEX A 101:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.