



wwPDB EM Validation Summary Report ⓘ

Mar 10, 2026 – 08:52 AM UTC

PDB ID : 9E87 / pdb_00009e87
EMDB ID : EMD-47709
Title : De Novo Mycobacterium tuberculosis transcription initiation pre-RPO promoter complex with open Beta' clamp (RNA Polymerase with Sigma-A, CarD, and RbpA)
Authors : Brewer, J.J.; Campbell, E.A.; Darst, S.A.
Deposited on : 2024-11-05
Resolution : 3.60 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	12797 (3.10 - 4.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	347	
1	B	347	
2	C	1177	
3	D	1333	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	E	110	<div><div><div></div><div></div><div></div></div><div>59%15%25%</div></div>
5	F	550	<div><div><div></div><div></div><div></div></div><div>42%13%45%</div></div>
6	J	111	<div><div><div></div><div></div><div></div></div><div>5%69%26%5%</div></div>
7	M	285	<div><div><div></div><div></div><div></div></div><div>31%40%15%45%</div></div>
8	N	127	<div><div><div></div><div></div><div></div></div><div>8%28%20%51%</div></div>
9	T	127	<div><div><div></div><div></div><div></div></div><div>8%22%20%57%</div></div>

2 Entry composition [i](#)

There are 11 unique types of molecules in this entry. The entry contains 57974 atoms, of which 28500 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase subunit alpha.

Mol	Chain	Residues	Atoms						AltConf	Trace
1	A	221	Total	C	H	N	O	S	0	0
			3400	1059	1720	288	331	2		
1	B	237	Total	C	H	N	O	S	0	0
			3559	1115	1794	301	346	3		

- Molecule 2 is a protein called DNA-directed RNA polymerase subunit beta.

Mol	Chain	Residues	Atoms						AltConf	Trace
2	C	1111	Total	C	H	N	O	S	0	0
			17090	5380	8508	1499	1664	39		

- Molecule 3 is a protein called DNA-directed RNA polymerase subunit beta'.

Mol	Chain	Residues	Atoms						AltConf	Trace
3	D	1269	Total	C	H	N	O	S	0	0
			19903	6209	9990	1801	1861	42		

There are 17 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	-8	LEU	-	expression tag	UNP P0A675
D	-7	ALA	-	expression tag	UNP P0A675
D	-6	ARG	-	expression tag	UNP P0A675
D	-5	HIS	-	expression tag	UNP P0A675
D	-4	GLY	-	expression tag	UNP P0A675
D	-3	GLY	-	expression tag	UNP P0A675
D	-2	SER	-	expression tag	UNP P0A675
D	-1	GLY	-	expression tag	UNP P0A675
D	0	ALA	-	expression tag	UNP P0A675
D	1317	HIS	-	expression tag	UNP P0A675
D	1318	HIS	-	expression tag	UNP P0A675
D	1319	HIS	-	expression tag	UNP P0A675
D	1320	HIS	-	expression tag	UNP P0A675

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	1321	HIS	-	expression tag	UNP P0A675
D	1322	HIS	-	expression tag	UNP P0A675
D	1323	HIS	-	expression tag	UNP P0A675
D	1324	HIS	-	expression tag	UNP P0A675

- Molecule 4 is a protein called DNA-directed RNA polymerase subunit omega.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	E	83	Total	C	H	N	O	0	0
			1294	414	645	108	127		

- Molecule 5 is a protein called RNA polymerase sigma factor SigA.

Mol	Chain	Residues	Atoms						AltConf	Trace
5	F	305	Total	C	H	N	O	S	0	0
			4867	1512	2450	437	459	9		

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	-21	HIS	-	expression tag	UNP P0A603
F	-20	HIS	-	expression tag	UNP P0A603
F	-19	HIS	-	expression tag	UNP P0A603
F	-18	HIS	-	expression tag	UNP P0A603
F	-17	HIS	-	expression tag	UNP P0A603
F	-16	HIS	-	expression tag	UNP P0A603
F	-15	HIS	-	expression tag	UNP P0A603
F	-14	HIS	-	expression tag	UNP P0A603
F	-13	HIS	-	expression tag	UNP P0A603
F	-12	HIS	-	expression tag	UNP P0A603
F	-11	SER	-	expression tag	UNP P0A603
F	-10	SER	-	expression tag	UNP P0A603
F	-9	GLY	-	expression tag	UNP P0A603
F	-8	LEU	-	expression tag	UNP P0A603
F	-7	GLU	-	expression tag	UNP P0A603
F	-6	VAL	-	expression tag	UNP P0A603
F	-5	LEU	-	expression tag	UNP P0A603
F	-4	PHE	-	expression tag	UNP P0A603
F	-3	GLN	-	expression tag	UNP P0A603
F	-2	GLY	-	expression tag	UNP P0A603
F	-1	PRO	-	expression tag	UNP P0A603
F	0	HIS	-	expression tag	UNP P0A603

- Molecule 6 is a protein called RNA polymerase-binding protein RbpA.

Mol	Chain	Residues	Atoms						AltConf	Trace
6	J	106	Total	C	H	N	O	S	0	0
			1687	529	830	160	165	3		

- Molecule 7 is a protein called Ubiquitin-like protein SMT3,RNA polymerase-binding transcription factor CarD.

Mol	Chain	Residues	Atoms						AltConf	Trace
7	M	158	Total	C	H	N	O	S	0	0
			2481	771	1248	223	238	1		

There are 25 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
M	-122	MET	-	expression tag	UNP Q12306
M	-121	GLY	-	expression tag	UNP Q12306
M	-120	HIS	-	expression tag	UNP Q12306
M	-119	HIS	-	expression tag	UNP Q12306
M	-118	HIS	-	expression tag	UNP Q12306
M	-117	HIS	-	expression tag	UNP Q12306
M	-116	HIS	-	expression tag	UNP Q12306
M	-115	HIS	-	expression tag	UNP Q12306
M	-114	HIS	-	expression tag	UNP Q12306
M	-113	HIS	-	expression tag	UNP Q12306
M	-112	HIS	-	expression tag	UNP Q12306
M	-111	HIS	-	expression tag	UNP Q12306
M	-110	SER	-	expression tag	UNP Q12306
M	-109	SER	-	expression tag	UNP Q12306
M	-108	GLY	-	expression tag	UNP Q12306
M	-107	HIS	-	expression tag	UNP Q12306
M	-106	ILE	-	expression tag	UNP Q12306
M	-105	GLU	-	expression tag	UNP Q12306
M	-104	GLY	-	expression tag	UNP Q12306
M	-103	ARG	-	expression tag	UNP Q12306
M	-102	HIS	-	expression tag	UNP Q12306
M	-101	MET	-	expression tag	UNP Q12306
M	-100	ALA	-	expression tag	UNP Q12306
M	-99	SER	-	expression tag	UNP Q12306
M	0	SER	-	linker	UNP Q12306

- Molecule 8 is a DNA chain called DNA (62-MER).

Mol	Chain	Residues	Atoms						AltConf	Trace
8	N	62	Total	C	H	N	O	P	0	0
			1974	610	699	239	364	62		

- Molecule 9 is a DNA chain called DNA (54-MER).


Mol	Chain	Residues	Atoms						AltConf	Trace
9	T	54	Total	C	H	N	O	P	0	0
			1716	530	616	184	332	54		

- Molecule 10 is ZINC ION (CCD ID: ZN) (formula: Zn).



Mol	Chain	Residues	Atoms		AltConf
10	D	2	Total	Zn	0
			2	2	


- Molecule 11 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		AltConf
11	D	1	Total	Mg	0
			1	1	




• Molecule 3: DNA-directed RNA polymerase subunit beta'







- Molecule 4: DNA-directed RNA polymerase subunit omega



- Molecule 5: RNA polymerase sigma factor SigA







4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	102319	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	51.83	Depositor
Minimum defocus (nm)	400	Depositor
Maximum defocus (nm)	2200	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	2.671	Depositor
Minimum map value	-1.807	Depositor
Average map value	0.004	Depositor
Map value standard deviation	0.063	Depositor
Recommended contour level	0.15	Depositor
Map size (Å)	275.456, 275.456, 275.456	wwPDB
Map dimensions	256, 256, 256	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.076, 1.076, 1.076	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.22	0/1706	0.45	0/2322
1	B	0.21	0/1792	0.50	2/2442 (0.1%)
2	C	0.24	0/8741	0.46	0/11857
3	D	0.23	0/10077	0.46	1/13623 (0.0%)
4	E	0.20	0/662	0.44	0/901
5	F	0.23	0/2448	0.46	0/3302
6	J	0.25	0/873	0.53	0/1179
7	M	0.19	0/1249	0.52	0/1689
8	N	0.37	0/1432	0.58	1/2205 (0.0%)
9	T	0.36	0/1228	0.56	0/1890
All	All	0.24	0/30208	0.48	4/41410 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
6	J	0	1
7	M	0	1
All	All	0	2

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	1069	ASP	N-CA-C	-7.97	103.36	113.72
8	N	29	DT	O4'-C1'-N1	6.10	117.55	108.40
1	B	28	PRO	CA-C-N	5.40	131.42	121.70
1	B	28	PRO	C-N-CA	5.40	131.42	121.70

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
6	J	106	ARG	Sidechain
7	M	109	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1680	1720	1720	42	0
1	B	1765	1794	1794	30	0
2	C	8582	8508	8500	206	0
3	D	9913	9990	9992	188	0
4	E	649	645	645	17	0
5	F	2417	2450	2448	68	0
6	J	857	830	828	25	0
7	M	1233	1248	1248	32	0
8	N	1275	699	701	43	0
9	T	1100	616	618	44	0
10	D	2	0	0	0	0
11	D	1	0	0	0	0
All	All	29474	28500	28494	631	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 631 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:N:16:DT:C2'	8:N:17:DT:H71	1.65	1.26
9:T:77:DA:C2'	9:T:78:DT:H72	1.76	1.15
9:T:66:DC:H2''	9:T:67:DT:C7	1.81	1.11
8:N:16:DT:H2''	8:N:17:DT:C7	1.83	1.09
9:T:77:DA:H2'	9:T:78:DT:H72	1.13	1.08

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	219/347 (63%)	203 (93%)	16 (7%)	0	100	100
1	B	235/347 (68%)	212 (90%)	23 (10%)	0	100	100
2	C	1109/1177 (94%)	1009 (91%)	99 (9%)	1 (0%)	48	79
3	D	1263/1333 (95%)	1177 (93%)	81 (6%)	5 (0%)	30	61
4	E	81/110 (74%)	73 (90%)	8 (10%)	0	100	100
5	F	303/550 (55%)	285 (94%)	18 (6%)	0	100	100
6	J	104/111 (94%)	99 (95%)	4 (4%)	1 (1%)	12	45
7	M	156/285 (55%)	152 (97%)	4 (3%)	0	100	100
All	All	3470/4260 (82%)	3210 (92%)	253 (7%)	7 (0%)	44	72

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	C	231	ARG
3	D	416	ASN
3	D	1008	THR
3	D	309	PRO
3	D	1070	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	190/297 (64%)	189 (100%)	1 (0%)	81	80

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	195/297 (66%)	194 (100%)	1 (0%)	81	80
2	C	933/997 (94%)	931 (100%)	2 (0%)	87	85
3	D	1049/1107 (95%)	1044 (100%)	5 (0%)	81	80
4	E	69/90 (77%)	68 (99%)	1 (1%)	59	71
5	F	255/447 (57%)	254 (100%)	1 (0%)	84	81
6	J	90/97 (93%)	90 (100%)	0	100	100
7	M	128/240 (53%)	128 (100%)	0	100	100
All	All	2909/3572 (81%)	2898 (100%)	11 (0%)	81	81

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	D	581	MET
3	D	676	LEU
5	F	281	MET
4	E	33	LEU
3	D	75	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
3	D	606	HIS
3	D	1104	HIS
7	M	133	GLN
5	F	234	GLN
5	F	353	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-47709. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)

6.2.1 Primary map



X Index: 128




Y Index: 128




Z Index: 128


6.2.2 Raw map



X Index: 128



Y Index: 128



Z Index: 128


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 109




Y Index: 126




Z Index: 142


6.3.2 Raw map



X Index: 126



Y Index: 134




Z Index: 141


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)

6.4.1 Primary map



6.4.2 Raw map



The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

This section was not generated.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 276 nm^3 ; this corresponds to an approximate mass of 249 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.278 \AA^{-1}

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.278 \AA^{-1}

8.2 Resolution estimates [i](#)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.60	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	4.06	6.15	4.11

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.06 differs from the reported value 3.6 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-47709 and PDB model 9E87. Per-residue inclusion information can be found in section 3 on page 8.

9.1 Map-model overlay [i](#)


This section was not generated.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.15).

9.4 Atom inclusion [i](#)



At the recommended contour level, 80% of all backbone atoms, 80% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.15) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div></div> 0.8000	<div></div> 0.3640
A	<div></div> 0.8840	<div></div> 0.4470
B	<div></div> 0.8310	<div></div> 0.3760
C	<div></div> 0.8590	<div></div> 0.4170
D	<div></div> 0.8230	<div></div> 0.3750
E	<div></div> 0.8030	<div></div> 0.4060
F	<div></div> 0.7920	<div></div> 0.3090
J	<div></div> 0.7330	<div></div> 0.2990
M	<div></div> 0.3690	<div></div> 0.1890
N	<div></div> 0.7490	<div></div> 0.2180
T	<div></div> 0.7100	<div></div> 0.2210

1.0

0.0

<0.0