Full wwPDB X-ray Structure Validation Report (i)

Oct 10, 2021 – 03:39 PM EDT

PDB ID	:	3EBI
Title	:	Structure of the M1 Alanylaminopeptidase from malaria complexed with the phosphinate dipeptide analog
Authors	:	McGowan, S.; Porter, C.J.; Buckle, A.M.; Whisstock, J.C.
Deposited on	:	2008-08-27
Resolution	:	2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.23.2
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.23.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	890	3% 93% 7%

2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 8120 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called M1 family aminopeptidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	889	7320	4707	1194	1392	27	18	5	0

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	213	GLN	ASN	engineered mutation	UNP O96935
A	223	GLN	ASN	engineered mutation	UNP O96935
A	378	PRO	HIS	engineered mutation	UNP O96935
A	501	GLN	ASN	engineered mutation	UNP O96935
A	745	GLN	ASN	engineered mutation	UNP O96935
A	795	GLN	ASN	engineered mutation	UNP O96935
A	1069	GLN	ASN	engineered mutation	UNP O96935

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
2	A	1	1	1	0	0

- Molecule 3 is (2S)-3-[(R)-[(1S)-1-amino-3-phenylpropyl](hydroxy)phosphoryl]-2-benzylpropanoic acid (three-letter code: BEY) (formula: $C_{19}H_{24}NO_4P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	25	19	1	4	1	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O			
4	A	1	6	3	3		0	0
			Total	C	O			
4	A	1	6	3	3		0	0
			Total	C	O			
4	A	1	6	3	3		0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	A	1	6	3	3	0	0
			Total	C	O		
4	A	1	6	3	3	0	0

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
5	A	4	4	4	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	A	740	740	740	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: M1 family aminopeptidase

Chain A:





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	75.85Å 108.63Å 118.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.61 – 2.00 28.60 – 2.00	Depositor EDS
% Data completeness (in resolution range)	90.7 (28.61-2.00) 90.7 (28.60-2.00)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.06	Depositor
$< I/\sigma(I) >$ [1]	2.65 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.175 , 0.221 0.174 , 0.220	Depositor DCC
R_{free} test set	3078 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å^2)	22.9	Xtriage
Anisotropy	0.774	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.37 , 56.1	EDS
L-test for twinning[2]	$< \|L\| > = 0.46$, $< L^2 > = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	8120	wwPDB-VP
Average B, all atoms (Å^2)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.97% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, BEY, MG, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.49	0/7499	0.56	0/10136

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7320	0	7262	31	0
2	A	1	0	0	0	0
3	A	25	0	22	4	0
4	A	30	0	40	1	0
5	A	4	0	0	0	0
6	A	740	0	0	7	0
All	All	8120	0	7324	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:265:GLU:OE2	6:A:1596:HOH:O	1.99	0.80
1:A:969[A]:ARG:HD3	6:A:1617:HOH:O	1.87	0.73
1:A:538:THR:HG21	1:A:819:GLY:HA3	1.77	0.67
1:A:969[B]:ARG:HD2	6:A:1154:HOH:O	1.99	0.63
1:A:811:LEU:HB3	1:A:867:MET:SD	2.42	0.60
1:A:745:GLN:NE2	6:A:1618:HOH:O	2.34	0.60
1:A:973:LYS:HG2	1:A:1009:ARG:HH21	1.65	0.60
1:A:809:TYR:HD2	1:A:810:LEU:HD12	1.68	0.57
1:A:398:THR:HG21	1:A:407:LYS:HE3	1.89	0.55
1:A:488:ALA:O	1:A:492:THR:HG23	2.08	0.53
1:A:907:LYS:HD2	6:A:1485:HOH:O	2.07	0.53
1:A:415:GLU:HG2	6:A:1142:HOH:O	2.11	0.51
1:A:228:ARG:HG2	1:A:293:GLU:HG3	1.94	0.50
1:A:451:LEU:HG	1:A:470:PHE:CE2	2.45	0.50
1:A:973:LYS:HA	1:A:1009:ARG:HE	1.78	0.48
1:A:281:LYS:HE3	1:A:282:PHE:CZ	2.49	0.48
1:A:891:GLN:OE1	4:A:1086:GOL:H2	2.14	0.48
1:A:489:ARG:NH1	3:A:1085:BEY:H6	2.30	0.46
1:A:442:LEU:HB3	1:A:502:TYR:CE2	2.51	0.46
1:A:827:LEU:HD23	1:A:849[B]:LYS:HD2	1.98	0.46
1:A:519:GLU:HA	1:A:519:GLU:OE1	2.16	0.45
1:A:271:TYR:CZ	1:A:273:ASN:HA	2.51	0.45
1:A:199:HIS:O	1:A:567:SER:HA	2.18	0.44
1:A:264:VAL:HG23	1:A:267:GLU:HB3	2.00	0.43
1:A:575:TYR:CE1	3:A:1085:BEY:H13	2.53	0.43
3:A:1085:BEY:H18	3:A:1085:BEY:H17	1.75	0.43
1:A:626:GLN:O	1:A:630:MET:HG3	2.19	0.42
1:A:1041:GLU:N	1:A:1042:PRO:CD	2.83	0.42
1:A:834:ILE:HG12	1:A:1048:LYS:HD3	2.02	0.41
1:A:492:THR:CG2	6:A:1291:HOH:O	2.69	0.41
1:A:496:HIS:CG	3:A:1085:BEY:H2	2.56	0.40
1:A:586:VAL:HA	1:A:589:MET:HE2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	892/890 (100%)	875 (98%)	17 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	822/822 (100%)	809 (98%)	13 (2%)	62	67

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	263	LEU
1	A	370	HIS
1	A	379	LEU
1	A	436	ASP
1	A	541	LEU
1	A	555	ASP
1	A	688	PHE
1	A	765	TYR
1	A	840	LEU
1	A	901	LEU
1	A	982	LEU
1	A	1037	THR
1	A	1045	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	199	HIS
1	A	212	ASN
1	A	260	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	481	ASN
1	A	527	ASN
1	A	543	HIS
1	A	915	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 11 ligands modelled in this entry, 5 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	GOL	A	2	-	5,5,5	0.36	0	5,5,5	0.56	0
4	GOL	A	4	-	5,5,5	0.34	0	5,5,5	0.49	0
4	GOL	A	5	-	5,5,5	0.37	0	5,5,5	0.33	0
4	GOL	A	1086	-	5,5,5	0.52	0	5,5,5	0.69	0
4	GOL	A	3	-	5,5,5	0.41	0	5,5,5	0.28	0
3	BEY	A	1085	2	19,26,26	0.87	1 (5%)	21,35,35	1.15	2 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	2	-	-	2/4/4/4	-
4	GOL	A	4	-	-	4/4/4/4	-
4	GOL	A	5	-	-	0/4/4/4	-
4	GOL	A	1086	-	-	3/4/4/4	-
4	GOL	A	3	-	-	0/4/4/4	-
3	BEY	A	1085	2	-	5/14/24/24	0/2/2/2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1085	BEY	P-C17	2.64	1.82	1.79

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	A	1085	BEY	O3-P-C19	-4.11	97.65	106.87
3	A	1085	BEY	C1-C7-C8	-2.16	111.02	113.81

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1085	BEY	C18-C19-P-O4
3	A	1085	BEY	C15-C18-C19-N
4	A	1086	GOL	O1-C1-C2-O2
4	A	1086	GOL	O1-C1-C2-C3
4	A	4	GOL	O2-C2-C3-O3
4	A	2	GOL	C1-C2-C3-O3
4	A	4	GOL	O1-C1-C2-C3
4	A	4	GOL	C1-C2-C3-O3
4	A	2	GOL	O2-C2-C3-O3
4	A	1086	GOL	O2-C2-C3-O3
4	A	4	GOL	O1-C1-C2-O2
3	A	1085	BEY	P-C17-C8-C9
3	A	1085	BEY	P-C17-C8-C7
3	A	1085	BEY	C8-C17-P-O4

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1086	GOL	1	0
3	A	1085	BEY	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand BEY A 1085



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	889/890 (99%)	0.02	25 (2%) 53 51	7, 17, 31, 47	9 (1%)

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	815	HIS	4.7
1	A	940	ASP	4.3
1	A	634	ASP	3.7
1	A	635	ASN	3.5
1	A	633	ALA	3.4
1	A	636	SER	3.3
1	A	403	TYR	3.0
1	A	200	TYR	2.9
1	A	196	PRO	2.8
1	A	584[A]	SER	2.7
1	A	1066	GLN	2.6
1	A	637	ALA	2.6
1	A	682	LYS	2.6
1	A	790	LYS	2.5
1	A	683	GLU	2.5
1	A	989	ASP	2.5
1	A	543	HIS	2.4
1	A	670[A]	HIS	2.3
1	A	516	THR	2.2
1	A	612	GLY	2.2
1	A	363	PHE	2.1
1	A	663	GLU	2.1
1	A	632	LYS	2.0
1	A	407	LYS	2.0
1	A	326	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
5	MG	A	8	1/1	0.72	0.11	40,40,40,40	0
5	MG	A	7	1/1	0.75	0.11	44,44,44,44	0
4	GOL	A	4	6/6	0.86	0.21	22,25,33,38	0
5	MG	A	10	1/1	0.89	0.30	54,54,54,54	0
4	GOL	A	5	6/6	0.90	0.21	40,41,42,43	0
4	GOL	A	2	6/6	0.90	0.16	24,26,34,37	0
5	MG	A	9	1/1	0.90	0.12	46,46,46,46	0
4	GOL	A	1086	6/6	0.91	0.14	12,13,16,21	0
4	GOL	A	3	6/6	0.93	0.14	33,34,35,38	0
3	BEY	A	1085	25/25	0.94	0.16	16,22,29,35	0
2	ZN	A	1	1/1	1.00	0.06	14,14,14,14	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around BEY A 1085:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.