Full wwPDB X-ray Structure Validation Report (i)

Jun 13, 2023 – 02:06 PM EDT

PDB ID	:	8F7V
Title	:	Macrocyclic plasmin inhibitor
Authors	:	Guojie, W.
Deposited on	:	2022-11-20
Resolution	:	1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.33
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.33

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1827 (1.66-1.66)
Clashscore	141614	1931 (1.66-1.66)
Ramachandran outliers	138981	1891 (1.66-1.66)
Sidechain outliers	138945	1891 (1.66-1.66)
RSRZ outliers	127900	1791 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	250	4% 90% 6% •
1	B	250	2% 83% 12% • •

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 4370 atoms, of which 100 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Plasminogen.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	242	1853	1175	327	337	14	0	1	1
1	B	242	1875	1193	332	334	16	0	3	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	741	ALA	SER	engineered mutation	UNP P00747
B	741	ALA	SER	engineered mutation	UNP P00747

- Molecule 2 is (6S,9R,19S,22R)-N-{[4-(aminomethyl)phenyl]methyl}-22-[(benzenesulfonyl)amino]-3,12,21-trioxo-2,6,9,13,20-pentaazatetracyclo[22.2.2.2 6,9 .2 14,17]dotriaconta-1(26),14,16,24,27,29-hexaene-19-carboxamide (three-letter code: EUA) (formula: $C_{42}H_{50}N_8O_6S$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	H	N	O	S		
2	A	1	107	42	50	8	6	1	0	0
2	B	1	107	42	50	8	6	1	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	O	S			
3	A	1	5	4	1		0	0
3	A	1	5	4	1		0	0
3	A	1	5	4	1		0	0
3	B	1	5	4	1		0	0
3	B	1	5	4	1		0	0
3	B	1	5	4	1		0	0
3	B	1	5	4	1		0	0
3	B	1	5	4	1		0	0
3	B	1	5	4	1		0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	B	1	6	3	3	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	151	151	151	0	0
			Total	O		
5	B	226	226	226	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Plasminogen



• Molecule 1: Plasminogen



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	48.68Å 50.19Å 92.02Å 90.00° 93.31° 90.00°	Depositor
Resolution (Å)	48.60 – 1.65 48.60 – 1.65	Depositor EDS
% Data completeness (in resolution range)	99.8 (48.60-1.65) 99.9 (48.60-1.65)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$< I/\sigma(I) >$ [1]	2.49 (at 1.65Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.173 , 0.187 0.171 , 0.185	Depositor DCC
R_{free} test set	2703 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å^2)	18.3	Xtriage
Anisotropy	0.205	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.35 , 47.1	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	4370	wwPDB-VP
Average B, all atoms (Å^2)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.33% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, EUA, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	1.43	3/1899 (0.2%)	0.90	0/2579
1	B	1.63	14/1931 (0.7%)	1.00	9/2618 (0.3%)
All	All	1.54	17/3830 (0.4%)	0.95	9/5197 (0.2%)

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	788[A]	MET	N-CA	9.91	1.66	1.46
1	B	788[B]	MET	N-CA	9.91	1.66	1.46
1	B	712[A]	ARG	C-N	-8.41	1.14	1.34
1	B	712[B]	ARG	C-N	-8.41	1.14	1.34
1	B	788[A]	MET	CA-C	6.43	1.69	1.52
1	B	788[B]	MET	CA-C	6.43	1.69	1.52
1	B	772	GLY	C-O	-6.06	1.14	1.23
1	B	594	SER	CB-OG	-5.91	1.34	1.42
1	B	712[A]	ARG	C-O	5.85	1.34	1.23
1	B	712[B]	ARG	C-O	5.85	1.34	1.23
1	A	679	GLU	CD-OE2	-5.76	1.19	1.25
1	B	743	GLY	C-O	-5.44	1.15	1.23
1	B	596	GLU	CD-OE2	-5.32	1.19	1.25
1	B	724	GLU	CD-OE1	-5.22	1.20	1.25
1	A	594	SER	CB-OG	-5.12	1.35	1.42
1	A	757	GLY	C-O	-5.09	1.15	1.23
1	B	641	GLU	CD-OE1	-5.05	1.20	1.25

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	712[A]	ARG	CA-C-N	9.15	137.33	117.20
1	B	712[B]	ARG	CA-C-N	9.15	137.33	117.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	712[A]	ARG	O-C-N	-8.19	109.59	122.70
1	B	712[B]	ARG	O-C-N	-8.19	109.59	122.70
1	B	676	ASP	CB-CG-OD1	5.62	123.36	118.30
1	B	712[A]	ARG	N-CA-C	5.33	125.40	111.00
1	B	712[B]	ARG	N-CA-C	5.33	125.40	111.00
1	B	610	ARG	NE-CZ-NH1	5.28	122.94	120.30
1	B	677	ARG	NE-CZ-NH1	-5.21	117.69	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1853	0	1828	9	0
1	B	1875	0	1868	17	0
2	A	57	50	0	0	0
2	B	57	50	0	0	0
3	A	15	0	0	0	0
3	B	30	0	0	0	0
4	B	6	0	8	0	0
5	A	151	0	0	2	0
5	B	226	0	0	2	0
All	All	4270	100	3704	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:577:VAL:HG13	1:B:616:VAL:HG13	1.53	0.91
1:B:606:GLU:OE1	1:B:607:LYS:HE2	1.84	0.77
1:B:577:VAL:CG1	1:B:616:VAL:HG13	2.16	0.75
1:B:597:TRP:CD1	1:B:788[B]:MET:HE3	2.25	0.71

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:623:GLU:HG2	1:B:685:TRP:CD1	2.38	0.57
1:A:577:VAL:HG22	1:A:590:GLY:C	2.29	0.54
1:A:596:GLU:HG2	5:A:1023:HOH:O	2.12	0.49
1:A:615:LYS:HE2	1:A:632:GLU:OE1	2.11	0.49
1:B:641:GLU:HB2	1:B:647:ILE:HG23	1.97	0.46
1:B:578:SER:HB3	1:B:617:ILE:HB	1.96	0.46
1:B:617:ILE:HG23	1:B:630:VAL:HG13	1.97	0.46
1:B:687:GLU:HG2	1:B:738:GLN:HB2	1.98	0.46
1:A:682:ILE:HD12	1:A:682:ILE:C	2.36	0.45
1:B:615:LYS:NZ	1:B:632:GLU:OE2	2.46	0.45
1:A:658:ILE:HD12	1:A:664:PRO:HD3	1.98	0.44
1:B:750:LYS:HE2	5:B:1086:HOH:O	2.17	0.43
1:B:610:ARG:NH1	5:B:906:HOH:O	2.47	0.43
1:B:577:VAL:HG22	1:B:618:LEU:CD2	2.48	0.43
1:A:732:GLY:HA2	5:A:953:HOH:O	2.19	0.42
1:B:682:ILE:C	1:B:682:ILE:HD12	2.40	0.42
1:A:555:PRO:CA	1:A:572:SER:HB2	2.50	0.42
1:B:617:ILE:HG23	1:B:630:VAL:CG1	2.50	0.41
1:A:682:ILE:HD12	1:A:682:ILE:O	2.21	0.40
1:A:759:THR:HA	1:A:774:TYR:CD2	2.56	0.40
1:B:658:ILE:HD12	1:B:664:PRO:HD3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	237/250 (95%)	226 (95%)	11 (5%)	0	100	100
1	B	237/250 (95%)	231 (98%)	6 (2%)	0	100	100
All	All	474/500 (95%)	457 (96%)	17 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	203/209 (97%)	201 (99%)	2 (1%)	76	62
1	B	206/209 (99%)	203 (98%)	3 (2%)	65	44
All	All	409/418 (98%)	404 (99%)	5 (1%)	71	53

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	615	LYS
1	A	628	PRO
1	B	547	ASP
1	B	623	GLU
1	B	637	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	625	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	EUA	B	801	-	62,62,62	2.68	22 (35%)	85,85,85	1.87	17 (20%)
3	SO4	B	803	-	4,4,4	0.15	0	6,6,6	0.23	0
3	SO4	A	802	-	4,4,4	0.13	0	6,6,6	0.08	0
3	SO4	A	803	-	4,4,4	0.13	0	6,6,6	0.08	0
3	SO4	B	807	-	4,4,4	0.60	0	6,6,6	0.37	0
4	GOL	B	808	-	5,5,5	0.83	0	5,5,5	1.10	1 (20%)
2	EUA	A	801	-	62,62,62	3.26	27 (43%)	85,85,85	1.75	19 (22%)
3	SO4	B	806	-	4,4,4	0.87	0	6,6,6	0.90	0
3	SO4	A	804	-	4,4,4	0.18	0	6,6,6	0.05	0
3	SO4	B	804	-	4,4,4	0.11	0	6,6,6	0.14	0
3	SO4	B	805	-	4,4,4	0.14	0	6,6,6	0.08	0
3	SO4	B	802	-	4,4,4	0.14	0	6,6,6	0.16	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EUA	B	801	-	-	12/56/66/66	0/5/6/6
2	EUA	A	801	-	-	10/56/66/66	0/5/6/6
4	GOL	B	808	-	-	0/4/4/4	-

All (49) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	EUA	O56-S49	-11.44	1.30	1.43
2	A	801	EUA	C41-N40	-8.24	1.28	1.47
2	B	801	EUA	C41-N40	-7.84	1.29	1.47
2	A	801	EUA	C36-N37	-7.19	1.30	1.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	EUA	O26-C15	-6.42	1.10	1.23
2	B	801	EUA	C15-N16	6.27	1.47	1.33
2	B	801	EUA	C36-N37	-5.98	1.33	1.47
2	A	801	EUA	C43-N40	-5.75	1.31	1.46
2	B	801	EUA	C34-N33	5.73	1.48	1.35
2	A	801	EUA	C15-N16	5.13	1.44	1.33
2	A	801	EUA	O57-S49	-4.82	1.38	1.43
2	A	801	EUA	C51-C50	-4.82	1.31	1.38
2	B	801	EUA	C50-S49	4.79	1.83	1.76
2	A	801	EUA	O01-C02	-4.76	1.13	1.23
2	B	801	EUA	C02-N03	4.72	1.44	1.34
2	B	801	EUA	C11-N10	4.62	1.45	1.35
2	B	801	EUA	O57-S49	4.59	1.48	1.43
2	A	801	EUA	C02-N03	4.59	1.44	1.34
2	A	801	EUA	O45-C34	-4.47	1.14	1.23
2	B	801	EUA	C36-C35	4.34	1.60	1.52
2	A	801	EUA	C38-N37	-4.18	1.35	1.46
2	A	801	EUA	C39-N40	-4.01	1.35	1.46
2	B	801	EUA	C17-C18	3.97	1.60	1.51
2	B	801	EUA	S49-N48	3.93	1.68	1.61
2	A	801	EUA	O12-C11	-3.93	1.15	1.23
2	B	801	EUA	C43-N40	-3.87	1.36	1.46
2	A	801	EUA	C27-N48	-3.84	1.39	1.46
2	A	801	EUA	C34-N33	3.61	1.43	1.35
2	A	801	EUA	C17-C18	3.59	1.59	1.51
2	A	801	EUA	C36-C35	3.55	1.59	1.52
2	B	801	EUA	C09-N10	3.55	1.48	1.41
2	A	801	EUA	C44-N37	-3.36	1.37	1.46
2	A	801	EUA	C09-N10	3.28	1.48	1.41
2	B	801	EUA	C38-N37	-3.17	1.38	1.46
2	B	801	EUA	O56-S49	3.09	1.47	1.43
2	B	801	EUA	C44-N37	-3.01	1.38	1.46
2	B	801	EUA	C39-N40	-2.99	1.38	1.46
2	A	801	EUA	C24-C21	-2.97	1.32	1.38
2	A	801	EUA	C11-N10	2.88	1.42	1.35
2	A	801	EUA	C19-C18	-2.85	1.32	1.38
2	A	801	EUA	C28-C29	2.65	1.57	1.51
2	B	801	EUA	C05-C06	2.52	1.57	1.51
2	A	801	EUA	C52-C51	-2.47	1.33	1.38
2	B	801	EUA	C32-N33	2.37	1.46	1.41
2	B	801	EUA	O26-C15	-2.29	1.18	1.23
2	B	801	EUA	C28-C29	2.28	1.56	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	EUA	C55-C50	-2.20	1.35	1.38
2	B	801	EUA	O12-C11	-2.20	1.18	1.23
2	A	801	EUA	S49-N48	2.02	1.65	1.61

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	B	801	EUA	O57-S49-O56	-9.87	107.42	119.55
2	A	801	EUA	O57-S49-O56	-6.57	111.48	119.55
2	B	801	EUA	C18-C17-N16	-5.53	101.20	113.05
2	A	801	EUA	C43-N40-C39	4.39	118.72	108.83
2	B	801	EUA	C27-N48-S49	-4.25	112.03	121.33
2	A	801	EUA	C27-N48-S49	-4.20	112.16	121.33
2	A	801	EUA	C18-C17-N16	-4.05	104.37	113.05
2	A	801	EUA	O57-S49-C50	3.40	112.15	107.97
2	B	801	EUA	C32-N33-C34	-3.36	121.62	127.50
2	B	801	EUA	C43-N40-C39	3.20	116.03	108.83
2	A	801	EUA	C54-C55-C50	-3.02	115.82	118.95
2	A	801	EUA	C51-C50-S49	-3.01	116.50	119.77
2	B	801	EUA	C50-S49-N48	2.90	111.80	107.78
2	A	801	EUA	C50-S49-N48	2.74	111.57	107.78
2	B	801	EUA	C28-C27-N48	2.72	113.88	110.59
2	A	801	EUA	O56-S49-C50	-2.68	104.66	107.97
2	B	801	EUA	C43-C44-N37	2.62	116.03	110.64
2	B	801	EUA	O57-S49-C50	2.57	111.13	107.97
2	B	801	EUA	C15-C04-N03	-2.52	104.29	111.16
2	A	801	EUA	O12-C11-N10	-2.38	119.29	123.63
2	B	801	EUA	C02-C27-N48	-2.33	104.90	111.17
2	A	801	EUA	C32-N33-C34	-2.31	123.45	127.50
2	B	801	EUA	C39-C38-N37	2.31	115.37	110.64
2	A	801	EUA	O45-C34-N33	-2.29	119.46	123.63
2	A	801	EUA	O57-S49-N48	2.25	110.96	106.88
2	A	801	EUA	C30-C31-C32	2.25	122.89	120.30
2	A	801	EUA	C41-N40-C39	2.23	116.95	111.23
2	B	801	EUA	O57-S49-N48	2.18	110.83	106.88
2	B	801	EUA	C24-C21-C20	2.18	121.59	118.17
4	B	808	GOL	C3-C2-C1	-2.12	103.46	111.70
2	A	801	EUA	O26-C15-C04	2.08	124.83	120.45
2	B	801	EUA	C05-C04-N03	-2.03	106.52	110.79
2	A	801	EUA	C31-C30-C29	-2.02	118.24	121.03
2	A	801	EUA	C52-C51-C50	2.02	121.05	118.95
2	B	801	EUA	C44-C43-N40	2.01	114.78	110.64

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	A	801	EUA	C09-N10-C11	-2.01	123.98	127.50
2	B	801	EUA	C09-N10-C11	-2.01	123.99	127.50

There are no chirality outliers.

All (22) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	801	EUA	C42-C41-N40-C39
2	B	801	EUA	C42-C41-N40-C43
2	A	801	EUA	C51-C50-S49-N48
2	A	801	EUA	C51-C50-S49-O56
2	A	801	EUA	C55-C50-S49-N48
2	A	801	EUA	C55-C50-S49-O56
2	B	801	EUA	C55-C50-S49-O56
2	B	801	EUA	C51-C50-S49-O56
2	B	801	EUA	C35-C36-N37-C44
2	B	801	EUA	O01-C02-C27-N48
2	B	801	EUA	N03-C02-C27-N48
2	A	801	EUA	O01-C02-C27-N48
2	A	801	EUA	N03-C02-C27-N48
2	B	801	EUA	O01-C02-C27-C28
2	A	801	EUA	O01-C02-C27-C28
2	B	801	EUA	N03-C02-C27-C28
2	B	801	EUA	C35-C36-N37-C38
2	B	801	EUA	C55-C50-S49-N48
2	A	801	EUA	N03-C02-C27-C28
2	B	801	EUA	C51-C50-S49-N48
2	A	801	EUA	C35-C36-N37-C44
2	A	801	EUA	C35-C36-N37-C38

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand EUA B 801



Ligand EUA A 801



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	B	2

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	561:ARG	C	562:VAL	N	19.20
1	B	712:ARG	C	713:TYR	N	1.14

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	242/250 (96%)	-0.14	10 (4%)	37	37	13, 25, 55, 83	0
1	B	242/250 (96%)	-0.32	5 (2%)	63	65	11, 17, 35, 56	0
All	All	484/500 (96%)	-0.23	15 (3%)	49	49	11, 20, 50, 83	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	692	PHE	7.2
1	A	688	THR	4.9
1	A	585	MET	4.4
1	B	688	THR	4.2
1	A	625	ASN	4.1
1	B	686	GLY	3.9
1	A	629	HIS	3.7
1	B	687	GLU	3.6
1	A	691	THR	3.5
1	A	628	PRO	3.5
1	A	627	GLU	3.1
1	B	629	HIS	3.0
1	A	690	GLY	2.7
1	A	626	LEU	2.3
1	B	628	PRO	2.3

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	GOL	B	808	6/6	0.84	0.15	22,27,33,35	0
3	SO4	B	804	5/5	0.91	0.14	37,53,60,65	0
3	SO4	B	805	5/5	0.93	0.21	40,48,66,73	0
3	SO4	B	807	5/5	0.94	0.14	38,41,48,48	0
2	EUA	B	801	57/57	0.94	0.10	14,23,43,62	0
3	SO4	A	802	5/5	0.95	0.11	50,56,67,69	0
3	SO4	B	806	5/5	0.95	0.15	49,53,55,67	0
2	EUA	A	801	57/57	0.96	0.08	13,22,33,35	0
3	SO4	A	804	5/5	0.97	0.15	30,40,41,41	0
3	SO4	B	803	5/5	0.98	0.10	29,31,36,37	0
3	SO4	A	803	5/5	0.98	0.13	42,48,58,60	0
3	SO4	B	802	5/5	0.99	0.07	29,32,38,42	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around EUA B 801:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around EUA A 801:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.