Full wwPDB X-ray Structure Validation Report (i)

Aug 27, 2023 – 02:37 PM EDT

PDB ID	:	3F8J
Title	:	Mouse UHRF1 SRA domain bound with hemi-methylated CpG, crystal structure in space group C222(1)
Authors	:	Hashimoto, H.; Horton, J.R.; Zhang, X.; Cheng, X.
Deposited on	:	2008-11-12
Resolution	:	1.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	212	5%, 88%, 8%
2	F	12	17%, 75%, 25%
3	G	12	8%, 50%, 50%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	B	191	-	X	-	-
4	GOL	B	192	-	X	-	X
4	GOL	B	193	-	X	-	-
4	GOL	B	194	-	X	-	X
4	GOL	B	195	-	X	-	-
4	GOL	B	196	-	X	-	X
4	GOL	B	197	-	X	-	-
4	GOL	B	198	-	X	-	-
4	GOL	B	199	-	X	-	X

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 2342 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called E3 ubiquitin-protein ligase UHRF1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	206	1640	1017	317	302	4	0	3	0

- Molecule 2 is a DNA chain called 5'-D(*DCP*DCP*DAP*DTP*DGP*(5CM)P*DGP*DCP*DTP*DGP*DAP*DC)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	F	12	241	116	44	70	11	0	0	0

- Molecule 3 is a DNA chain called 5'-D(*DGP*DTP*DCP*DAP*DGP*DCP*DGP*DCP*DAP*DTP*DGP*DG)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	G	12	246	117	48	70	11	0	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	B	1	6	3	3	0	0
			Total	C	O		
4	B	1	6	3	3	0	0
			Total	C	O		
4	B	1	6	3	3	0	0
			Total	C	O		
4	B	1	6	3	3	0	0
			Total	C	O		
4	B	1	6	3	3	0	0
			Total	C	O		
4	B	1	6	3	3	0	0
			Total	C	O		
4	B	1	6	3	3	0	0
			Total	C	O		
4	B	1	6	3	3	0	0
			Total	C	O		
4	B	1	6	3	3	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	B	123	123	123	0	0
			Total	O		
5	F	22	22	22	0	0
			Total	O		
5	G	16	16	16	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: E3 ubiquitin-protein ligase UHRF1

Chain B:



• Molecule 2: 5'-D(*DCP*DCP*DAP*DTP*DGP*(5CM)P*DGP*DCP*DTP*DGP*DAP*DC)-3'

Chain F:



• Molecule 3: 5'-D(*DGP*DTP*DCP*DAP*DGP*DCP*DGP*DCP*DAP*DTP*DGP*DG)-3'

Chain G:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α, β, γ	61.67Å 91.59Å 118.51Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.91 – 1.99 46.97 – 1.99	Depositor EDS
% Data completeness (in resolution range)	95.5 (29.91-1.99) 95.3 (46.97-1.99)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.47 (at 1.98Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.198 , 0.232 0.186 , 0.213	Depositor DCC
R_{free} test set	1127 reflections (4.80%)	wwPDB-VP
Wilson B-factor (Å^2)	22.8	Xtriage
Anisotropy	0.617	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.34 , 50.0	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	2342	wwPDB-VP
Average B, all atoms (Å^2)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.70% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, 5CM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	B	0.31	0/1682	0.57	0/2275
2	F	0.28	0/246	0.68	0/375
3	G	0.30	0/276	0.73	0/425
All	All	0.31	0/2204	0.61	0/3075

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	G	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	G	433	DG	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1640	0	1562	15	0
2	F	241	0	138	1	0
3	G	246	0	136	7	0
4	B	54	0	36	5	0
5	B	123	0	0	2	0
5	F	22	0	0	0	0
5	G	16	0	0	0	0
All	All	2342	0	1872	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:486:SER:H	1:B:510:ASN:HD21	1.17	0.91
1:B:505:LYS:HA	4:B:192:GOL:H11	1.64	0.78
3:G:424:DA:H2"	3:G:425:DG:H5'	1.73	0.69
3:G:424:DA:H2"	3:G:425:DG:C5'	2.28	0.64
3:G:422:DT:H2"	3:G:423:DC:O5'	2.00	0.61
1:B:523:GLU:HG3	5:B:157:HOH:O	1.99	0.61
1:B:460:ASP:HB2	4:B:199:GOL:H12	1.82	0.61
1:B:453:GLY:O	1:B:465:LEU:HA	2.09	0.53
1:B:486:SER:N	1:B:510:ASN:HD21	1.96	0.51
1:B:602:THR:HG23	1:B:607:LEU:HB2	1.93	0.51
1:B:458:SER:O	1:B:511[B]:ARG:NH1	2.43	0.50
1:B:535:LYS:HE2	5:B:30:HOH:O	2.12	0.48
2:F:412:DA:H2"	2:F:413:DC:O5'	2.14	0.47
1:B:570:PRO:HB2	1:B:578:LEU:HD11	1.96	0.46
3:G:421:DG:H2"	3:G:422:DT:H5'	1.98	0.45
3:G:421:DG:H2"	3:G:422:DT:C5'	2.46	0.45
3:G:422:DT:H5'	3:G:422:DT:H6	1.83	0.44
1:B:588:ASP:HA	4:B:194:GOL:H11	1.99	0.43
1:B:599:LYS:HG2	4:B:193:GOL:O2	2.19	0.42
1:B:439:VAL:O	1:B:443:GLU:HG3	2.20	0.42
1:B:531:TRP:NE1	1:B:532[A]:ARG:NH1	2.69	0.41
3:G:421:DG:C2'	3:G:422:DT:H71	2.51	0.41
1:B:505:LYS:HA	4:B:192:GOL:C1	2.44	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	B	207/212 (98%)	199 (96%)	7 (3%)	1 (0%)	29 23

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	570	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	B	165/173 (95%)	164 (99%)	1 (1%)	86 90

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	523	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	477	ASN
1	B	510	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	5CM	F	407	2	17,21,22	0.48	0	24,30,33	0.86	2 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	5CM	F	407	2	-	0/7/21/22	0/2/2/2

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	F	407	5CM	C5-C6-N1	-2.27	121.00	123.34
2	F	407	5CM	C5A-C5-C6	-2.16	119.97	122.85

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	GOL	B	199	-	5,5,5	4.52	5 (100%)	5,5,5	5.69	3 (60%)
4	GOL	B	191	-	5,5,5	4.56	5 (100%)	5,5,5	5.71	3 (60%)
4	GOL	B	193	-	5,5,5	4.56	5 (100%)	5,5,5	5.68	3 (60%)
4	GOL	B	192	-	5,5,5	4.52	5 (100%)	5,5,5	5.73	3 (60%)
4	GOL	B	197	-	5,5,5	4.51	5 (100%)	5,5,5	5.71	3 (60%)
4	GOL	B	195	-	5,5,5	4.50	5 (100%)	5,5,5	5.71	3 (60%)
4	GOL	B	196	-	5,5,5	4.51	5 (100%)	5,5,5	5.72	3 (60%)
4	GOL	B	194	-	5,5,5	4.51	5 (100%)	5,5,5	5.69	3 (60%)
4	GOL	B	198	-	5,5,5	4.57	5 (100%)	5,5,5	5.71	3 (60%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	B	199	-	-	2/4/4/4	-
4	GOL	B	191	-	-	3/4/4/4	-
4	GOL	B	193	-	-	3/4/4/4	-
4	GOL	B	192	-	-	4/4/4/4	-
4	GOL	B	197	-	-	2/4/4/4	-
4	GOL	B	195	-	-	3/4/4/4	-
4	GOL	B	196	-	-	2/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	B	194	-	-	2/4/4/4	-
4	GOL	B	198	-	-	2/4/4/4	-

All (45) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	191	GOL	C3-C2	-7.53	1.20	1.51
4	B	198	GOL	C3-C2	-7.51	1.20	1.51
4	B	199	GOL	C3-C2	-7.47	1.21	1.51
4	B	194	GOL	C3-C2	-7.47	1.21	1.51
4	B	193	GOL	C3-C2	-7.44	1.21	1.51
4	B	197	GOL	C3-C2	-7.40	1.21	1.51
4	B	196	GOL	C3-C2	-7.37	1.21	1.51
4	B	195	GOL	C3-C2	-7.37	1.21	1.51
4	B	192	GOL	C3-C2	-7.32	1.21	1.51
4	B	198	GOL	O1-C1	4.58	1.61	1.42
4	B	196	GOL	O1-C1	4.58	1.61	1.42
4	B	191	GOL	O1-C1	4.55	1.61	1.42
4	B	195	GOL	O1-C1	4.54	1.61	1.42
4	B	192	GOL	O1-C1	4.53	1.61	1.42
4	B	193	GOL	O1-C1	4.53	1.61	1.42
4	B	194	GOL	O1-C1	4.44	1.61	1.42
4	B	197	GOL	O1-C1	4.40	1.61	1.42
4	B	199	GOL	O1-C1	4.18	1.60	1.42
4	B	192	GOL	O3-C3	3.42	1.56	1.42
4	B	195	GOL	O3-C3	3.39	1.56	1.42
4	B	196	GOL	O3-C3	3.37	1.56	1.42
4	B	193	GOL	O3-C3	3.36	1.56	1.42
4	B	191	GOL	O3-C3	3.34	1.56	1.42
4	B	197	GOL	O3-C3	3.32	1.56	1.42
4	B	194	GOL	O3-C3	3.28	1.56	1.42
4	B	199	GOL	O3-C3	3.27	1.56	1.42
4	B	198	GOL	O3-C3	3.22	1.56	1.42
4	B	199	GOL	C1-C2	-3.05	1.39	1.51
4	B	199	GOL	O2-C2	-3.00	1.34	1.43
4	B	197	GOL	C1-C2	-2.92	1.39	1.51
4	B	193	GOL	O2-C2	-2.91	1.34	1.43
4	B	193	GOL	C1-C2	-2.89	1.39	1.51
4	B	198	GOL	O2-C2	-2.88	1.34	1.43
4	B	192	GOL	C1-C2	-2.88	1.39	1.51
4	B	194	GOL	C1-C2	-2.88	1.39	1.51
4	B	198	GOL	C1-C2	-2.87	1.39	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	192	GOL	O2-C2	-2.87	1.34	1.43
4	B	195	GOL	C1-C2	-2.79	1.40	1.51
4	B	197	GOL	O2-C2	-2.79	1.35	1.43
4	B	191	GOL	C1-C2	-2.78	1.40	1.51
4	B	191	GOL	O2-C2	-2.75	1.35	1.43
4	B	196	GOL	O2-C2	-2.74	1.35	1.43
4	B	194	GOL	O2-C2	-2.72	1.35	1.43
4	B	196	GOL	C1-C2	-2.71	1.40	1.51
4	B	195	GOL	O2-C2	-2.65	1.35	1.43

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
4	B	192	GOL	O3-C3-C2	10.50	160.56	110.20
4	B	198	GOL	O3-C3-C2	10.44	160.24	110.20
4	B	195	GOL	O3-C3-C2	10.43	160.23	110.20
4	B	196	GOL	O3-C3-C2	10.43	160.20	110.20
4	B	193	GOL	O3-C3-C2	10.40	160.07	110.20
4	B	197	GOL	O3-C3-C2	10.39	160.00	110.20
4	B	191	GOL	O3-C3-C2	10.34	159.80	110.20
4	B	199	GOL	O3-C3-C2	10.33	159.74	110.20
4	B	194	GOL	O3-C3-C2	10.32	159.68	110.20
4	B	197	GOL	O2-C2-C3	6.66	138.44	109.12
4	B	199	GOL	O2-C2-C3	6.65	138.40	109.12
4	B	191	GOL	O2-C2-C3	6.64	138.35	109.12
4	B	194	GOL	O2-C2-C3	6.63	138.32	109.12
4	B	196	GOL	O2-C2-C3	6.59	138.17	109.12
4	B	198	GOL	O2-C2-C3	6.54	137.94	109.12
4	B	195	GOL	O2-C2-C3	6.52	137.85	109.12
4	B	193	GOL	O2-C2-C3	6.49	137.72	109.12
4	B	192	GOL	O2-C2-C3	6.48	137.67	109.12
4	B	192	GOL	O1-C1-C2	3.36	126.30	110.20
4	B	196	GOL	O1-C1-C2	3.31	126.06	110.20
4	B	191	GOL	O1-C1-C2	3.28	125.94	110.20
4	B	195	GOL	O1-C1-C2	3.27	125.87	110.20
4	B	198	GOL	O1-C1-C2	3.26	125.82	110.20
4	B	199	GOL	O1-C1-C2	3.25	125.78	110.20
4	B	194	GOL	O1-C1-C2	3.25	125.77	110.20
4	B	193	GOL	O1-C1-C2	3.22	125.64	110.20
4	B	197	GOL	O1-C1-C2	3.15	125.31	110.20

There are no chirality outliers.

All (23) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	191	GOL	C1-C2-C3-O3
4	B	192	GOL	C1-C2-C3-O3
4	B	193	GOL	O1-C1-C2-C3
4	B	193	GOL	O2-C2-C3-O3
4	B	196	GOL	C1-C2-C3-O3
4	B	195	GOL	O1-C1-C2-O2
4	B	197	GOL	O1-C1-C2-O2
4	B	192	GOL	O1-C1-C2-C3
4	B	194	GOL	C1-C2-C3-O3
4	B	195	GOL	C1-C2-C3-O3
4	B	198	GOL	O1-C1-C2-C3
4	B	198	GOL	C1-C2-C3-O3
4	B	192	GOL	O1-C1-C2-O2
4	B	194	GOL	O1-C1-C2-O2
4	B	196	GOL	O1-C1-C2-O2
4	B	199	GOL	O1-C1-C2-O2
4	B	199	GOL	O2-C2-C3-O3
4	B	191	GOL	O2-C2-C3-O3
4	B	191	GOL	O1-C1-C2-O2
4	B	195	GOL	O1-C1-C2-C3
4	B	192	GOL	O2-C2-C3-O3
4	B	197	GOL	O2-C2-C3-O3
4	B	193	GOL	C1-C2-C3-O3

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	199	GOL	1	0
4	B	193	GOL	1	0
4	B	192	GOL	2	0
4	B	194	GOL	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	B	206/212 (97%)	0.20	10 (4%)	29	28	12, 21, 40, 74	0
2	F	11/12 (91%)	0.61	2 (18%)	1	1	22, 34, 64, 69	0
3	G	12/12 (100%)	0.53	1 (8%)	11	10	21, 33, 74, 80	0
All	All	229/236 (97%)	0.23	13 (5%)	23	23	12, 22, 58, 80	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	620	ALA	7.7
1	B	618	ALA	6.9
1	B	616	LEU	4.4
1	B	619	LEU	3.8
1	B	622	LYS	3.7
1	B	614	GLY	3.3
1	B	417	ILE	3.2
3	G	421	DG	3.2
2	F	413	DC	3.0
1	B	621	ASN	2.8
1	B	613	GLU	2.7
1	B	615	TYR	2.3
2	F	411	DG	2.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	5CM	F	407	20/21	0.97	0.10	11,16,19,21	0

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	GOL	B	192	6/6	0.29	0.53	71,72,72,74	0
4	GOL	B	194	6/6	0.41	0.46	56,60,61,64	0
4	GOL	B	196	6/6	0.57	0.52	62,65,66,67	0
4	GOL	B	199	6/6	0.61	0.45	45,48,50,52	0
4	GOL	B	198	6/6	0.65	0.36	61,64,64,65	0
4	GOL	B	197	6/6	0.73	0.39	39,45,48,49	0
4	GOL	B	191	6/6	0.74	0.23	70,71,71,71	0
4	GOL	B	195	6/6	0.74	0.39	60,62,63,64	0
4	GOL	B	193	6/6	0.79	0.29	54,55,56,58	0

6.5 Other polymers (i)

There are no such residues in this entry.