Full wwPDB X-ray Structure Validation Report (i)

Feb 6, 2024 – 01:26 PM EST

PDB ID	:	2FGR
Title	:	High resolution Xray structure of Omp32
Authors	:	Zeth, K.; Zachariae, U.; Engelhardt, H.
Deposited on	:	2005-12-22
Resolution	:	1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	332	2%, 87%, 10%, •
2	B	8	12%, 62%, 12%, 12%, 12%

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 2948 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Outer membrane porin protein 32.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	332	2462	1546	430	484	2	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLU	GLN	SEE REMARK 999	UNP P24305

- Molecule 2 is a protein called PAP.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	8	65	37	12	16	0	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
3	A	3	3	3	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	380	Total O 380 380	0	0
5	B	28	Total O 28 28	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Outer membrane porin protein 32

Chain A:



• Molecule 2: PAP

Chain B:

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α, β, γ	106.41Å 106.41Å 93.79Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 1.50 19.79 – 1.50	Depositor EDS
% Data completeness (in resolution range)	96.1 (20.00-1.50) 96.1 (19.79-1.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.30 (at 1.50Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.150 , 0.180 0.156 , 0.153	Depositor DCC
R_{free} test set	4646 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å^2)	15.0	Xtriage
Anisotropy	0.241	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.45 , 80.1	EDS
L-test for twinning[2]	$< \|L\| > = 0.48$, $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	0.039 for h,-h-k,-l	Xtriage
F_o,F_c correlation	0.97	EDS
Total number of atoms	2948	wwPDB-VP
Average B, all atoms (Å^2)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.56% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.82	4/2506 (0.2%)	1.02	13/3379 (0.4%)
2	B	1.08	0/66	1.08	1/88 (1.1%)
All	All	0.83	4/2572 (0.2%)	1.02	14/3467 (0.4%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	242	MET	SD-CE	-10.05	1.21	1.77
1	A	159	LYS	CE-NZ	7.10	1.66	1.49
1	A	329	ARG	CD-NE	-5.53	1.37	1.46
1	A	258	GLN	CB-CG	-5.12	1.38	1.52

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	329	ARG	NE-CZ-NH1	17.34	128.97	120.30
1	A	329	ARG	NE-CZ-NH2	-17.22	111.69	120.30
1	A	89	ARG	NE-CZ-NH2	-7.96	116.32	120.30
1	A	299	ASP	CB-CG-OD2	7.46	125.01	118.30
1	A	329	ARG	CD-NE-CZ	6.50	132.70	123.60
1	A	268	ASP	CB-CG-OD2	5.91	123.62	118.30
1	A	92	ARG	NE-CZ-NH1	5.61	123.11	120.30
2	B	401	ASP	CB-CG-OD2	5.40	123.16	118.30
1	A	42	ARG	NE-CZ-NH2	-5.30	117.65	120.30
1	A	257	LEU	CB-CG-CD2	5.20	119.84	111.00
1	A	106	LEU	CB-CG-CD1	5.11	119.68	111.00
1	A	105	ASP	CB-CG-OD1	5.06	122.85	118.30
1	A	65	ASP	CB-CG-OD1	5.06	122.85	118.30
1	A	329	ARG	CG-CD-NE	-5.06	101.18	111.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2462	0	2398	49	0
2	B	65	0	48	7	0
3	A	3	0	0	0	0
4	A	10	0	0	0	0
5	A	380	0	0	19	0
5	B	28	0	0	3	0
All	All	2948	0	2446	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:252:VAL:N	5:A:931:HOH:O	1.78	1.13
1:A:161:THR:HB	5:A:917:HOH:O	0.92	1.09
1:A:101:LEU:HG	5:A:894:HOH:O	0.93	1.09
1:A:242:MET:HE3	1:A:258:GLN:CG	1.84	1.07
1:A:242:MET:CE	1:A:258:GLN:HG3	1.86	1.06
1:A:242:MET:HE3	1:A:258:GLN:CD	1.80	1.02
1:A:305:LEU:H	1:A:321:GLN:HE22	1.15	0.94
2:B:401:ASP:HB3	5:B:319:HOH:O	1.69	0.92
2:B:401:ASP:N	5:B:312:HOH:O	2.05	0.88
1:A:242:MET:CE	1:A:258:GLN:CG	2.48	0.86
1:A:281:ASN:OD1	5:A:911:HOH:O	1.95	0.84
1:A:101:LEU:CD1	1:A:207:THR:HB	2.08	0.82
1:A:101:LEU:HD13	1:A:207:THR:CG2	2.10	0.82
1:A:101:LEU:HD13	1:A:207:THR:HB	1.62	0.81
1:A:242:MET:HE2	1:A:258:GLN:HG3	1.62	0.81
1:A:329:ARG:HD2	5:A:657:HOH:O	1.79	0.81
1:A:101:LEU:HD13	1:A:207:THR:CB	2.12	0.80

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:329:ARG:CD	5:A:657:HOH:O	2.33	0.77
1:A:251:GLY:O	1:A:252:VAL:HB	1.85	0.75
1:A:251:GLY:CA	5:A:931:HOH:O	2.34	0.75
1:A:159:LYS:O	1:A:161:THR:HG23	1.90	0.71
1:A:305:LEU:H	1:A:321:GLN:NE2	1.89	0.70
1:A:250:GLY:O	1:A:251:GLY:O	2.10	0.69
1:A:106:LEU:H	1:A:258:GLN:HE22	1.40	0.67
5:A:965:HOH:O	2:B:407:THR:HG23	1.95	0.66
1:A:44:THR:HB	5:A:905:HOH:O	1.97	0.64
1:A:101:LEU:HD13	1:A:207:THR:HG21	1.79	0.64
2:B:407:THR:O	2:B:408:SER:OG	2.21	0.59
1:A:101:LEU:HD11	1:A:207:THR:HB	1.85	0.59
1:A:101:LEU:CG	5:A:894:HOH:O	1.78	0.59
1:A:286:THR:OG1	1:A:330:HIS:HD2	1.86	0.58
1:A:18:ASN:ND2	1:A:322:THR:H	2.04	0.55
1:A:161:THR:HG22	1:A:166:ASP:HA	1.89	0.54
1:A:242:MET:HE3	1:A:258:GLN:NE2	2.23	0.53
1:A:251:GLY:C	5:A:931:HOH:O	2.20	0.53
1:A:113:GLY:HA3	1:A:275:THR:HG21	1.90	0.52
2:B:401:ASP:N	2:B:401:ASP:OD1	2.43	0.52
1:A:242:MET:CE	1:A:258:GLN:CD	2.69	0.52
1:A:113:GLY:CA	1:A:275:THR:HG21	2.41	0.50
1:A:84:ASN:HD22	1:A:85:PHE:N	2.11	0.49
1:A:316:GLN:HB2	1:A:319:GLU:HG3	1.93	0.49
1:A:84:ASN:HD22	1:A:84:ASN:C	2.16	0.48
2:B:407:THR:C	2:B:408:SER:OG	2.52	0.48
1:A:161:THR:CB	5:A:917:HOH:O	1.81	0.48
2:B:401:ASP:N	5:B:192:HOH:O	2.48	0.47
1:A:116:MET:HE1	5:A:658:HOH:O	2.16	0.45
1:A:329:ARG:HD3	5:A:657:HOH:O	2.05	0.45
1:A:242:MET:CE	1:A:258:GLN:NE2	2.80	0.45
1:A:116:MET:CE	5:A:658:HOH:O	2.65	0.44
1:A:242:MET:HE2	1:A:243:LEU:C	2.37	0.44
1:A:251:GLY:HA2	5:A:931:HOH:O	2.06	0.44
1:A:189:LEU:HB3	5:A:902:HOH:O	2.17	0.44
1:A:101:LEU:CD1	5:A:894:HOH:O	2.31	0.43
1:A:189:LEU:HD22	5:A:902:HOH:O	2.19	0.42
1:A:18:ASN:HD21	1:A:321:GLN:HA	1.85	0.41
1:A:170:ARG:O	1:A:192:GLN:HA	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	330/332 (99%)	322 (98%)	6 (2%)	2 (1%)	25 7
2	B	6/8 (75%)	6 (100%)	0	0	100 100
All	All	336/340 (99%)	328 (98%)	6 (2%)	2 (1%)	25 7

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	251	GLY
1	A	252	VAL

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	245/245 (100%)	237 (97%)	8 (3%)	38 10
2	B	7/7 (100%)	5 (71%)	2 (29%)	0 0
All	All	252/252 (100%)	242 (96%)	10 (4%)	31 6

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	ASN
1	A	81	LEU
1	A	84	ASN
1	A	90	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	106	LEU
1	A	130	ASN
1	A	257	LEU
1	A	288	LEU
2	B	401	ASP
2	B	408	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	18	ASN
1	A	84	ASN
1	A	130	ASN
1	A	135	ASN
1	A	160	GLN
1	A	191	GLN
1	A	224	GLN
1	A	258	GLN
1	A	264	GLN
1	A	291	ASN
1	A	297	ASN
1	A	306	GLN
1	A	316	GLN
1	A	321	GLN
1	A	330	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	SO4	A	601	-	4,4,4	0.24	0	6,6,6	0.40	0
4	SO4	A	600	-	4,4,4	0.34	0	6,6,6	0.35	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	332/332 (100%)	-0.14	6 (1%) 68 73	10, 17, 26, 42	0
2	B	8/8 (100%)	0.62	1 (12%) 3 3	13, 21, 29, 32	0
All	All	340/340 (100%)	-0.12	7 (2%) 63 68	10, 17, 27, 42	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	250	GLY	4.4
1	A	251	GLY	4.3
1	A	1	GLU	3.4
2	B	408	SER	2.8
1	A	27	TYR	2.2
1	A	216	VAL	2.2
1	A	284	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	SO4	A	601	5/5	0.95	0.29	45,49,51,53	0
4	SO4	A	600	5/5	0.97	0.15	34,34,38,40	0
3	CA	A	503	1/1	1.00	0.05	22,22,22,22	0
3	CA	A	501	1/1	1.00	0.04	12,12,12,12	0
3	CA	A	502	1/1	1.00	0.15	34,34,34,34	1

6.5 Other polymers (i)

There are no such residues in this entry.