wwPDB X-ray Structure Validation Summary Report (i)

Jun 19, 2024 – 06:41 AM EDT

PDB ID	:	4FGI
Title	:	Structure of the effector - immunity system Tse1 / Tsi1 from Pseudomonas aeruginosa
Authors	:	Benz, J.; Sendlmeier, C.; Barends, T.R.M.; Meinhart, A.
Deposited on	:	2012-06-04
Resolution	:	3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1133 (3.20-3.20)
Clashscore	141614	1253 (3.20-3.20)
Ramachandran outliers	138981	1234 (3.20-3.20)
Sidechain outliers	138945	1233 (3.20-3.20)
RSRZ outliers	127900	1095 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	165	% 82% 7% 11%
1	C	165	79% 9% 12%
1	E	165	79% 10% 12%
1	G	165	2% 80% 7% • 12%
2	B	151	81% 15% • •

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	151	87% 9%
2	F	151	78% 19%
2	H	151	% 81% 14%

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 8893 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tse1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	147	1097	685	196	207	6	3	0	0	0
1	C	146	1091	682	195	205	6	3	0	0	0
1	E	146	1091	682	195	205	6	3	0	0	0
1	G	145	1085	679	194	203	6	3	0	0	0

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	155	ALA	-	EXPRESSION TAG	UNP Q9I2Q1
A	156	ALA	-	EXPRESSION TAG	UNP Q9I2Q1
A	157	ALA	-	EXPRESSION TAG	UNP Q9I2Q1
A	158	LEU	-	EXPRESSION TAG	UNP Q9I2Q1
A	159	GLU	-	EXPRESSION TAG	UNP Q9I2Q1
A	160	HIS	-	EXPRESSION TAG	UNP Q9I2Q1
A	161	HIS	-	EXPRESSION TAG	UNP Q9I2Q1
A	162	HIS	-	EXPRESSION TAG	UNP Q9I2Q1
A	163	HIS	-	EXPRESSION TAG	UNP Q9I2Q1
A	164	HIS	-	EXPRESSION TAG	UNP Q9I2Q1
A	165	HIS	-	EXPRESSION TAG	UNP Q9I2Q1
C	155	ALA	-	EXPRESSION TAG	UNP Q9I2Q1
C	156	ALA	-	EXPRESSION TAG	UNP Q9I2Q1
C	157	ALA	-	EXPRESSION TAG	UNP Q9I2Q1
C	158	LEU	-	EXPRESSION TAG	UNP Q9I2Q1
C	159	GLU	-	EXPRESSION TAG	UNP Q9I2Q1
C	160	HIS	-	EXPRESSION TAG	UNP Q9I2Q1
C	161	HIS	-	EXPRESSION TAG	UNP Q9I2Q1
C	162	HIS	-	EXPRESSION TAG	UNP Q9I2Q1
C	163	HIS	-	EXPRESSION TAG	UNP Q9I2Q1
C	164	HIS	-	EXPRESSION TAG	UNP Q9I2Q1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	165	HIS	-	EXPRESSION TAG	UNP Q9I2Q1
E	155	ALA	-	EXPRESSION TAG	UNP Q9I2Q1
E	156	ALA	-	EXPRESSION TAG	UNP Q9I2Q1
E	157	ALA	-	EXPRESSION TAG	UNP Q9I2Q1
E	158	LEU	-	EXPRESSION TAG	UNP Q9I2Q1
E	159	GLU	-	EXPRESSION TAG	UNP Q9I2Q1
E	160	HIS	-	EXPRESSION TAG	UNP Q9I2Q1
E	161	HIS	-	EXPRESSION TAG	UNP Q9I2Q1
E	162	HIS	-	EXPRESSION TAG	UNP Q9I2Q1
E	163	HIS	-	EXPRESSION TAG	UNP Q9I2Q1
E	164	HIS	-	EXPRESSION TAG	UNP Q9I2Q1
E	165	HIS	-	EXPRESSION TAG	UNP Q9I2Q1
G	155	ALA	-	EXPRESSION TAG	UNP Q9I2Q1
G	156	ALA	-	EXPRESSION TAG	UNP Q9I2Q1
G	157	ALA	-	EXPRESSION TAG	UNP Q9I2Q1
G	158	LEU	-	EXPRESSION TAG	UNP Q9I2Q1
G	159	GLU	-	EXPRESSION TAG	UNP Q9I2Q1
G	160	HIS	-	EXPRESSION TAG	UNP Q9I2Q1
G	161	HIS	-	EXPRESSION TAG	UNP Q9I2Q1
G	162	HIS	-	EXPRESSION TAG	UNP Q9I2Q1
G	163	HIS	-	EXPRESSION TAG	UNP Q9I2Q1
G	164	HIS	-	EXPRESSION TAG	UNP Q9I2Q1
G	165	HIS	-	EXPRESSION TAG	UNP Q9I2Q1

- Molecule 2 is a protein called Tsi1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
2	B	147	1134	697	201	228	5	3	0	0	0
			Total	C	N	O	S	Se			
2	D	147	1134	697	201	228	5	3	0	0	0
			Total	C	N	O	S	Se			
2	F	147	1134	697	201	228	5	3	0	0	0
			Total	C	N	O	S	Se			
2	H	146	1127	693	200	226	5	3	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	22	MSE	-	EXPRESSION TAG	UNP Q9I2Q0
B	23	ALA	-	EXPRESSION TAG	UNP Q9I2Q0
D	22	MSE	-	EXPRESSION TAG	UNP Q9I2Q0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	23	ALA	-	EXPRESSION TAG	UNP Q9I2Q0
F	22	MSE	-	EXPRESSION TAG	UNP Q9I2Q0
F	23	ALA	-	EXPRESSION TAG	UNP Q9I2Q0
H	22	MSE	-	EXPRESSION TAG	UNP Q9I2Q0
H	23	ALA	-	EXPRESSION TAG	UNP Q9I2Q0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tse1

Chain A:



- Molecule 1: Tse1

Chain C:



- Molecule 1: Tse1

Chain E:

- Molecule 1: Tse1

Chain G:



- Molecule 2: Tsi1

Chain B:

- Molecule 2: Tsi1

Chain D:



• Molecule 2: Tsi1

Chain F:



• Molecule 2: Tsi1

Chain H:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 41 2 2	Depositor
Cell constants a, b, c, α, β, γ	97.56Å 97.56Å 423.74Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.33 – 3.20 49.35 – 3.20	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.33-3.20) 99.9 (49.35-3.20)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	5.50 (at 3.19Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.224 , 0.263 0.221 , 0.261	Depositor DCC
R_{free} test set	1753 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å^2)	62.6	Xtriage
Anisotropy	0.034	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.28 , 20.6	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.91	EDS
Total number of atoms	8893	wwPDB-VP
Average B, all atoms (Å^2)	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.48% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.32	0/1116	0.47	0/1507
1	C	0.32	0/1110	0.48	0/1499
1	E	0.33	0/1110	0.48	0/1499
1	G	0.32	0/1104	0.48	0/1491
2	B	0.37	0/1155	0.55	0/1555
2	D	0.37	0/1155	0.51	0/1555
2	F	0.34	0/1155	0.51	0/1555
2	H	0.37	0/1148	0.55	0/1545
All	All	0.35	0/9053	0.50	0/12206

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1097	0	1074	7	0
1	C	1091	0	1069	6	0
1	E	1091	0	1069	8	0
1	G	1085	0	1064	6	0
2	B	1134	0	1056	16	0
2	D	1134	0	1056	8	0
2	F	1134	0	1056	15	0
2	H	1127	0	1049	17	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	8893	0	8493	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 73 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:23:ALA:N	2:D:167:CYS:HG	1.55	1.04
2:F:23:ALA:N	2:F:167:CYS:HG	1.74	0.86
2:B:23:ALA:N	2:B:167:CYS:HG	1.82	0.78
1:E:87:ARG:HH11	2:H:99:ARG:HE	1.40	0.70
2:H:99:ARG:HD2	2:H:100:ARG:HH12	1.58	0.69

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	145/165 (88%)	141 (97%)	4 (3%)	0	100	100
1	C	144/165 (87%)	136 (94%)	8 (6%)	0	100	100
1	E	144/165 (87%)	137 (95%)	7 (5%)	0	100	100
1	G	143/165 (87%)	137 (96%)	6 (4%)	0	100	100
2	B	145/151 (96%)	140 (97%)	5 (3%)	0	100	100
2	D	145/151 (96%)	139 (96%)	5 (3%)	1 (1%)	22	61
2	F	145/151 (96%)	138 (95%)	7 (5%)	0	100	100
2	H	144/151 (95%)	138 (96%)	6 (4%)	0	100	100
All	All	1155/1264 (91%)	1106 (96%)	48 (4%)	1 (0%)	51	83

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	168	GLN

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	114/124 (92%)	111 (97%)	3 (3%)	46	76
1	C	113/124 (91%)	108 (96%)	5 (4%)	28	64
1	E	113/124 (91%)	104 (92%)	9 (8%)	12	42
1	G	112/124 (90%)	106 (95%)	6 (5%)	22	58
2	B	125/124 (101%)	119 (95%)	6 (5%)	25	61
2	D	125/124 (101%)	123 (98%)	2 (2%)	62	84
2	F	125/124 (101%)	117 (94%)	8 (6%)	17	52
2	H	124/124 (100%)	117 (94%)	7 (6%)	21	57
All	All	951/992 (96%)	905 (95%)	46 (5%)	25	61

5 of 46 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	70	LEU
1	G	7	CYS
2	F	122	LYS
2	F	150	GLU
1	G	87	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	145	GLN
1	C	6	GLN
2	F	145	GLN
1	G	6	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	144/165 (87%)	0.05	1 (0%)	87	81	46, 73, 102, 117	1 (0%)
1	C	143/165 (86%)	-0.27	0	100	100	41, 67, 95, 123	1 (0%)
1	E	143/165 (86%)	-0.13	0	100	100	53, 89, 132, 157	1 (0%)
1	G	142/165 (86%)	0.12	3 (2%)	63	49	53, 84, 122, 130	1 (0%)
2	B	144/151 (95%)	-0.16	0	100	100	40, 56, 81, 91	0
2	D	144/151 (95%)	-0.10	0	100	100	37, 57, 87, 99	0
2	F	144/151 (95%)	-0.05	0	100	100	42, 65, 106, 127	0
2	H	143/151 (94%)	0.06	2 (1%)	75	63	48, 76, 104, 117	0
All	All	1147/1264 (90%)	-0.06	6 (0%)	91	86	37, 70, 115, 157	4 (0%)

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	14	ASN	3.3
2	H	44	GLU	3.0
2	H	34	GLY	2.8
1	G	143	TYR	2.3
1	A	143	TYR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.