wwPDB NMR Structure Validation Summary Report (i)

Feb 12, 2022 – 06:39 PM EST

PDB ID	:	1FSH
Title	:	STRUCTURAL BASIS OF THE RECOGNITION OF THE DISHEVELLED DEP DOMAIN IN THE WNT SIGNALING PATHWAY
Authors	:	Wong, H.C.; Zheng, J.
Deposited on	:	2000-09-08

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	2.26
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.26

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	105	13% 54% 17% • • 10%

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 10 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest target function*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:14-A:105 (92)	0.34	10

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 4, 6, 7, 8, 10, 11, 12, 14, 16, 17, 18, 19, 20
2	5, 9
3	13, 15

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1538 atoms, of which 776 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called DISHEVELLED-1.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	94	1538	487	776	133	137	5	0

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: DISHEVELLED-1

Chain A: 13% 54% 17% • • 10%

THR SER SER VAL PRO GLY ALA PRO GLN LEU GLU E12 A13 P14 L15 T16 V17 K18 S19 D20 M21 S22 A23 I24 V25 R26 V27 M28 Q29 L30 S33 G34 L35 E36 I37 R38 D39 R40 M41 W42 L43 K44 I45 T46 I47 A48 N49 A50 V51 I52 G53 A54 D55 V56 V57 D58 W59 L60 Y61

T62 H63 V64 E65 G66 F67 K68 E69 R70 R71 E72 Y76 A77 S78 S79 M80 L81 K82 H83 G84 F85 L86 R87 H88 T89 V90 N91 K92 I93 T94 F95 S96 E97 Q98 C99 Y100 Y101 V102 F103 G104 D105

4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 10. Colouring as in section 4.1 above.

- Molecule 1: DISHEVELLED-1

Chain A: 11% 52% 21% • • 10%

THR SER SER VAL PRO GLY ALA PRO GLN LEU GLU E12 A13 P14 L15 T16 V17 K18 S19 D20 M21 S22 A23 I24 V25 R26 V27 L30 P31 D32 S33 G34 L35 E36 I37 R38 D39 R40 M41 W42 L43 K44 I45 A48 N49 A50 V51 I52 G53 A54 D55 V56 V57 D58 W59 L60 Y61 T62

H63 V64 E65 G66 F67 K68 E69 R70 R71 E72 K75 Y76 A77 S78 S79 M80 L81 K82 H83 G84 F85 L86 R87 H88 T89 V90 N91 K92 I93 T94 F95 S96 E97 Q98 C99 Y100 Y101 V102 F103 G104 D105

5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *torsion angle dynamics*.

Of the 1000 calculated structures, 20 were deposited, based on the following criterion: *target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
DYANA	refinement	1.5
DYANA	structure solution	1.5

No chemical shift data was provided.

6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	748	764	753	172±15
All	All	14960	15280	15054	3444

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 115.

5 of 803 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:28:MET:CE	1:A:56:VAL:HG13	1.67	1.20	19	19
1:A:17:VAL:HG22	1:A:67:PHE:CE1	1.66	1.10	2	3
1:A:28:MET:HE1	1:A:56:VAL:CG1	1.60	1.25	4	19
1:A:24:ILE:HG13	1:A:76:TYR:CE2	1.60	1.29	12	10
1:A:33:SER:HB2	1:A:59:TRP:CZ2	1.59	1.26	5	13

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	91/105 (87%)	56±2 (61±2%)	24±2 (27±3%)	11±1 (12±1%)	1 7
All	All	1820/2100 (87%)	1113 (61%)	488 (27%)	219 (12%)	1 7

5 of 26 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	17	VAL	20
1	A	38	ARG	20
1	A	49	ASN	20
1	A	51	VAL	20
1	A	52	ILE	20

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	81/91 (89%)	54±3 (67±4%)	27±3 (33±4%)	1 12
All	All	1620/1820 (89%)	1088 (67%)	532 (33%)	1 12

5 of 57 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	16	THR	20
1	A	43	LEU	20
1	A	51	VAL	20
1	A	82	LYS	20
1	A	20	ASP	19

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

No chemical shift data were provided