wwPDB X-ray Structure Validation Summary Report (i)

Apr 20, 2024 – 01:33 pm BST

PDB ID	:	6FX1
Title	:	Crystal structure of Pholiota squarrosa lectin in complex with an octasaccharide
Authors	:	Cabanettes, A.; Varrot, A.
Deposited on	:	2018-03-08
Resolution	:	2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	43	2%, 86%, 7%, 7%
1	B	43	5%, 95%, • •
1	C	43	88%, 5%, 7%
1	D	43	86%, 5%, 9%
1	E	43	5%, 93%, 5%, •

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	43	2% 93% 5%
1	G	43	2% 81% 12% 7%
1	H	43	2% 93% 7%
1	I	43	2% 98%
1	J	43	5% 84% 9% 7%
1	K	43	7% 91% 9%
1	L	43	2% 86% 9% 5%
2	M	6	100%
2	O	6	33% 67%
2	R	6	50% 50%
2	X	6	17% 83%
3	N	8	12% 88%
3	U	8	50% 50%
4	P	5	20% 80%
5	Q	7	29% 71%
5	S	7	14% 86%
6	T	4	25% 75%
6	V	4	75% 25%
6	W	4	100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	MLI	D	108	-	X	-	-

2 Entry composition (i)

There are 9 unique types of molecules in this entry. The entry contains 5109 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called lectin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	40	314	201	51	60	2	0	0	0
1	B	42	322	207	50	62	3	0	0	0
1	C	40	314	201	51	60	2	0	0	0
1	D	39	309	199	50	58	2	0	0	0
1	E	42	330	211	52	64	3	0	1	0
1	F	41	322	206	52	61	3	0	0	0
1	G	40	314	201	51	60	2	0	0	0
1	H	40	322	205	53	62	2	0	1	0
1	I	43	328	210	52	63	3	0	0	0
1	J	40	309	199	48	60	2	0	0	0
1	K	39	301	195	46	58	2	0	0	0
1	L	41	318	206	49	60	3	0	0	0

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]1-azido-beta-N-acetyl-D-glucosamine.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	M	6	75	42	4	29	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	O	6	77	42	6	29	0	0	0
2	R	6	77	42	6	29	0	0	0
2	X	6	75	42	4	29	0	0	0

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]1-azido-beta-N-acetyl-D-glucosamine.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	N	8	100	56	5	39	0	0	0
3	U	8	100	56	5	39	0	0	0

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]1-azido-beta-N-acetyl-D-glucosamine.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	P	5	61	34	3	24	0	0	0

- Molecule 5 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]1-azido-beta-N-acetyl-D-glucosamine.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
5	Q	7	86	48	4	34	0	0	0
5	S	7	88	48	6	34	0	0	0

- Molecule 6 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]1-azido-beta-N-acetyl-D-glucosamine.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
6	T	4	50	28	3	19	0	0	0
			Total	C	N	O			
6	V	4	50	28	3	19	0	0	0
			Total	C	N	O			
6	W	4	50	28	3	19	0	0	0

- Molecule 7 is MALONATE ION (three-letter code: MLI) (formula: $C_3H_2O_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O			
7	A	1	7	3	4		0	0
			Total	C	O			
7	B	1	7	3	4		0	0
			Total	C	O			
7	C	1	7	3	4		0	0
			Total	C	O			
7	D	1	7	3	4		0	0
			Total	C	O			
7	D	1	7	3	4		0	0
			Total	C	O			
7	D	1	7	3	4		0	0
			Total	C	O			
7	E	1	7	3	4		0	0
			Total	C	O			
7	F	1	7	3	4		0	0
			Total	C	O			
7	H	1	7	3	4		0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	H	1	Total C O 7 3 4	0	0
7	I	1	Total C O 7 3 4	0	0
7	K	1	Total C O 7 3 4	0	0

- Molecule 8 is 3-OXOOCTANOIC ACID (three-letter code: OOA) (formula: $C_8H_{14}O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total C O 11 8 3	0	0

- Molecule 9 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	27	Total O 27 27	0	0
9	B	22	Total O 22 22	0	0
9	C	29	Total O 31 31	0	2
9	D	36	Total O 36 36	0	0
9	E	32	Total O 33 33	0	1

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
9	F	34	35	35	0	1
			Total	O		
9	G	34	34	34	0	0
			Total	O		
9	H	25	26	26	0	1
			Total	O		
9	I	34	35	35	0	1
			Total	O		
9	J	13	13	13	0	0
			Total	O		
9	K	15	15	15	0	0
			Total	O		
9	L	15	15	15	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: lectin

Chain A:

• Molecule 1: lectin

Chain B:





• Molecule 1: lectin

Chain C:

• Molecule 1: lectin

Chain D:



• Molecule 1: lectin

Chain E:





• Molecule 1: lectin

Chain F:





• Molecule 1: lectin

Chain G:



• Molecule 1: lectin

Chain H:





• Molecule 1: lectin

Chain I:



• Molecule 1: lectin

Chain J:



• Molecule 1: lectin

Chain K:





• Molecule 1: lectin

Chain L:



• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]1-azido-beta-N-acetyl-D-glucosamine

Chain M: 100%

• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]1-azido-beta-N-acetyl-D-glucosamine

Chain O: 33% 67%

• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]1-azido-beta-N-acetyl-D-glucosamine

Chain R: 50% 50%



• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]1-azido-beta-N-acetyl-D-glucosamine

Chain X: 17% 83%

• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]1-azido-beta-N-acetyl-D-glucosamine

Chain N: 12% 88%

• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]1-azido-beta-N-acetyl-D-glucosamine

Chain U: 50% 50%



• Molecule 4: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]1-azido-beta-N-acetyl-D-glucosamine

Chain P: 20% 80%

C4W1 NAG2 BMA3 MAN4 FUC5

• Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]1-azido-beta-N-acetyl-D-glucosamine

Chain Q: 29% 71%

C4W1 NAG2 BMA3 MAN4 NAG5 MAN6 FUC7

• Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]1-azido-beta-N-acetyl-D-glucosamine

Chain S: 14% 86%

C4W1 NAG2 BMA3 MAN4 NAG5 MAN6 FUC7

• Molecule 6: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]1-azido-beta-N-acetyl-D-glucosamine

Chain T: 25% 75%

C4W1 NAG2 BMA3 FUC4

• Molecule 6: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]1-azido-beta-N-acetyl-D-glucosamine

Chain V: 75% 25%

C4W1 NAG2 BMA3 FUC4

• Molecule 6: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]1-azido-beta-N-acetyl-D-glucosamine

Chain W: 100%

C4W1 NAG2 BMA3 FUC4

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 2 21	Depositor
Cell constants a, b, c, α, β, γ	91.00Å 95.59Å 110.63Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.88 – 2.10 47.88 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.9 (47.88-2.10) 99.9 (47.88-2.10)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.08 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.184 , 0.219 0.191 , 0.222	Depositor DCC
R_{free} test set	2785 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å^2)	40.1	Xtriage
Anisotropy	0.069	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.37 , 37.7	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.013 for k,h,-l	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	5109	wwPDB-VP
Average B, all atoms (Å^2)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 15.12% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, FUC, C4W, OOA, MAN, BMA, MLI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.80	0/324	0.92	1/443 (0.2%)
1	B	0.70	0/332	0.81	0/454
1	C	0.80	0/324	0.96	3/443 (0.7%)
1	D	0.86	0/319	1.02	2/438 (0.5%)
1	E	0.79	0/340	0.82	0/465
1	F	0.71	0/332	0.91	1/453 (0.2%)
1	G	0.88	0/324	1.11	4/443 (0.9%)
1	H	0.85	0/332	0.83	0/454
1	I	0.80	0/338	0.83	0/461
1	J	0.82	0/319	0.97	3/437 (0.7%)
1	K	0.76	0/311	0.77	0/429
1	L	0.74	0/328	0.81	0/450
All	All	0.79	0/3923	0.90	14/5370 (0.3%)

There are no bond length outliers.

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	D	27	ARG	NE-CZ-NH1	9.56	125.08	120.30
1	J	33	ASP	CB-CG-OD1	9.34	126.70	118.30
1	G	27	ARG	NE-CZ-NH1	-8.78	115.91	120.30
1	G	27	ARG	NE-CZ-NH2	8.50	124.55	120.30
1	D	27	ARG	NE-CZ-NH2	-7.39	116.61	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	314	0	289	2	0
1	B	322	0	294	1	0
1	C	314	0	289	0	0
1	D	309	0	286	1	0
1	E	330	0	299	2	0
1	F	322	0	298	0	0
1	G	314	0	289	2	0
1	H	322	0	294	0	0
1	I	328	0	301	1	0
1	J	309	0	277	1	0
1	K	301	0	268	0	0
1	L	318	0	293	3	0
2	M	75	0	53	1	0
2	O	77	0	53	0	0
2	R	77	0	53	0	0
2	X	75	0	53	0	0
3	N	100	0	74	0	0
3	U	100	0	74	0	0
4	P	61	0	41	1	0
5	Q	86	0	62	0	0
5	S	88	0	62	1	0
6	T	50	0	32	0	0
6	V	50	0	32	0	0
6	W	50	0	32	0	0
7	A	7	0	2	0	0
7	B	7	0	2	0	0
7	C	7	0	2	0	0
7	D	21	0	6	0	0
7	E	7	0	2	0	0
7	F	7	0	2	0	0
7	H	14	0	4	0	0
7	I	7	0	2	0	0
7	K	7	0	2	0	0
8	A	11	0	13	2	0
9	A	27	0	0	0	0
9	B	22	0	0	0	0
9	C	31	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
9	D	36	0	0	0	0
9	E	33	0	0	1	0
9	F	35	0	0	0	0
9	G	34	0	0	0	0
9	H	26	0	0	0	0
9	I	35	0	0	0	0
9	J	13	0	0	0	0
9	K	15	0	0	0	0
9	L	15	0	0	1	0
All	All	5109	0	4135	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 11 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:35[B]:ASN:ND2	9:E:201:HOH:O	2.29	0.65
1:B:6:THR:HB	1:L:6:THR:HB	1.82	0.60
1:A:25:ASP:HA	8:A:108:OOA:H7C1	1.85	0.59
1:E:6:THR:HB	1:I:6:THR:HB	1.89	0.54
1:J:39:THR:OG1	1:J:40:GLY:N	2.44	0.50

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	38/43 (88%)	38 (100%)	0	0	100	100
1	B	40/43 (93%)	40 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	38/43 (88%)	38 (100%)	0	0	100	100
1	D	37/43 (86%)	36 (97%)	1 (3%)	0	100	100
1	E	41/43 (95%)	41 (100%)	0	0	100	100
1	F	39/43 (91%)	39 (100%)	0	0	100	100
1	G	38/43 (88%)	37 (97%)	1 (3%)	0	100	100
1	H	39/43 (91%)	38 (97%)	1 (3%)	0	100	100
1	I	41/43 (95%)	40 (98%)	1 (2%)	0	100	100
1	J	38/43 (88%)	35 (92%)	3 (8%)	0	100	100
1	K	37/43 (86%)	37 (100%)	0	0	100	100
1	L	39/43 (91%)	38 (97%)	1 (3%)	0	100	100
All	All	465/516 (90%)	457 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	33/34 (97%)	33 (100%)	0	100	100
1	B	33/34 (97%)	33 (100%)	0	100	100
1	C	33/34 (97%)	33 (100%)	0	100	100
1	D	33/34 (97%)	33 (100%)	0	100	100
1	E	34/34 (100%)	34 (100%)	0	100	100
1	F	34/34 (100%)	34 (100%)	0	100	100
1	G	33/34 (97%)	33 (100%)	0	100	100
1	H	34/34 (100%)	34 (100%)	0	100	100
1	I	33/34 (97%)	33 (100%)	0	100	100
1	J	31/34 (91%)	31 (100%)	0	100	100
1	K	31/34 (91%)	31 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	33/34 (97%)	33 (100%)	0	100	100
All	All	395/408 (97%)	395 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

71 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	C4W	M	1	2	14,15,17	1.29	1 (7%)	18,21,23	3.24	4 (22%)
2	NAG	M	2	2	14,14,15	0.45	0	17,19,21	1.12	1 (5%)
2	BMA	M	3	2	11,11,12	0.63	0	15,15,17	1.55	4 (26%)
2	MAN	M	4	2	11,11,12	0.46	0	15,15,17	1.36	2 (13%)
2	NAG	M	5	2	14,14,15	0.42	0	17,19,21	1.00	1 (5%)
2	FUC	M	6	2	10,10,11	0.53	0	14,14,16	0.85	0
3	C4W	N	1	3	14,15,17	0.79	1 (7%)	18,21,23	1.64	3 (16%)
3	NAG	N	2	3	14,14,15	0.50	0	17,19,21	1.07	1 (5%)
3	BMA	N	3	3	11,11,12	0.61	0	15,15,17	1.13	1 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	MAN	N	4	3	11,11,12	0.35	0	15,15,17	1.22	2 (13%)
3	NAG	N	5	3	14,14,15	0.60	0	17,19,21	1.17	3 (17%)
3	MAN	N	6	3	11,11,12	0.56	0	15,15,17	1.49	2 (13%)
3	NAG	N	7	3	14,14,15	0.39	0	17,19,21	1.06	1 (5%)
3	FUC	N	8	3	10,10,11	0.57	0	14,14,16	0.77	0
2	C4W	O	1	2	16,17,17	1.04	1 (6%)	22,23,23	1.74	6 (27%)
2	NAG	O	2	2	14,14,15	0.57	0	17,19,21	1.10	2 (11%)
2	BMA	O	3	2	11,11,12	0.54	0	15,15,17	0.64	0
2	MAN	O	4	2	11,11,12	0.56	0	15,15,17	1.39	3 (20%)
2	NAG	O	5	2	14,14,15	0.28	0	17,19,21	0.82	1 (5%)
2	FUC	O	6	2	10,10,11	0.54	0	14,14,16	0.71	0
4	C4W	P	1	4	14,15,17	0.88	0	18,21,23	1.99	5 (27%)
4	NAG	P	2	4	14,14,15	0.41	0	17,19,21	0.94	0
4	BMA	P	3	4	11,11,12	0.48	0	15,15,17	1.78	6 (40%)
4	MAN	P	4	4	11,11,12	0.55	0	15,15,17	1.25	1 (6%)
4	FUC	P	5	4	10,10,11	0.49	0	14,14,16	0.78	0
5	C4W	Q	1	5	14,15,17	1.13	1 (7%)	18,21,23	1.98	4 (22%)
5	NAG	Q	2	5	14,14,15	0.52	0	17,19,21	1.02	1 (5%)
5	BMA	Q	3	5	11,11,12	0.57	0	15,15,17	0.79	0
5	MAN	Q	4	5	11,11,12	0.72	0	15,15,17	1.19	2 (13%)
5	NAG	Q	5	5	14,14,15	0.44	0	17,19,21	0.91	1 (5%)
5	MAN	Q	6	5	11,11,12	0.75	0	15,15,17	1.42	2 (13%)
5	FUC	Q	7	5	10,10,11	0.48	0	14,14,16	0.45	0
2	C4W	R	1	2	16,17,17	0.81	1 (6%)	22,23,23	1.76	4 (18%)
2	NAG	R	2	2	14,14,15	0.51	0	17,19,21	1.09	2 (11%)
2	BMA	R	3	2	11,11,12	0.35	0	15,15,17	0.79	0
2	MAN	R	4	2	11,11,12	0.54	0	15,15,17	2.03	5 (33%)
2	NAG	R	5	2	14,14,15	0.43	0	17,19,21	0.79	0
2	FUC	R	6	2	10,10,11	0.44	0	14,14,16	0.86	0
5	C4W	S	1	5	16,17,17	1.02	1 (6%)	22,23,23	1.93	4 (18%)
5	NAG	S	2	5	14,14,15	0.70	0	17,19,21	0.94	0
5	BMA	S	3	5	11,11,12	0.61	0	15,15,17	0.97	1 (6%)
5	MAN	S	4	5	11,11,12	0.56	0	15,15,17	1.23	3 (20%)
5	NAG	S	5	5	14,14,15	0.37	0	17,19,21	1.38	2 (11%)
5	MAN	S	6	5	11,11,12	0.61	0	15,15,17	1.69	3 (20%)
5	FUC	S	7	5	10,10,11	0.39	0	14,14,16	0.70	0
6	C4W	T	1	6	14,15,17	0.76	0	18,21,23	2.08	5 (27%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
6	NAG	T	2	6	14,14,15	0.60	0	17,19,21	1.09	1 (5%)
6	BMA	T	3	6	11,11,12	0.91	0	15,15,17	1.36	2 (13%)
6	FUC	T	4	6	10,10,11	0.47	0	14,14,16	0.81	0
3	C4W	U	1	3	14,15,17	0.87	0	18,21,23	1.78	4 (22%)
3	NAG	U	2	3	14,14,15	0.57	0	17,19,21	0.88	0
3	BMA	U	3	3	11,11,12	0.50	0	15,15,17	0.98	1 (6%)
3	MAN	U	4	3	11,11,12	0.52	0	15,15,17	1.16	0
3	NAG	U	5	3	14,14,15	0.51	0	17,19,21	1.02	0
3	MAN	U	6	3	11,11,12	0.44	0	15,15,17	1.03	1 (6%)
3	NAG	U	7	3	14,14,15	0.57	0	17,19,21	1.76	5 (29%)
3	FUC	U	8	3	10,10,11	0.51	0	14,14,16	0.87	0
6	C4W	V	1	6	14,15,17	0.74	0	18,21,23	1.90	4 (22%)
6	NAG	V	2	6	14,14,15	0.40	0	17,19,21	0.91	0
6	BMA	V	3	6	11,11,12	1.09	0	15,15,17	0.92	0
6	FUC	V	4	6	10,10,11	0.35	0	14,14,16	0.49	0
6	C4W	W	1	6	14,15,17	1.14	1 (7%)	18,21,23	2.24	5 (27%)
6	NAG	W	2	6	14,14,15	0.42	0	17,19,21	1.16	2 (11%)
6	BMA	W	3	6	11,11,12	0.78	0	15,15,17	0.97	1 (6%)
6	FUC	W	4	6	10,10,11	0.40	0	14,14,16	1.05	1 (7%)
2	C4W	X	1	2	14,15,17	1.28	1 (7%)	18,21,23	2.19	5 (27%)
2	NAG	X	2	2	14,14,15	0.41	0	17,19,21	1.13	1 (5%)
2	BMA	X	3	2	11,11,12	0.83	0	15,15,17	1.06	1 (6%)
2	MAN	X	4	2	11,11,12	0.57	0	15,15,17	1.39	2 (13%)
2	NAG	X	5	2	14,14,15	0.53	0	17,19,21	1.10	2 (11%)
2	FUC	X	6	2	10,10,11	0.47	0	14,14,16	0.63	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	C4W	M	1	2	-	0/6/26/29	0/1/1/1
2	NAG	M	2	2	-	0/6/23/26	0/1/1/1
2	BMA	M	3	2	-	0/2/19/22	0/1/1/1
2	MAN	M	4	2	-	0/2/19/22	0/1/1/1
2	NAG	M	5	2	-	2/6/23/26	0/1/1/1
2	FUC	M	6	2	-	-	0/1/1/1
3	C4W	N	1	3	-	0/6/26/29	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	N	2	3	-	2/6/23/26	0/1/1/1
3	BMA	N	3	3	-	0/2/19/22	0/1/1/1
3	MAN	N	4	3	-	1/2/19/22	0/1/1/1
3	NAG	N	5	3	-	0/6/23/26	0/1/1/1
3	MAN	N	6	3	-	1/2/19/22	0/1/1/1
3	NAG	N	7	3	-	0/6/23/26	0/1/1/1
3	FUC	N	8	3	-	-	0/1/1/1
2	C4W	O	1	2	-	0/8/29/29	0/1/1/1
2	NAG	O	2	2	-	0/6/23/26	0/1/1/1
2	BMA	O	3	2	-	0/2/19/22	0/1/1/1
2	MAN	O	4	2	-	2/2/19/22	0/1/1/1
2	NAG	O	5	2	-	0/6/23/26	0/1/1/1
2	FUC	O	6	2	-	-	0/1/1/1
4	C4W	P	1	4	-	0/6/26/29	0/1/1/1
4	NAG	P	2	4	-	0/6/23/26	0/1/1/1
4	BMA	P	3	4	-	2/2/19/22	0/1/1/1
4	MAN	P	4	4	-	1/2/19/22	0/1/1/1
4	FUC	P	5	4	-	-	0/1/1/1
5	C4W	Q	1	5	-	0/6/26/29	0/1/1/1
5	NAG	Q	2	5	-	0/6/23/26	0/1/1/1
5	BMA	Q	3	5	-	0/2/19/22	0/1/1/1
5	MAN	Q	4	5	-	0/2/19/22	0/1/1/1
5	NAG	Q	5	5	-	0/6/23/26	0/1/1/1
5	MAN	Q	6	5	-	0/2/19/22	0/1/1/1
5	FUC	Q	7	5	-	-	0/1/1/1
2	C4W	R	1	2	-	0/8/29/29	0/1/1/1
2	NAG	R	2	2	-	0/6/23/26	0/1/1/1
2	BMA	R	3	2	-	0/2/19/22	0/1/1/1
2	MAN	R	4	2	-	2/2/19/22	0/1/1/1
2	NAG	R	5	2	-	0/6/23/26	0/1/1/1
2	FUC	R	6	2	-	-	0/1/1/1
5	C4W	S	1	5	-	0/8/29/29	0/1/1/1
5	NAG	S	2	5	-	0/6/23/26	0/1/1/1
5	BMA	S	3	5	-	0/2/19/22	0/1/1/1
5	MAN	S	4	5	-	0/2/19/22	0/1/1/1
5	NAG	S	5	5	-	1/6/23/26	0/1/1/1
5	MAN	S	6	5	-	0/2/19/22	0/1/1/1
5	FUC	S	7	5	-	-	0/1/1/1
6	C4W	T	1	6	-	0/6/26/29	0/1/1/1
6	NAG	T	2	6	-	0/6/23/26	0/1/1/1
6	BMA	T	3	6	-	2/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	FUC	T	4	6	-	-	0/1/1/1
3	C4W	U	1	3	-	0/6/26/29	0/1/1/1
3	NAG	U	2	3	-	0/6/23/26	0/1/1/1
3	BMA	U	3	3	-	0/2/19/22	0/1/1/1
3	MAN	U	4	3	-	0/2/19/22	0/1/1/1
3	NAG	U	5	3	-	2/6/23/26	0/1/1/1
3	MAN	U	6	3	-	0/2/19/22	0/1/1/1
3	NAG	U	7	3	-	2/6/23/26	0/1/1/1
3	FUC	U	8	3	-	-	0/1/1/1
6	C4W	V	1	6	-	0/6/26/29	0/1/1/1
6	NAG	V	2	6	-	0/6/23/26	0/1/1/1
6	BMA	V	3	6	-	1/2/19/22	0/1/1/1
6	FUC	V	4	6	-	-	0/1/1/1
6	C4W	W	1	6	-	0/6/26/29	0/1/1/1
6	NAG	W	2	6	-	2/6/23/26	0/1/1/1
6	BMA	W	3	6	-	0/2/19/22	0/1/1/1
6	FUC	W	4	6	-	-	0/1/1/1
2	C4W	X	1	2	-	0/6/26/29	0/1/1/1
2	NAG	X	2	2	-	0/6/23/26	0/1/1/1
2	BMA	X	3	2	-	0/2/19/22	0/1/1/1
2	MAN	X	4	2	-	0/2/19/22	0/1/1/1
2	NAG	X	5	2	-	0/6/23/26	0/1/1/1
2	FUC	X	6	2	-	-	0/1/1/1

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	M	1	C4W	C1-C2	3.47	1.56	1.53
2	X	1	C4W	C1-C2	3.27	1.56	1.53
5	Q	1	C4W	C1-C2	2.91	1.55	1.53
5	S	1	C4W	O5-C1	2.72	1.46	1.41
6	W	1	C4W	C1-C2	2.66	1.55	1.53

The worst 5 of 126 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	M	1	C4W	C1-C2-N2	-8.69	100.52	111.12
2	M	1	C4W	C2-N2-C7	7.23	140.75	123.18
2	M	1	C4W	C3-C2-N2	6.48	122.85	110.62
2	X	1	C4W	C8-C7-N2	5.72	125.79	116.10
6	W	1	C4W	C3-C2-N2	5.10	120.24	110.62

There are no chirality outliers.

5 of 23 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	U	7	NAG	O5-C5-C6-O6
6	W	2	NAG	O5-C5-C6-O6
6	W	2	NAG	C4-C5-C6-O6
3	U	5	NAG	O5-C5-C6-O6
3	U	7	NAG	C4-C5-C6-O6

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	M	6	FUC	1	0
5	S	7	FUC	1	0
4	P	5	FUC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

Oligosaccharide Chain M



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain O



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain R



Oligosaccharide Chain X



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain N



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain U



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain P



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain Q



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain S



Oligosaccharide Chain T



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain V



Oligosaccharide Chain W



5.6 Ligand geometry (i)

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
7	MLI	I	109	-	6,6,6	1.10	0	7,7,7	0.85	0
7	MLI	E	108	-	6,6,6	1.46	0	7,7,7	1.59	1 (14%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
8	OOA	A	108	-	10,10,10	1.49	2 (20%)	10,11,11	1.30	1 (10%)
7	MLI	D	108	-	6,6,6	1.55	1 (16%)	7,7,7	1.53	2 (28%)
7	MLI	F	107	-	6,6,6	1.60	0	7,7,7	0.66	0
7	MLI	A	107	-	6,6,6	1.41	1 (16%)	7,7,7	1.06	0
7	MLI	C	107	-	6,6,6	1.39	0	7,7,7	0.41	0
7	MLI	B	109	-	6,6,6	1.31	0	7,7,7	1.27	1 (14%)
7	MLI	H	106	-	6,6,6	1.16	0	7,7,7	1.42	1 (14%)
7	MLI	K	105	-	6,6,6	1.19	0	7,7,7	0.63	0
7	MLI	D	107	-	6,6,6	1.48	1 (16%)	7,7,7	1.05	0
7	MLI	D	106	-	6,6,6	1.20	0	7,7,7	1.55	3 (42%)
7	MLI	H	105	-	6,6,6	1.22	0	7,7,7	1.14	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	MLI	I	109	-	-	4/4/4/4	-
7	MLI	E	108	-	-	0/4/4/4	-
8	OOA	A	108	-	-	3/9/9/9	-
7	MLI	D	108	-	-	4/4/4/4	-
7	MLI	F	107	-	-	0/4/4/4	-
7	MLI	A	107	-	-	2/4/4/4	-
7	MLI	C	107	-	-	0/4/4/4	-
7	MLI	B	109	-	-	0/4/4/4	-
7	MLI	H	106	-	-	2/4/4/4	-
7	MLI	K	105	-	-	0/4/4/4	-
7	MLI	D	107	-	-	0/4/4/4	-
7	MLI	D	106	-	-	2/4/4/4	-
7	MLI	H	105	-	-	3/4/4/4	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	A	108	OOA	O1-C3	3.12	1.27	1.21
8	A	108	OOA	C2-C1	2.13	1.54	1.51
7	D	107	MLI	C1-C2	2.12	1.54	1.51
7	D	108	MLI	C1-C3	2.05	1.54	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	107	MLI	O7-C2	-2.02	1.24	1.30

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
8	A	108	OOA	C3-C2-C1	2.94	118.47	112.38
7	E	108	MLI	O9-C3-C1	2.49	122.50	114.54
7	D	108	MLI	O7-C2-C1	2.17	121.47	114.54
7	D	106	MLI	O6-C2-C1	-2.09	115.96	122.08
7	H	106	MLI	O7-C2-C1	2.08	121.18	114.54

There are no chirality outliers.

5 of 20 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	A	108	OOA	C3-C4-C5-C6
7	I	109	MLI	C3-C1-C2-O7
7	D	106	MLI	C2-C1-C3-O8
7	D	108	MLI	C2-C1-C3-O8
7	I	109	MLI	C3-C1-C2-O6

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	A	108	OOA	2	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	40/43 (93%)	0.56	1 (2%)	57	62	30, 38, 50, 58	1 (2%)
1	B	42/43 (97%)	0.33	2 (4%)	30	36	35, 41, 50, 59	0
1	C	40/43 (93%)	0.35	0	100	100	33, 38, 50, 53	0
1	D	39/43 (90%)	0.55	0	100	100	29, 37, 47, 48	0
1	E	42/43 (97%)	0.51	2 (4%)	30	36	29, 33, 47, 57	0
1	F	41/43 (95%)	0.33	1 (2%)	59	64	29, 34, 45, 59	0
1	G	40/43 (93%)	0.43	1 (2%)	57	62	28, 35, 48, 54	0
1	H	40/43 (93%)	0.64	1 (2%)	57	62	29, 38, 49, 62	0
1	I	43/43 (100%)	0.46	1 (2%)	60	65	29, 34, 45, 56	0
1	J	40/43 (93%)	0.40	2 (5%)	28	34	36, 45, 61, 70	0
1	K	39/43 (90%)	0.59	3 (7%)	13	17	40, 49, 61, 68	0
1	L	41/43 (95%)	0.42	1 (2%)	59	64	40, 47, 59, 68	0
All	All	487/516 (94%)	0.46	15 (3%)	49	55	28, 39, 57, 70	1 (0%)

The worst 5 of 15 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	40	GLY	5.6
1	I	40	GLY	4.2
1	A	40	GLY	3.9
1	G	40	GLY	3.9
1	B	40	GLY	3.4

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
6	BMA	W	3	11/12	0.73	0.23	64,68,73,76	0
2	BMA	X	3	11/12	0.75	0.29	52,56,58,60	0
6	BMA	V	3	11/12	0.78	0.31	51,57,68,68	0
5	NAG	S	5	14/15	0.80	0.28	54,59,63,64	14
6	BMA	T	3	11/12	0.80	0.15	57,59,65,65	0
4	BMA	P	3	11/12	0.80	0.21	57,61,63,64	0
5	MAN	S	4	11/12	0.80	0.25	60,65,68,69	11
5	MAN	Q	6	11/12	0.82	0.23	56,59,61,64	0
2	BMA	M	3	11/12	0.82	0.22	57,59,61,65	0
4	MAN	P	4	11/12	0.82	0.21	59,59,62,63	11
3	NAG	U	5	14/15	0.85	0.34	53,57,60,61	0
3	NAG	N	5	14/15	0.87	0.18	60,62,65,65	0
2	NAG	X	2	14/15	0.88	0.26	49,58,63,63	0
2	MAN	R	4	11/12	0.88	0.22	55,57,61,62	0
2	NAG	M	5	14/15	0.89	0.25	52,55,65,65	0
3	MAN	U	4	11/12	0.90	0.23	51,55,59,59	0
5	MAN	Q	4	11/12	0.90	0.13	45,52,55,55	0
2	MAN	X	4	11/12	0.90	0.16	50,54,57,59	0
2	BMA	R	3	11/12	0.90	0.20	54,56,61,67	0
2	MAN	M	4	11/12	0.91	0.22	54,59,62,63	0
5	MAN	S	6	11/12	0.92	0.21	49,54,56,57	11
6	C4W	W	1	15/17	0.92	0.25	49,54,67,68	0
2	C4W	X	1	15/17	0.92	0.17	53,59,66,67	0
2	C4W	M	1	15/17	0.93	0.17	44,47,65,65	0
2	NAG	X	5	14/15	0.93	0.31	45,50,57,57	0
2	FUC	X	6	10/11	0.93	0.17	50,53,53,54	0
4	C4W	P	1	15/17	0.93	0.16	38,46,61,62	0
6	NAG	W	2	14/15	0.93	0.17	53,57,66,67	0
3	MAN	N	4	11/12	0.93	0.20	56,57,59,60	0
2	MAN	O	4	11/12	0.94	0.18	51,55,62,62	0
3	NAG	U	7	14/15	0.94	0.17	39,46,51,57	0
2	NAG	M	2	14/15	0.94	0.14	44,48,56,62	0
2	NAG	O	2	14/15	0.94	0.12	37,40,47,47	0
3	MAN	N	6	11/12	0.94	0.11	48,52,56,58	0
3	NAG	N	7	14/15	0.94	0.16	40,48,62,64	0
2	BMA	O	3	11/12	0.94	0.17	47,51,56,60	0
5	BMA	S	3	11/12	0.94	0.15	43,50,55,63	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	C4W	R	1	17/17	0.95	0.13	36,44,53,59	0
2	C4W	O	1	17/17	0.95	0.12	33,37,42,48	0
4	FUC	P	5	10/11	0.95	0.12	38,40,41,41	0
6	C4W	T	1	15/17	0.95	0.15	38,45,58,59	0
5	NAG	Q	2	14/15	0.95	0.13	33,35,37,38	0
6	NAG	V	2	14/15	0.95	0.14	43,44,46,52	0
2	NAG	O	5	14/15	0.95	0.13	47,49,60,67	0
5	NAG	Q	5	14/15	0.95	0.17	46,51,63,66	0
3	C4W	N	1	15/17	0.95	0.14	37,44,53,56	0
4	NAG	P	2	14/15	0.95	0.17	39,43,54,56	0
5	FUC	Q	7	10/11	0.96	0.14	31,34,36,36	0
5	C4W	S	1	17/17	0.96	0.14	32,38,48,55	0
3	FUC	U	8	10/11	0.96	0.11	33,35,40,42	0
2	NAG	R	2	14/15	0.96	0.14	40,42,50,50	0
2	NAG	R	5	14/15	0.96	0.16	51,55,62,62	0
3	FUC	N	8	10/11	0.96	0.12	39,43,46,47	0
3	C4W	U	1	15/17	0.96	0.13	33,40,52,53	0
3	BMA	U	3	11/12	0.96	0.12	37,41,44,45	0
6	C4W	V	1	15/17	0.96	0.12	39,45,51,51	0
5	C4W	Q	1	15/17	0.96	0.11	33,36,47,49	0
3	BMA	N	3	11/12	0.96	0.13	44,47,51,55	0
2	FUC	R	6	10/11	0.96	0.11	37,38,40,43	0
3	MAN	U	6	11/12	0.96	0.12	43,49,52,55	0
2	FUC	M	6	10/11	0.96	0.10	39,41,43,44	0
6	FUC	W	4	10/11	0.96	0.11	47,49,50,55	0
5	NAG	S	2	14/15	0.97	0.12	34,37,40,41	0
5	BMA	Q	3	11/12	0.97	0.14	39,45,50,53	0
6	FUC	V	4	10/11	0.97	0.10	40,42,45,46	0
6	NAG	T	2	14/15	0.97	0.14	40,43,52,55	0
3	NAG	U	2	14/15	0.97	0.11	32,35,37,39	0
6	FUC	T	4	10/11	0.97	0.13	36,38,39,40	0
3	NAG	N	2	14/15	0.97	0.13	39,41,47,55	0
5	FUC	S	7	10/11	0.98	0.14	32,35,39,39	0
2	FUC	O	6	10/11	0.99	0.14	35,36,39,40	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain M:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain O:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain R:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain X:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain N:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain U:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain P:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain Q:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain S:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain T:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain V:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain W:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
7	MLI	F	107	7/7	0.67	0.32	59,61,64,66	0
8	OOA	A	108	11/11	0.73	0.23	38,52,61,63	0
7	MLI	D	108	7/7	0.85	0.15	59,60,65,66	0
7	MLI	K	105	7/7	0.86	0.41	59,60,63,63	0
7	MLI	D	106	7/7	0.91	0.27	50,54,57,60	0
7	MLI	D	107	7/7	0.92	0.22	49,55,55,56	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
7	MLI	H	105	7/7	0.93	0.28	47,55,57,59	0
7	MLI	I	109	7/7	0.93	0.38	48,49,55,55	0
7	MLI	B	109	7/7	0.94	0.19	49,52,53,55	0
7	MLI	C	107	7/7	0.94	0.20	45,50,52,54	0
7	MLI	A	107	7/7	0.95	0.13	47,53,54,54	0
7	MLI	H	106	7/7	0.97	0.24	46,54,56,60	0
7	MLI	E	108	7/7	0.97	0.23	44,51,55,60	0

6.5 Other polymers (i)

There are no such residues in this entry.