Full wwPDB X-ray Structure Validation Report (i)

May 18, 2020 – 06:35 am BST

PDB ID	:	6FYK
Title	:	X-Ray structure of CLK2-KD(136-496)/Indazole1 at 2.39A
Authors	:	Kallen, J.
Deposited on	:	2018-03-12
Resolution	:	2.39 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	363	5% 90% 6% •
1	B	363	4% 91% 5% •
1	C	363	5% 90% 6% •

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 9306 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dual specificity protein kinase CLK2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	349	2920	1857	520	529	14	0	4	0
1	B	349	2928	1861	520	533	14	0	6	0
1	C	349	2920	1857	520	529	14	0	4	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	134	GLY	-	expression tag	UNP P49760
A	135	PRO	-	expression tag	UNP P49760
B	134	GLY	-	expression tag	UNP P49760
B	135	PRO	-	expression tag	UNP P49760
C	134	GLY	-	expression tag	UNP P49760
C	135	PRO	-	expression tag	UNP P49760

- Molecule 2 is (2 {S})-4-methyl-1-[5-(3-methyl-2 {H}-indazol-5-yl)pyridin-3-yl]oxy-pentan-2-amine (three-letter code: EAZ) (formula: $C_{19}H_{24}N_4O$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			24	19	4	1		
2	B	1	Total	C	N	O	0	0
			24	19	4	1		
2	C	1	Total	C	N	O	0	0
			24	19	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	189	Total	O	0	0
			189	189		
3	B	149	Total	O	0	0
			149	149		
3	C	128	Total	O	0	0
			128	128		

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Dual specificity protein kinase CLK2

Chain A:



- Molecule 1: Dual specificity protein kinase CLK2

Chain B:





- Molecule 1: Dual specificity protein kinase CLK2

Chain C:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α, β, γ	95.63Å 95.63Å 222.86Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.68 – 2.39 19.68 – 2.39	Depositor EDS
% Data completeness (in resolution range)	99.8 (19.68-2.39) 100.0 (19.68-2.39)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.92 (at 2.38Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.195 , 0.228 0.194 , 0.227	Depositor DCC
R_{free} test set	2376 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å^2)	34.3	Xtriage
Anisotropy	0.022	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.33 , 31.5	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.028 for -h,-k,l	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	9306	wwPDB-VP
Average B, all atoms (Å^2)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.76% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: EAZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.37	0/3004	0.51	0/4056
1	B	0.37	0/3018	0.51	0/4074
1	C	0.37	0/3004	0.50	0/4056
All	All	0.37	0/9026	0.51	0/12186

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2920	0	2861	13	0
1	B	2928	0	2868	8	0
1	C	2920	0	2861	12	0
2	A	24	0	0	1	0
2	B	24	0	0	0	0
2	C	24	0	0	0	0
3	A	189	0	0	1	0
3	B	149	0	0	0	0
3	C	128	0	0	2	0
All	All	9306	0	8590	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:251:PHE:HB2	1:B:293:PRO:HB2	1.52	0.90
1:A:251:PHE:HB2	1:A:293:PRO:HB2	1.59	0.85
1:C:239:MET:HG3	3:C:625:HOH:O	1.82	0.80
1:B:439:ARG:NH2	1:B:454:ASP:OD1	2.23	0.71
1:C:195[B]:ILE:HD13	1:C:205:ALA:HB2	1.78	0.63
1:C:195[B]:ILE:HD13	1:C:205:ALA:CB	2.31	0.61
1:A:203:GLU:OE2	1:A:206:ARG:NH2	2.35	0.58
1:C:439:ARG:NH2	1:C:454:ASP:OD1	2.36	0.58
1:A:439:ARG:NH2	1:A:454:ASP:OD1	2.37	0.57
1:B:138:ARG:HH11	1:B:146:ASP:HB3	1.70	0.55
1:A:195[B]:ILE:HD13	1:A:205:ALA:HB2	1.87	0.55
1:C:251:PHE:HB2	1:C:293:PRO:HB2	1.91	0.53
1:B:164:GLU:OE1	1:B:183:HIS:HE1	1.92	0.51
1:C:297:LEU:HD22	1:C:326:VAL:HG11	1.92	0.51
1:B:297:LEU:HD22	1:B:326:VAL:HG11	1.93	0.50
1:C:165:ILE:HG23	1:C:178[B]:VAL:CG1	2.42	0.49
1:A:480:ALA:C	1:A:482:LEU:H	2.15	0.49
1:C:165:ILE:HG23	1:C:178[B]:VAL:HG11	1.95	0.49
1:C:453:PHE:O	1:C:457:GLU:HB2	2.12	0.48
1:B:356:LEU:HD22	1:B:407:LYS:HG2	1.95	0.48
1:C:394:GLU:HA	1:C:398:GLY:O	2.13	0.48
1:B:260:LEU:HD13	1:B:315:GLU:HG2	1.96	0.46
1:A:208:GLU:HG3	1:A:328:PHE:HB2	1.98	0.46
1:A:195[B]:ILE:HD13	1:A:205:ALA:CB	2.47	0.45
1:B:405:ILE:O	1:B:408:THR:HG22	2.16	0.45
1:A:297:LEU:HD22	1:A:326:VAL:HG11	2.00	0.44
1:A:364:VAL:HG11	1:A:468:LEU:O	2.16	0.44
1:C:293:PRO:HD3	1:C:370:ILE:HG12	2.00	0.42
1:A:327:ASP:HB2	2:A:501:EAZ:C4	2.49	0.42
1:A:146:ASP:HB2	3:A:694:HOH:O	2.20	0.41
1:A:227:VAL:HB	1:A:326:VAL:HG12	2.03	0.41
1:A:459:MET:O	1:A:467:ARG:HD3	2.21	0.40
1:C:185:ARG:HD3	3:C:708:HOH:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	351/363 (97%)	341 (97%)	8 (2%)	2 (1%)	25	36
1	B	353/363 (97%)	344 (98%)	9 (2%)	0	100	100
1	C	351/363 (97%)	335 (95%)	14 (4%)	2 (1%)	25	36
All	All	1055/1089 (97%)	1020 (97%)	31 (3%)	4 (0%)	34	48

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	481	ARG
1	C	483	ARG
1	C	484	ALA
1	A	327	ASP

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	318/327 (97%)	317 (100%)	1 (0%)	92	97
1	B	320/327 (98%)	317 (99%)	3 (1%)	78	90
1	C	318/327 (97%)	314 (99%)	4 (1%)	69	84
All	All	956/981 (98%)	948 (99%)	8 (1%)	81	91

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	483	ARG
1	B	146	ASP
1	B	184	ARG
1	B	305	LEU
1	C	146	ASP
1	C	309	LEU
1	C	409	ARG
1	C	434	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	183	HIS
1	A	228	GLN
1	A	451	GLN
1	B	154	HIS
1	B	179	GLN
1	B	183	HIS
1	B	224	ASN
1	B	228	GLN
1	B	451	GLN
1	C	160	GLN
1	C	179	GLN
1	C	228	GLN
1	C	451	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	EAZ	A	501	-	24,26,26	1.36	2 (8%)	27,36,36	1.64	6 (22%)
2	EAZ	B	501	-	24,26,26	1.43	2 (8%)	27,36,36	1.80	7 (25%)
2	EAZ	C	501	-	24,26,26	1.41	1 (4%)	27,36,36	1.60	6 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EAZ	A	501	-	-	2/13/13/13	0/3/3/3
2	EAZ	B	501	-	-	4/13/13/13	0/3/3/3
2	EAZ	C	501	-	-	4/13/13/13	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	EAZ	C14-N15	-4.68	1.27	1.34
2	C	501	EAZ	C14-N15	-4.54	1.28	1.34
2	A	501	EAZ	C14-N15	-4.49	1.28	1.34
2	B	501	EAZ	C20-C19	2.11	1.56	1.53
2	A	501	EAZ	C10-C11	2.02	1.40	1.36

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	B	501	EAZ	C18-O7-C5	3.86	126.12	117.93
2	B	501	EAZ	C4-N3-C2	3.74	122.58	117.48
2	A	501	EAZ	C18-O7-C5	3.63	125.62	117.93

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	C	501	EAZ	C4-N3-C2	3.54	122.31	117.48
2	B	501	EAZ	C11-C10-C12	-3.42	116.53	120.84
2	B	501	EAZ	C9-C13-C12	3.38	122.89	118.26
2	A	501	EAZ	C11-C10-C12	-3.31	116.67	120.84
2	C	501	EAZ	C9-C13-C12	3.30	122.77	118.26
2	A	501	EAZ	C9-C13-C12	3.28	122.74	118.26
2	C	501	EAZ	C11-C10-C12	-3.04	117.01	120.84
2	A	501	EAZ	C4-N3-C2	3.02	121.59	117.48
2	C	501	EAZ	C18-O7-C5	3.00	124.29	117.93
2	C	501	EAZ	C17-C14-N15	2.77	125.74	119.65
2	B	501	EAZ	C17-C14-N15	2.72	125.64	119.65
2	B	501	EAZ	C8-C9-C13	-2.64	116.53	122.30
2	A	501	EAZ	C8-C9-C13	-2.63	116.54	122.30
2	C	501	EAZ	C8-C9-C13	-2.56	116.70	122.30
2	A	501	EAZ	C17-C14-N15	2.51	125.16	119.65
2	B	501	EAZ	O7-C18-C19	-2.09	103.68	108.29

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	501	EAZ	C19-C20-C21-C22
2	C	501	EAZ	C19-C20-C21-C22
2	C	501	EAZ	C19-C20-C21-C23
2	B	501	EAZ	C19-C20-C21-C23
2	A	501	EAZ	C19-C20-C21-C22
2	A	501	EAZ	C19-C20-C21-C23
2	C	501	EAZ	C6-C5-O7-C18
2	B	501	EAZ	C4-C5-O7-C18
2	B	501	EAZ	C6-C5-O7-C18
2	C	501	EAZ	C4-C5-O7-C18

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	EAZ	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand EAZ A 501



Ligand EAZ B 501



Ligand EAZ C 501



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	349/363 (96%)	0.04	18 (5%)	27	26	18, 27, 60, 119	0
1	B	349/363 (96%)	0.21	16 (4%)	32	31	21, 34, 58, 154	0
1	C	349/363 (96%)	0.23	17 (4%)	29	28	22, 37, 64, 115	0
All	All	1047/1089 (96%)	0.16	51 (4%)	29	28	18, 33, 61, 154	0

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	484	ALA	13.9
1	B	485	GLU	13.5
1	B	483	ARG	12.4
1	C	484	ALA	10.2
1	C	485	GLU	9.5
1	A	309	LEU	7.8
1	A	485	GLU	6.9
1	A	483	ARG	6.6
1	A	307	TYR	6.3
1	A	484	ALA	6.2
1	A	312	LYS	6.0
1	B	482	LEU	5.3
1	B	481	ARG	5.0
1	A	306	THR	4.8
1	A	313	ARG	4.8
1	B	415	TYR	4.4
1	A	482	LEU	4.3
1	C	221	ASP	4.1
1	C	313	ARG	4.0
1	B	309	LEU	3.9
1	C	223	LYS	3.7
1	A	305	LEU	3.6
1	C	220	PRO	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	310	GLU	3.3
1	A	192	LEU	3.3
1	B	310	GLU	3.2
1	A	308	ASN	3.1
1	A	314	ASP	3.0
1	C	483	ARG	2.9
1	B	313	ARG	2.8
1	A	481	ARG	2.8
1	B	409	ARG	2.7
1	C	310	GLU	2.6
1	C	241	ILE	2.6
1	C	200	LYS	2.5
1	C	192	LEU	2.5
1	A	311	LYS	2.5
1	B	184	ARG	2.4
1	A	137	SER	2.4
1	B	241	ILE	2.4
1	C	429	ARG	2.3
1	A	199	GLU	2.3
1	B	185	ARG	2.2
1	C	222	ASN	2.2
1	C	409	ARG	2.2
1	B	221	ASP	2.2
1	B	465	ALA	2.2
1	C	144	GLU	2.1
1	B	416	ARG	2.1
1	C	481	ARG	2.1
1	C	137	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	EAZ	B	501	24/24	0.96	0.15	21,26,26,27	0
2	EAZ	A	501	24/24	0.97	0.12	19,22,25,25	0
2	EAZ	C	501	24/24	0.97	0.12	24,25,26,27	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around EAZ B 501:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around EAZ A 501:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around EAZ C 501:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.