Full wwPDB X-ray Structure Validation Report (i)

Aug 20, 2023 – 03:56 AM EDT

PDB ID	:	2G9H
Title	:	Crystal Structure of Staphylococcal Enterotoxin I (SEI) in Complex with a Human MHC class II Molecule
Authors	:	Fernandez, M.M.; Guan, R.; Malchiodi, E.L.; Mariuzza, R.A.
Deposited on	:	2006-03-06
Resolution	:	2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	182	2%, 93%, • • •
2	B	190	4%, 86%, 12%, •
3	C	13	8%, 85%, 8%, 8%
4	D	218	4%, 83%, 12%, • •

2 Entry composition (i)

There are 9 unique types of molecules in this entry. The entry contains 5364 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class II histocompatibility antigen, DR alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	178	1464	949	238	272	5	0	0	0

- Molecule 2 is a protein called HLA class II histocompatibility antigen, DRB1-1 beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	190	1557	979	279	293	6	0	0	0

- Molecule 3 is a protein called Hemagglutinin.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	13	106	69	18	19	0	0	0

- Molecule 4 is a protein called extracellular enterotoxin type I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	213	1729	1112	284	331	2	0	0	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	24	LEU	VAL	SEE REMARK 999	UNP O85383
D	25	ILE	THR	SEE REMARK 999	UNP O85383
D	31	SER	ILE	SEE REMARK 999	UNP O85383
D	41	ILE	THR	SEE REMARK 999	UNP O85383
D	114	SER	ALA	SEE REMARK 999	UNP O85383
D	144	SER	ASN	SEE REMARK 999	UNP O85383
D	163	LYS	ASN	SEE REMARK 999	UNP O85383
D	172	ASP	ASN	SEE REMARK 999	UNP O85383

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	187	VAL	LEU	SEE REMARK 999	UNP O85383

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	O	S		0	0
			5	4	1			
5	A	1	Total	O	S		0	0
			5	4	1			
5	C	1	Total	O	S		0	0
			5	4	1			
5	D	1	Total	O	S		0	0
			5	4	1			

- Molecule 6 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (three-letter code: EPE) (formula: $C_8H_{18}N_2O_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
6	A	1	15	8	2	4	1	0	0

- Molecule 7 is 1,4-DIETHYLENE DIOXIDE (three-letter code: DIO) (formula: $C_4H_8O_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	O				
7	A	1	6	4	2			0	0
			Total	C	O				
7	A	1	6	4	2			0	0

- Molecule 8 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
8	D	1	1	1	0	0

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
9	A	160	160	160	0	0
			Total	O		
9	B	132	132	132	0	0
			Total	O		
9	C	11	11	11	0	0
			Total	O		
9	D	157	157	157	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: HLA class II histocompatibility antigen, DR alpha chain

Chain A:





• Molecule 2: HLA class II histocompatibility antigen, DRB1-1 beta chain

Chain B:



• Molecule 3: Hemagglutinin

Chain C:



• Molecule 4: extracellular enterotoxin type I

Chain D:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α, β, γ	150.40Å 99.93Å 72.92Å 90.00° 92.00° 90.00°	Depositor
Resolution (Å)	29.79 – 2.00 29.78 – 2.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (29.79-2.00) 99.2 (29.78-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.95 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.213 , 0.252 0.214 , 0.250	Depositor DCC
R_{free} test set	2949 reflections (4.09%)	wwPDB-VP
Wilson B-factor (Å^2)	30.2	Xtriage
Anisotropy	0.327	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.33 , 33.9	EDS
L-test for twinning[2]	$< \|L\| > = 0.48$, $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	0.000 for -h,-k,l	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	5364	wwPDB-VP
Average B, all atoms (Å^2)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.16% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, EPE, DIO, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.55	0/1509	0.70	1/2058 (0.0%)
2	B	0.54	0/1597	0.66	0/2168
3	C	0.46	0/107	0.77	0/141
4	D	0.61	0/1768	0.71	1/2377 (0.0%)
All	All	0.57	0/4981	0.69	2/6744 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	146	ARG	NE-CZ-NH2	-5.89	117.35	120.30
4	D	44	LEU	CA-CB-CG	5.47	127.88	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1464	0	1401	6	0
2	B	1557	0	1488	18	0
3	C	106	0	119	1	0
4	D	1729	0	1702	16	0
5	A	10	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	C	5	0	0	0	0
5	D	5	0	0	0	0
6	A	15	0	18	3	0
7	A	12	0	16	3	0
8	D	1	0	0	0	0
9	A	160	0	0	4	0
9	B	132	0	0	2	0
9	C	11	0	0	0	0
9	D	157	0	0	1	0
All	All	5364	0	4744	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:106:THR:HG23	2:B:108:PRO:HD3	1.27	1.11
4:D:139:ILE:HD11	4:D:212:ILE:HD12	1.57	0.84
2:B:46:GLU:OE2	2:B:48:ARG:NH1	2.21	0.73
2:B:112:HIS:O	9:B:322:HOH:O	2.09	0.70
2:B:145:THR:HG22	2:B:146:GLY:O	1.96	0.65
6:A:601:EPE:H21	9:A:810:HOH:O	1.99	0.63
2:B:100:THR:HG22	9:B:252:HOH:O	1.99	0.62
4:D:24:LEU:HD12	4:D:64:ILE:HD13	1.84	0.59
1:A:134:GLU:OE1	1:A:147:LYS:NZ	2.37	0.58
4:D:139:ILE:CD1	4:D:212:ILE:HD12	2.34	0.56
1:A:69:ASN:HB3	3:C:316:LEU:HD22	1.88	0.56
2:B:104:SER:HB3	2:B:106:THR:O	2.07	0.54
4:D:19:ILE:HD11	4:D:68:ASP:HB2	1.88	0.54
2:B:106:THR:O	2:B:106:THR:CG2	2.56	0.53
9:A:846:HOH:O	2:B:100:THR:HG21	2.09	0.53
2:B:106:THR:HG23	2:B:108:PRO:CD	2.20	0.53
1:A:177:HIS:HE1	9:A:854:HOH:O	1.91	0.53
7:A:701:DIO:H22	2:B:3:THR:HA	1.92	0.51
2:B:145:THR:CG2	2:B:146:GLY:O	2.58	0.51
4:D:63:ASP:C	4:D:64:ILE:HD12	2.31	0.51
2:B:149:GLN:HG3	2:B:155:PHE:CE2	2.46	0.51
7:A:701:DIO:H1'1	9:A:826:HOH:O	2.10	0.50
7:A:701:DIO:H1'2	2:B:4:ARG:O	2.12	0.49
4:D:139:ILE:HD11	4:D:212:ILE:CD1	2.37	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:40:PHE:HB2	2:B:47:TYR:CE1	2.48	0.49
1:A:123:ARG:HD2	5:A:803:SO4:O4	2.14	0.48
4:D:216:ASP:N	4:D:216:ASP:OD1	2.47	0.47
2:B:187:GLU:OE1	2:B:189:ARG:NH1	2.47	0.47
1:A:98:GLU:OE1	6:A:601:EPE:H32	2.15	0.47
4:D:65:PHE:CD2	4:D:124:GLU:HA	2.51	0.46
4:D:65:PHE:CE2	4:D:124:GLU:HA	2.51	0.45
4:D:64:ILE:HD12	4:D:64:ILE:N	2.31	0.45
4:D:171:ASN:ND2	9:D:809:HOH:O	2.49	0.44
1:A:98:GLU:CD	6:A:601:EPE:H32	2.38	0.44
2:B:18:PHE:CD1	2:B:23:ARG:HG2	2.52	0.44
4:D:37:PHE:HE1	4:D:62:LEU:HD12	1.82	0.44
4:D:6:VAL:HG13	4:D:157:PHE:CD1	2.55	0.42
2:B:9:TRP:CH2	2:B:30:CYS:HB3	2.54	0.42
4:D:6:VAL:CG2	4:D:156:LYS:HD3	2.50	0.42
4:D:193:LYS:O	4:D:196:GLU:HG2	2.20	0.42
2:B:74:ALA:O	2:B:78:TYR:HB3	2.21	0.41
4:D:72:PRO:O	4:D:73:CYS:HB2	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	176/182 (97%)	173 (98%)	3 (2%)	0	100	100
2	B	188/190 (99%)	182 (97%)	6 (3%)	0	100	100
3	C	11/13 (85%)	11 (100%)	0	0	100	100
4	D	211/218 (97%)	201 (95%)	10 (5%)	0	100	100
All	All	586/603 (97%)	567 (97%)	19 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	163/166 (98%)	160 (98%)	3 (2%)	59 63
2	B	171/171 (100%)	162 (95%)	9 (5%)	22 18
3	C	12/12 (100%)	10 (83%)	2 (17%)	2 1
4	D	190/194 (98%)	176 (93%)	14 (7%)	13 9
All	All	536/543 (99%)	508 (95%)	28 (5%)	23 19

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	126	LYS
1	A	177	HIS
1	A	181	ASP
2	B	23	ARG
2	B	53	LEU
2	B	98	LYS
2	B	100	THR
2	B	105	LYS
2	B	106	THR
2	B	109	LEU
2	B	145	THR
2	B	189	ARG
3	C	310	LYS
3	C	316	LEU
4	D	22	LYS
4	D	37	PHE
4	D	56	LYS
4	D	76	LYS
4	D	85	SER
4	D	106	LYS
4	D	139	ILE
4	D	144	SER
4	D	172	ASP
4	D	174	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
4	D	187	VAL
4	D	193	LYS
4	D	198	ASN
4	D	216	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	149	HIS
1	A	177	HIS
2	B	156	GLN
4	D	8	ASN
4	D	42	ASN
4	D	171	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 8 ligands modelled in this entry, 1 is monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
7	DIO	A	702	-	6,6,6	0.45	0	6,6,6	0.68	0
5	SO4	D	801	-	4,4,4	0.23	0	6,6,6	0.19	0
5	SO4	A	802	-	4,4,4	0.16	0	6,6,6	0.26	0
7	DIO	A	701	-	6,6,6	0.42	0	6,6,6	1.00	0
5	SO4	C	804	-	4,4,4	0.15	0	6,6,6	0.22	0
5	SO4	A	803	-	4,4,4	0.19	0	6,6,6	0.16	0
6	EPE	A	601	-	15,15,15	1.90	1 (6%)	18,20,20	2.46	7 (38%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	DIO	A	702	-	-	-	0/1/1/1
7	DIO	A	701	-	-	-	0/1/1/1
6	EPE	A	601	-	-	2/9/19/19	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	601	EPE	C10-S	-7.08	1.67	1.77

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
6	A	601	EPE	C5-N4-C3	5.80	121.88	108.83
6	A	601	EPE	O3S-S-C10	4.24	112.63	105.77
6	A	601	EPE	C7-N4-C3	3.98	121.42	111.23
6	A	601	EPE	C5-C6-N1	3.83	118.50	110.64
6	A	601	EPE	O1S-S-C10	2.18	109.54	106.92
6	A	601	EPE	C6-N1-C2	2.15	113.67	108.83
6	A	601	EPE	C7-N4-C5	2.13	116.67	111.23

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	601	EPE	C8-C7-N4-C3
6	A	601	EPE	S-C10-C9-N1

There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	701	DIO	3	0
5	A	803	SO4	1	0
6	A	601	EPE	3	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	178/182 (97%)	0.01	3 (1%)	70	68	27, 33, 42, 51	0
2	B	190/190 (100%)	0.23	7 (3%)	41	41	24, 33, 42, 50	0
3	C	13/13 (100%)	0.37	1 (7%)	13	12	30, 33, 37, 38	0
4	D	213/218 (97%)	0.19	9 (4%)	36	35	23, 34, 43, 50	0
All	All	594/603 (98%)	0.15	20 (3%)	45	44	23, 33, 43, 51	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	106	THR	6.0
2	B	18	PHE	3.6
4	D	153	TYR	3.6
4	D	216	ASP	3.6
2	B	65	LYS	3.1
4	D	215	VAL	3.1
4	D	90	ASN	2.8
2	B	78	TYR	2.6
4	D	166	VAL	2.5
1	A	39	LYS	2.5
1	A	50	ARG	2.4
4	D	208	LEU	2.4
2	B	19	ASN	2.4
2	B	109	LEU	2.3
2	B	13	PHE	2.2
3	C	309	VAL	2.2
4	D	154	LYS	2.2
4	D	97	ILE	2.1
1	A	24	PHE	2.1
4	D	125	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
7	DIO	A	701	6/6	0.83	0.23	23,24,25,26	6
6	EPE	A	601	15/15	0.92	0.19	25,28,32,32	0
7	DIO	A	702	6/6	0.93	0.12	21,22,23,23	6
5	SO4	C	804	5/5	0.94	0.18	34,35,36,37	5
5	SO4	A	803	5/5	0.95	0.14	38,38,38,39	5
5	SO4	D	801	5/5	0.96	0.12	25,26,28,29	5
5	SO4	A	802	5/5	0.96	0.14	36,37,38,39	5
8	ZN	D	501	1/1	0.99	0.05	30,30,30,30	1

6.5 Other polymers (i)

There are no such residues in this entry.