Full wwPDB X-ray Structure Validation Report (i)

Jan 7, 2024 – 09:04 am GMT

PDB ID	:	6G98
Title	:	Three dimensional structure of human carbonic anhydrase IX in complex with sulfonamide
Authors	:	Leitans, J.; Tars, K.
Deposited on	:	2018-04-10
Resolution	:	2.47 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.47 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5857 (2.50-2.46)
Clashscore	141614	6594 (2.50-2.46)
Ramachandran outliers	138981	6469 (2.50-2.46)
Sidechain outliers	138945	6471 (2.50-2.46)
RSRZ outliers	127900	5738 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	257	.% 88% 5% • 6%
1	B	257	2% 87% 8% 5%
1	C	257	2% 88% • • 7%
1	D	257	8% 88% 5% 7%

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 7975 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Carbonic anhydrase 9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	241	1851	1176	328	343	4	0	0	0
1	B	243	1872	1192	331	345	4	0	0	0
1	C	239	1839	1168	326	341	4	0	0	0
1	D	239	1835	1165	325	341	4	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	3	PRO	ASP	conflict	UNP Q16790
A	41	SER	CYS	engineered mutation	UNP Q16790
B	3	PRO	ASP	conflict	UNP Q16790
B	41	SER	CYS	engineered mutation	UNP Q16790
C	3	PRO	ASP	conflict	UNP Q16790
C	41	SER	CYS	engineered mutation	UNP Q16790
D	3	PRO	ASP	conflict	UNP Q16790
D	41	SER	CYS	engineered mutation	UNP Q16790

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
2	A	1	1	1	0	0
2	B	1	1	1	0	0
2	C	1	1	1	0	0
2	D	1	1	1	0	0

- Molecule 3 is 4-[2-[3-(cyclododecylamino)-2,5,6-tris(fluoranyl)-4-sulfamoyl-phenyl]sulfanylethyl]benzoic acid (three-letter code: ER5) (formula: $C_{27}H_{35}F_3N_2O_4S_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	F	N	O	S		
3	A	1	38	27	3	2	4	2	0	0
3	B	1	38	27	3	2	4	2	0	0
3	C	1	38	27	3	2	4	2	0	0
3	D	1	38	27	3	2	4	2	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	146	146	146	0	0
4	B	98	98	98	0	0
4	C	90	90	90	0	0
4	D	88	88	88	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Carbonic anhydrase 9

Chain A:



• Molecule 1: Carbonic anhydrase 9

Chain B:



• Molecule 1: Carbonic anhydrase 9

Chain C:



• Molecule 1: Carbonic anhydrase 9

Chain D:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α, β, γ	152.39Å 152.39Å 171.83Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	76.19 – 2.47 72.00 – 2.47	Depositor EDS
% Data completeness (in resolution range)	99.8 (76.19-2.47) 99.7 (72.00-2.47)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.80 (at 2.48Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.169 , 0.209 0.176 , 0.211	Depositor DCC
R_{free} test set	2703 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å^2)	35.3	Xtriage
Anisotropy	0.038	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.36 , 39.7	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.023 for h,-h-k,-l	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	7975	wwPDB-VP
Average B, all atoms (Å^2)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.14% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ER5, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.96	2/1899 (0.1%)	0.77	2/2587 (0.1%)
1	B	0.52	0/1921	0.72	1/2615 (0.0%)
1	C	0.79	1/1886 (0.1%)	0.77	1/2566 (0.0%)
1	D	0.51	0/1881	0.71	0/2559
All	All	0.72	3/7587 (0.0%)	0.74	4/10327 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	82	GLY	C-N	29.16	1.89	1.34
1	C	125	THR	C-N	27.05	1.96	1.34
1	A	125	THR	C-N	19.14	1.78	1.34

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	C	125	THR	O-C-N	-14.45	99.58	122.70
1	A	125	THR	O-C-N	-8.75	108.70	122.70
1	A	82	GLY	O-C-N	-6.92	107.95	121.10
1	B	89	ARG	NE-CZ-NH1	5.54	123.07	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1851	0	1810	12	0
1	B	1872	0	1827	16	0
1	C	1839	0	1799	10	0
1	D	1835	0	1788	8	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	38	0	0	0	0
3	B	38	0	0	0	0
3	C	38	0	0	1	0
3	D	38	0	0	0	0
4	A	146	0	0	0	0
4	B	98	0	0	0	0
4	C	90	0	0	0	0
4	D	88	0	0	0	0
All	All	7975	0	7224	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:125:THR:C	1:A:127:ALA:N	1.78	1.36
1:A:82:GLY:C	1:A:84:PRO:N	1.89	1.25
1:C:125:THR:C	1:C:127:ALA:N	1.96	1.18
1:C:72:PRO:C	1:C:76:GLY:N	2.04	1.10
1:A:72:PRO:C	1:A:76:GLY:N	2.08	1.06
1:D:125:THR:C	1:D:127:ALA:N	2.10	1.04
1:D:72:PRO:C	1:D:76:GLY:N	2.14	1.00
1:B:28:GLN:HE21	1:B:205:GLN:HE21	1.08	0.95
1:D:82:GLY:C	1:D:84:PRO:N	2.21	0.94
1:B:72:PRO:C	1:B:76:GLY:N	2.20	0.94
1:B:82:GLY:C	1:B:84:PRO:N	2.25	0.90
1:C:82:GLY:C	1:C:84:PRO:N	2.25	0.88
1:B:125:THR:C	1:B:127:ALA:N	2.32	0.82
1:A:72:PRO:O	1:A:76:GLY:N	2.23	0.71
1:C:72:PRO:O	1:C:76:GLY:N	2.24	0.69
1:D:72:PRO:O	1:D:76:GLY:N	2.31	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:125:THR:O	1:B:127:ALA:N	2.31	0.64
1:D:125:THR:O	1:D:127:ALA:N	2.34	0.60
1:D:53:LEU:HD21	1:D:77:LEU:HD11	1.85	0.57
1:A:124:SER:HG	1:A:127:ALA:N	2.05	0.54
1:C:72:PRO:CA	1:C:76:GLY:N	2.69	0.53
1:B:53:LEU:HD21	1:B:77:LEU:HD11	1.90	0.53
1:A:125:THR:C	1:A:127:ALA:CA	2.72	0.52
1:A:82:GLY:CA	1:A:84:PRO:N	2.72	0.51
1:D:124:SER:HG	1:D:127:ALA:N	2.11	0.49
1:C:47:LEU:HD11	1:C:210:THR:HG21	1.93	0.49
1:A:53:LEU:HD21	1:A:77:LEU:HD11	1.95	0.48
1:B:28:GLN:NE2	1:B:205:GLN:HE21	1.92	0.48
1:A:88:TYR:HB3	1:A:122:HIS:HB3	1.96	0.47
1:B:72:PRO:CA	1:B:76:GLY:N	2.77	0.47
1:B:82:GLY:CA	1:B:84:PRO:N	2.77	0.47
1:B:143:VAL:HG11	1:B:209:TRP:CZ3	2.50	0.47
1:A:82:GLY:C	1:A:84:PRO:CA	2.79	0.46
1:B:124:SER:HG	1:B:127:ALA:N	2.14	0.46
1:B:72:PRO:O	1:B:76:GLY:N	2.49	0.44
1:A:72:PRO:CA	1:A:76:GLY:N	2.80	0.44
1:D:72:PRO:CA	1:D:76:GLY:N	2.82	0.43
1:C:125:THR:C	1:C:127:ALA:CA	2.82	0.42
1:C:82:GLY:C	1:C:84:PRO:CA	2.87	0.42
1:B:98:GLY:HA2	1:B:243:LEU:HD23	2.01	0.42
1:C:131:VAL:HG11	3:C:302:ER5:C23	2.50	0.41
1:C:72:PRO:HA	1:C:76:GLY:N	2.35	0.41
1:B:125:THR:O	1:B:127:ALA:CA	2.69	0.41
1:A:77:LEU:HD12	1:A:77:LEU:N	2.36	0.41
1:B:88:TYR:HB3	1:B:122:HIS:HB3	2.02	0.41
1:B:125:THR:N	1:B:127:ALA:N	2.69	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	235/257 (91%)	228 (97%)	7 (3%)	0	100	100
1	B	233/257 (91%)	227 (97%)	6 (3%)	0	100	100
1	C	231/257 (90%)	225 (97%)	6 (3%)	0	100	100
1	D	229/257 (89%)	223 (97%)	5 (2%)	1 (0%)	34	52
All	All	928/1028 (90%)	903 (97%)	24 (3%)	1 (0%)	51	71

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	238	ASP

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	195/210 (93%)	191 (98%)	4 (2%)	53	76
1	B	197/210 (94%)	195 (99%)	2 (1%)	76	89
1	C	194/210 (92%)	190 (98%)	4 (2%)	53	76
1	D	193/210 (92%)	191 (99%)	2 (1%)	76	89
All	All	779/840 (93%)	767 (98%)	12 (2%)	65	83

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	92	GLN
1	A	169	GLU
1	A	187	SER
1	A	213	ASN
1	B	92	GLN
1	B	213	ASN
1	C	36	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	84	PRO
1	C	92	GLN
1	C	132	ASP
1	D	60	ARG
1	D	92	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	36	GLN
1	B	159	GLN
1	B	205	GLN
1	C	213	ASN
1	D	159	GLN
1	D	175	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	ER5	D	302	2	39,40,40	3.01	7 (17%)	51,55,55	1.48	7 (13%)
3	ER5	B	302	2	39,40,40	3.02	7 (17%)	51,55,55	1.62	9 (17%)
3	ER5	A	302	2	39,40,40	2.90	7 (17%)	51,55,55	1.79	7 (13%)
3	ER5	C	302	2	39,40,40	2.94	7 (17%)	51,55,55	1.55	6 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ER5	D	302	2	-	6/34/34/34	0/3/3/3
3	ER5	B	302	2	-	15/34/34/34	1/3/3/3
3	ER5	A	302	2	-	14/34/34/34	1/3/3/3
3	ER5	C	302	2	-	9/34/34/34	1/3/3/3

All (28) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	302	ER5	C7-S37	-11.68	1.60	1.79
3	B	302	ER5	C7-S37	-10.45	1.62	1.79
3	C	302	ER5	C7-S37	-10.17	1.62	1.79
3	A	302	ER5	C7-S37	-10.17	1.62	1.79
3	A	302	ER5	C10-S30	-8.13	1.68	1.77
3	B	302	ER5	C8-C9	6.97	1.50	1.39
3	B	302	ER5	C10-C9	6.93	1.49	1.39
3	C	302	ER5	C10-S30	-6.56	1.70	1.77
3	D	302	ER5	C8-C9	6.52	1.50	1.39
3	C	302	ER5	C10-C11	6.19	1.48	1.39
3	B	302	ER5	C10-S30	-6.16	1.70	1.77
3	C	302	ER5	C10-C9	6.16	1.48	1.39
3	C	302	ER5	C12-C11	5.95	1.48	1.37
3	B	302	ER5	C10-C11	5.95	1.48	1.39
3	D	302	ER5	C10-C9	5.94	1.48	1.39
3	D	302	ER5	C10-S30	-5.86	1.71	1.77
3	C	302	ER5	C8-C9	5.81	1.48	1.39
3	D	302	ER5	C10-C11	5.80	1.47	1.39
3	C	302	ER5	C12-C7	5.77	1.49	1.39
3	A	302	ER5	C12-C7	5.72	1.49	1.39
3	B	302	ER5	C12-C7	5.62	1.49	1.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	302	ER5	C12-C11	5.61	1.47	1.37
3	A	302	ER5	C10-C9	5.60	1.47	1.39
3	A	302	ER5	C8-C9	5.55	1.48	1.39
3	D	302	ER5	C12-C7	5.36	1.48	1.39
3	D	302	ER5	C12-C11	5.33	1.47	1.37
3	A	302	ER5	C12-C11	4.97	1.46	1.37
3	A	302	ER5	C10-C11	4.94	1.46	1.39

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	A	302	ER5	O39-S37-O38	-6.57	107.95	118.76
3	D	302	ER5	O39-S37-O38	-6.34	108.33	118.76
3	B	302	ER5	O39-S37-O38	-5.75	109.31	118.76
3	C	302	ER5	O39-S37-O38	-5.60	109.55	118.76
3	B	302	ER5	C29-C18-N16	5.17	117.82	110.53
3	C	302	ER5	O39-S37-C7	4.93	114.62	107.30
3	A	302	ER5	C29-C18-N16	-4.89	103.64	110.53
3	A	302	ER5	C20-C19-C18	4.67	123.04	111.96
3	A	302	ER5	O38-S37-C7	4.50	113.99	107.30
3	B	302	ER5	O38-S37-C7	3.37	112.31	107.30
3	D	302	ER5	O38-S37-C7	3.18	112.03	107.30
3	D	302	ER5	F15-C9-C10	-2.95	115.93	119.86
3	B	302	ER5	C19-C18-N16	2.88	114.60	110.53
3	C	302	ER5	C20-C19-C18	2.83	118.68	111.96
3	D	302	ER5	C31-C32-C33	-2.72	109.08	113.97
3	B	302	ER5	O41-C39-O40	-2.68	117.41	123.35
3	C	302	ER5	O38-S37-N40	2.53	111.11	107.36
3	D	302	ER5	F15-C9-C8	2.51	123.31	119.56
3	C	302	ER5	O39-S37-N40	-2.45	103.73	107.36
3	B	302	ER5	C31-C32-C33	-2.37	109.70	113.97
3	D	302	ER5	O41-C39-O40	-2.29	118.26	123.35
3	A	302	ER5	C11-C10-C9	2.28	120.09	116.18
3	C	302	ER5	C31-C32-C33	-2.23	109.96	113.97
3	B	302	ER5	O41-C39-C36	2.23	120.64	114.85
3	A	302	ER5	C31-C32-C33	-2.22	109.98	113.97
3	A	302	ER5	O39-S37-N40	2.17	110.58	107.36
3	D	302	ER5	C7-S37-N40	2.06	112.07	108.26
3	B	302	ER5	C11-C10-S30	-2.03	117.33	121.80
3	B	302	ER5	C11-C10-C9	2.02	119.64	116.18

There are no chirality outliers.

All (44) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	302	ER5	C12-C7-S37-O39
3	A	302	ER5	C29-C18-C19-C20
3	A	302	ER5	C19-C18-C29-C28
3	B	302	ER5	C29-C18-N16-C8
3	B	302	ER5	C29-C18-C19-C20
3	B	302	ER5	C20-C21-C22-C23
3	B	302	ER5	C18-C19-C20-C21
3	C	302	ER5	C18-C19-C20-C21
3	A	302	ER5	N16-C18-C19-C20
3	A	302	ER5	N16-C18-C29-C28
3	B	302	ER5	N16-C18-C19-C20
3	B	302	ER5	C25-C26-C27-C28
3	C	302	ER5	C26-C27-C28-C29
3	A	302	ER5	C23-C24-C25-C26
3	A	302	ER5	C18-C19-C20-C21
3	B	302	ER5	C8-C7-S37-O39
3	C	302	ER5	C19-C20-C21-C22
3	B	302	ER5	C12-C7-S37-N40
3	B	302	ER5	C8-C7-S37-N40
3	D	302	ER5	C8-C7-S37-N40
3	C	302	ER5	C24-C25-C26-C27
3	C	302	ER5	C25-C26-C27-C28
3	A	302	ER5	C25-C26-C27-C28
3	D	302	ER5	C8-C7-S37-O39
3	D	302	ER5	C25-C26-C27-C28
3	B	302	ER5	C24-C25-C26-C27
3	A	302	ER5	C22-C23-C24-C25
3	A	302	ER5	C24-C25-C26-C27
3	C	302	ER5	C8-C7-S37-O38
3	B	302	ER5	C26-C27-C28-C29
3	A	302	ER5	C8-C7-S37-O39
3	A	302	ER5	C19-C20-C21-C22
3	A	302	ER5	C26-C27-C28-C29
3	B	302	ER5	C12-C7-S37-O38
3	C	302	ER5	C12-C7-S37-O38
3	D	302	ER5	C22-C23-C24-C25
3	C	302	ER5	C20-C21-C22-C23
3	B	302	ER5	C27-C28-C29-C18
3	A	302	ER5	C8-C7-S37-N40
3	C	302	ER5	C8-C7-S37-N40
3	D	302	ER5	C21-C22-C23-C24
3	D	302	ER5	C12-C7-S37-N40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	B	302	ER5	C22-C23-C24-C25
3	B	302	ER5	C7-C8-N16-C18

All (3) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	302	ER5	C18-C19-C20-C21-C22-C23-C24-C25-C26-C27-C28-C29
3	A	302	ER5	C18-C19-C20-C21-C22-C23-C24-C25-C26-C27-C28-C29
3	C	302	ER5	C18-C19-C20-C21-C22-C23-C24-C25-C26-C27-C28-C29

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	302	ER5	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand ER5 D 302



Ligand ER5 B 302



Ligand ER5 A 302



Ligand ER5 C 302



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	D	4
1	C	4
1	B	4
1	A	4

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	D	232:PRO	C	237:GLY	N	5.24

Continued on next page...

Continued from previous page...

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	C	232:PRO	C	237:GLY	N	3.33
1	B	232:PRO	C	237:GLY	N	3.17
1	A	232:PRO	C	237:GLY	N	3.11
1	B	125:THR	C	127:ALA	N	2.32
1	B	82:GLY	C	84:PRO	N	2.25
1	C	82:GLY	C	84:PRO	N	2.25
1	D	82:GLY	C	84:PRO	N	2.21
1	B	72:PRO	C	76:GLY	N	2.20
1	D	72:PRO	C	76:GLY	N	2.14
1	D	125:THR	C	127:ALA	N	2.10
1	A	72:PRO	C	76:GLY	N	2.08
1	C	72:PRO	C	76:GLY	N	2.04
1	C	125:THR	C	127:ALA	N	1.96
1	A	82:GLY	C	84:PRO	N	1.89
1	A	125:THR	C	127:ALA	N	1.78

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	241/257 (93%)	-0.08	3 (1%)	79	80	20, 30, 55, 118	0
1	B	243/257 (94%)	-0.08	4 (1%)	72	73	21, 35, 73, 127	0
1	C	239/257 (92%)	-0.06	6 (2%)	57	59	26, 40, 64, 107	0
1	D	239/257 (92%)	0.11	21 (8%)	10	9	25, 42, 70, 110	0
All	All	962/1028 (93%)	-0.03	34 (3%)	44	46	20, 37, 67, 127	0

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	237	GLY	9.5
1	C	237	GLY	6.6
1	B	16	TRP	5.4
1	A	18	ARG	4.8
1	D	238	ASP	4.4
1	B	237	GLY	4.0
1	D	230	TRP	4.0
1	C	238	ASP	3.9
1	C	152	GLU	3.4
1	D	239	SER	3.3
1	B	19	VAL	3.1
1	D	169	GLU	3.1
1	D	100	ALA	2.9
1	D	227(A)	ASP	2.9
1	D	240	ARG	2.9
1	D	101	GLY	2.6
1	A	237	GLY	2.6
1	D	243	LEU	2.6
1	D	64	HIS	2.6
1	C	227(A)	ASP	2.5
1	D	231	GLY	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	224	HIS	2.4
1	B	230	TRP	2.4
1	D	165	GLU	2.3
1	D	229	LEU	2.3
1	D	232	PRO	2.3
1	A	19	VAL	2.3
1	D	102	ARG	2.1
1	D	241	LEU	2.1
1	C	151(A)	PRO	2.1
1	C	100	ALA	2.1
1	D	242	GLN	2.1
1	D	174	THR	2.0
1	D	245	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	ER5	D	302	38/38	0.94	0.19	39,47,54,58	0
3	ER5	B	302	38/38	0.95	0.20	38,51,74,74	0
3	ER5	A	302	38/38	0.95	0.15	38,49,56,56	0
3	ER5	C	302	38/38	0.96	0.18	39,45,65,67	0
2	ZN	D	301	1/1	0.99	0.09	34,34,34,34	0
2	ZN	C	301	1/1	1.00	0.11	29,29,29,29	0
2	ZN	A	301	1/1	1.00	0.13	25,25,25,25	0
2	ZN	B	301	1/1	1.00	0.11	27,27,27,27	0

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ER5 D 302:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







Electron density around ER5 B 302:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around ER5 A 302:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around ER5 C 302:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







6.5 Other polymers (i)

There are no such residues in this entry.