wwPDB X-ray Structure Validation Summary Report (i)

Aug 26, 2023 – 11:34 PM EDT

PDB ID	:	3GFQ
Title	:	Structure of YhdA, K109L variant
Authors	:	Staunig, N.; Gruber, K.
Deposited on	:	2009-02-27
Resolution	:	3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	174	3% 71% 24% • •
1	B	174	2% 74% 20% • •
1	C	174	9% 67% 25% • •
1	D	174	4% 70% 26% • •

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 5248 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FMN-dependent NADPH-azoreductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	168	1284	819	219	241	5	0	0	0
1	B	167	1280	817	218	240	5	0	0	0
1	C	167	1280	817	218	240	5	0	0	0
1	D	167	1280	817	218	240	5	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	109	LEU	LYS	engineered mutation	UNP O07529
B	109	LEU	LYS	engineered mutation	UNP O07529
C	109	LEU	LYS	engineered mutation	UNP O07529
D	109	LEU	LYS	engineered mutation	UNP O07529

- Molecule 2 is FLAVIN MONONUCLEOTIDE (three-letter code: FMN) (formula: $C_{17}H_{21}N_4O_9P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
2	B	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
2	C	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
2	D	1	Total	C	N	O	P	0	0
			31	17	4	9	1		

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: FMN-dependent NADPH-azoreductase

Chain A:



• Molecule 1: FMN-dependent NADPH-azoreductase

Chain B:



• Molecule 1: FMN-dependent NADPH-azoreductase

Chain C:



• Molecule 1: FMN-dependent NADPH-azoreductase

Chain D:


MET ASN M3
N7 G8 T9 P10
H13 G14 R15 T16
S21 Y22 I23 A24
T29
L34
L39
N43
A46 E47 Q48 S49 E50 L51
L57 K58 Q59 R60 V61
L70 S71 P72 E73 Y74
S79 G80 A81
S91 E92 Q93 F94 K95 Y96
L102
A114
L115 N116 N117 M118
R123
V135 L136 D137 P138 V139 H140
E144
N151 I152 R153 E154 S155 I156
V160
L163 S164
K168 A169 GLY ASN PRO GLY VAL

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, α, β, γ	47.54Å 66.24Å 220.75Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.10 – 3.00 29.10 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.5 (29.10-3.00) 99.5 (29.10-3.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.18	Depositor
$< I/\sigma(I) >$ [1]	1.73 (at 3.00Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.222 , 0.274 0.219 , 0.268	Depositor DCC
R_{free} test set	747 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å^2)	59.7	Xtriage
Anisotropy	0.457	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.36 , 45.6	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.93	EDS
Total number of atoms	5248	wwPDB-VP
Average B, all atoms (Å^2)	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.60% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: FMN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.23	0/1303	0.40	0/1762
1	B	0.24	0/1299	0.41	0/1757
1	C	0.23	0/1299	0.41	0/1757
1	D	0.24	0/1299	0.41	0/1757
All	All	0.24	0/5200	0.41	0/7033

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1284	0	1324	33	0
1	B	1280	0	1321	28	0
1	C	1280	0	1321	36	0
1	D	1280	0	1321	30	0
2	A	31	0	19	2	0
2	B	31	0	19	1	0
2	C	31	0	19	1	0
2	D	31	0	19	2	0
All	All	5248	0	5363	121	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 121 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:136:LEU:HD22	1:A:152:ILE:HD11	1.58	0.84
1:A:109:LEU:H	1:A:109:LEU:HD23	1.43	0.82
1:B:131:PRO:HG3	1:D:123:ARG:HD2	1.70	0.72
1:D:114:ALA:O	1:D:118:MET:HG3	1.92	0.70
1:C:7:ASN:OD1	1:C:9:THR:HG22	1.90	0.69

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	166/174 (95%)	161 (97%)	4 (2%)	1 (1%)	25	64
1	B	165/174 (95%)	159 (96%)	4 (2%)	2 (1%)	13	48
1	C	165/174 (95%)	158 (96%)	7 (4%)	0	100	100
1	D	165/174 (95%)	157 (95%)	6 (4%)	2 (1%)	13	48
All	All	661/696 (95%)	635 (96%)	21 (3%)	5 (1%)	19	57

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	143	VAL
1	D	22	TYR
1	B	107	GLY
1	D	23	ILE
1	B	108	GLY

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	138/143 (96%)	135 (98%)	3 (2%)	52	81
1	B	138/143 (96%)	133 (96%)	5 (4%)	35	70
1	C	138/143 (96%)	127 (92%)	11 (8%)	12	40
1	D	138/143 (96%)	135 (98%)	3 (2%)	52	81
All	All	552/572 (96%)	530 (96%)	22 (4%)	31	68

5 of 22 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	117	ASN
1	C	148	VAL
1	C	140	HIS
1	C	152	ILE
1	B	151	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	55	GLN
1	B	140	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	FMN	D	200	-	33,33,33	1.10	2 (6%)	48,50,50	1.23	6 (12%)
2	FMN	A	200	-	33,33,33	1.09	2 (6%)	48,50,50	1.29	6 (12%)
2	FMN	C	200	-	33,33,33	1.08	2 (6%)	48,50,50	1.19	6 (12%)
2	FMN	B	200	-	33,33,33	1.07	2 (6%)	48,50,50	1.28	7 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FMN	D	200	-	-	8/18/18/18	0/3/3/3
2	FMN	A	200	-	-	5/18/18/18	0/3/3/3
2	FMN	C	200	-	-	3/18/18/18	0/3/3/3
2	FMN	B	200	-	-	5/18/18/18	0/3/3/3

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	200	FMN	C4A-N5	4.04	1.38	1.30
2	C	200	FMN	C4A-N5	3.98	1.38	1.30
2	A	200	FMN	C4A-N5	3.95	1.38	1.30
2	D	200	FMN	C4A-N5	3.91	1.38	1.30
2	D	200	FMN	C10-N1	2.63	1.38	1.33

The worst 5 of 25 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	C	200	FMN	C4-N3-C2	-2.86	120.36	125.64
2	A	200	FMN	C9A-C5A-N5	-2.83	119.35	122.43
2	D	200	FMN	C9A-C5A-N5	-2.81	119.38	122.43
2	D	200	FMN	C4-N3-C2	-2.80	120.48	125.64
2	B	200	FMN	C10-C4A-N5	-2.77	118.99	124.86

There are no chirality outliers.

5 of 21 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	200	FMN	C3'-C4'-C5'-O5'
2	A	200	FMN	O4'-C4'-C5'-O5'
2	A	200	FMN	C5'-O5'-P-O1P
2	A	200	FMN	C5'-O5'-P-O2P
2	A	200	FMN	C5'-O5'-P-O3P

There are no ring outliers.

4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	200	FMN	2	0
2	A	200	FMN	2	0
2	C	200	FMN	1	0
2	B	200	FMN	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand FMN D 200



Bond lengths



Bond angles



Torsions



Rings

Ligand FMN A 200



Bond lengths



Bond angles



Torsions



Rings

Ligand FMN C 200



Bond lengths



Bond angles



Torsions



Rings

Ligand FMN B 200



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	168/174 (96%)	0.18	6 (3%)	42	17	30, 53, 99, 128	0
1	B	167/174 (95%)	0.15	4 (2%)	59	30	28, 50, 90, 138	0
1	C	167/174 (95%)	0.50	16 (9%)	8	2	33, 68, 109, 172	0
1	D	167/174 (95%)	0.11	7 (4%)	36	14	25, 49, 77, 133	0
All	All	669/696 (96%)	0.24	33 (4%)	29	11	25, 55, 100, 172	0

The worst 5 of 33 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	106	GLY	7.3
1	C	108	GLY	6.8
1	B	96	TYR	3.7
1	C	109	LEU	3.6
1	C	157	LYS	3.6

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	FMN	B	200	31/31	0.94	0.16	57,57,57,57	0
2	FMN	A	200	31/31	0.95	0.15	43,43,43,43	0
2	FMN	C	200	31/31	0.96	0.17	54,54,54,54	0
2	FMN	D	200	31/31	0.96	0.17	55,55,55,55	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FMN B 200:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative) and green (positive)



Electron density around FMN A 200:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around FMN C 200:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around FMN D 200:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







6.5 Other polymers (i)

There are no such residues in this entry.