



wwPDB X-ray Structure Validation Summary Report

Apr 28, 2024 – 06:55 am BST

PDB ID : 1GVI
Title : Thermus maltogenic amylase in complex with beta-CD
Authors : Kim, M.-S.; Kim, J.-I.; Oh, B.-H.
Deposited on : 2002-02-14
Resolution : 3.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1205 (3.34-3.26)
Ramachandran outliers	138981	1183 (3.34-3.26)
Sidechain outliers	138945	1182 (3.34-3.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	588	43% (green), 48% (yellow), 9% (orange), 0% (red)
1	B	588	41% (green), 49% (yellow), 9% (orange), 0% (red)
2	C	7	71% (yellow), 29% (orange)
2	D	7	86% (yellow), 14% (orange)

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 9810 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called MALTOGENIC AMYLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	588	4828	3119	819	869	21	0	0	0
1	B	588	4828	3119	819	869	21	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	357	LEU	GLU	conflict	UNP O69007
B	357	LEU	GLU	conflict	UNP O69007

- Molecule 2 is an oligosaccharide called Cycloheptakis-(1-4)-(alpha-D-glucopyranose).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	C	7	77	42	35	0	0	0
2	D	7	77	42	35	0	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: MALTOGENIC AMYLASE


Chain A: 



- Molecule 1: MALTOGENIC AMYLASE

Chain B: 






● Molecule 2: Cycloheptakis-(1-4)-(alpha-D-glucopyranose)



● Molecule 2: Cycloheptakis-(1-4)-(alpha-D-glucopyranose)



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	119.09Å 119.09Å 270.20Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 3.30	Depositor
% Data completeness (in resolution range)	89.3 (20.00-3.30)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS	Depositor
R, R_{free}	0.192 , 0.243	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	9810	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.41	0/4982	0.67	0/6770
1	B	0.42	0/4982	0.68	1/6770 (0.0%)
All	All	0.41	0/9964	0.68	1/13540 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	170	PHE	N-CA-C	6.04	127.32	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4828	0	4629	347	0
1	B	4828	0	4629	349	0
2	C	77	0	63	2	0
2	D	77	0	63	3	0
All	All	9810	0	9384	683	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 36.

The worst 5 of 683 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:281:GLU:HB2	1:A:282:PRO:HD3	1.33	1.08
1:B:281:GLU:HB2	1:B:282:PRO:CD	1.93	0.99
1:B:275:GLU:HG3	1:B:277:PRO:HD2	1.46	0.97
1:B:155:LYS:HG3	1:B:170:PHE:HB2	1.47	0.96
1:A:253:ALA:HB3	1:A:254:PRO:HD3	1.44	0.95

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	586/588 (100%)	486 (83%)	76 (13%)	24 (4%)	3	17
1	B	586/588 (100%)	467 (80%)	93 (16%)	26 (4%)	2	16
All	All	1172/1176 (100%)	953 (81%)	169 (14%)	50 (4%)	2	16

5 of 50 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	80	ARG
1	A	118	PRO
1	A	156	GLY
1	A	158	ARG
1	A	159	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	505/505 (100%)	452 (90%)	53 (10%)	7	25
1	B	505/505 (100%)	457 (90%)	48 (10%)	8	29
All	All	1010/1010 (100%)	909 (90%)	101 (10%)	7	27

5 of 101 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	60	ARG
1	B	200	ARG
1	B	570	THR
1	B	67	LEU
1	B	147	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 41 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	299	ASN
1	B	418	ASN
1	B	302	HIS
1	B	360	HIS
1	B	482	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

14 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLC	C	1	2	11,11,12	1.52	3 (27%)	15,15,17	1.33	2 (13%)
2	GLC	C	2	2	11,11,12	1.60	4 (36%)	15,15,17	1.89	3 (20%)
2	GLC	C	3	2	11,11,12	0.86	0	15,15,17	1.83	5 (33%)
2	GLC	C	4	2	11,11,12	1.21	2 (18%)	15,15,17	0.57	0
2	GLC	C	5	2	11,11,12	1.18	1 (9%)	15,15,17	1.06	1 (6%)
2	GLC	C	6	2	11,11,12	1.34	1 (9%)	15,15,17	0.94	1 (6%)
2	GLC	C	7	2	11,11,12	1.60	4 (36%)	15,15,17	1.06	2 (13%)
2	GLC	D	1	2	11,11,12	1.57	2 (18%)	15,15,17	1.32	1 (6%)
2	GLC	D	2	2	11,11,12	1.61	4 (36%)	15,15,17	1.88	3 (20%)
2	GLC	D	3	2	11,11,12	1.08	0	15,15,17	1.81	4 (26%)
2	GLC	D	4	2	11,11,12	1.04	0	15,15,17	0.57	0
2	GLC	D	5	2	11,11,12	1.09	1 (9%)	15,15,17	0.99	1 (6%)
2	GLC	D	6	2	11,11,12	1.31	1 (9%)	15,15,17	1.10	1 (6%)
2	GLC	D	7	2	11,11,12	1.82	5 (45%)	15,15,17	1.03	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	C	1	2	-	1/2/19/22	0/1/1/1
2	GLC	C	2	2	-	0/2/19/22	0/1/1/1
2	GLC	C	3	2	-	0/2/19/22	0/1/1/1
2	GLC	C	4	2	-	0/2/19/22	0/1/1/1
2	GLC	C	5	2	-	2/2/19/22	0/1/1/1
2	GLC	C	6	2	-	0/2/19/22	0/1/1/1
2	GLC	C	7	2	-	2/2/19/22	0/1/1/1
2	GLC	D	1	2	-	2/2/19/22	0/1/1/1
2	GLC	D	2	2	-	2/2/19/22	0/1/1/1
2	GLC	D	3	2	-	0/2/19/22	0/1/1/1
2	GLC	D	4	2	-	0/2/19/22	0/1/1/1
2	GLC	D	5	2	-	2/2/19/22	0/1/1/1
2	GLC	D	6	2	-	1/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	D	7	2	-	0/2/19/22	0/1/1/1

The worst 5 of 28 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	7	GLC	O4-C4	3.01	1.50	1.43
2	D	7	GLC	O5-C5	2.64	1.48	1.43
2	D	1	GLC	O5-C5	2.52	1.48	1.43
2	C	7	GLC	O4-C4	2.50	1.48	1.43
2	D	1	GLC	C4-C5	2.50	1.58	1.53

The worst 5 of 26 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	2	GLC	O4-C4-C3	-5.48	97.69	110.35
2	C	2	GLC	O4-C4-C3	-5.40	97.86	110.35
2	D	3	GLC	O4-C4-C5	-4.14	99.02	109.30
2	D	1	GLC	C1-O5-C5	4.00	117.61	112.19
2	C	3	GLC	O4-C4-C5	-3.99	99.39	109.30

There are no chirality outliers.

5 of 12 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	D	5	GLC	C4-C5-C6-O6
2	D	2	GLC	C4-C5-C6-O6
2	D	1	GLC	O5-C5-C6-O6
2	D	5	GLC	O5-C5-C6-O6
2	D	2	GLC	O5-C5-C6-O6


There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1	GLC	1	0
2	C	4	GLC	1	0
2	D	5	GLC	2	0
2	D	4	GLC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.