wwPDB X-ray Structure Validation Summary Report (i)

Mar 23, 2024 – 12:49 PM EDT

PDB ID	:	4H25
Title	:	TCR interaction with peptide mimics of nickel offers structure insights to nickel contact allergy
Authors	:	Kappler, J.W.; Yin, L.; Dai, S.; Marrack, P.; Crawford, F.
Deposited on	:	2012-09-12
Resolution	:	2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	180	3% 81% 19%
1	D	180	7% 81% 19%
2	B	188	2% 72% 27% •
2	E	188	4% 72% 24% • •
3	C	22	9% 73% 23% 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	22	9% 77% 23%
4	G	2	50% 50%
4	H	2	100%
4	I	2	100%
4	J	2	100%
4	K	2	50% 50%
4	L	2	50% 50%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NAG	G	2	-	-	-	X
4	NAG	L	2	-	-	-	X

2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 6910 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class II histocompatibility antigen, DR alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	180	1479	957	240	277	5	0	0	0
1	D	180	1479	957	240	277	5	0	0	0

- Molecule 2 is a protein called MHC class II antigen.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	188	1549	978	279	287	5	0	0	0
2	E	182	1496	945	267	279	5	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	3	THR	-	expression tag	UNP B8YAC7
B	4	ARG	-	expression tag	UNP B8YAC7
B	5	PRO	-	expression tag	UNP B8YAC7
B	77	THR	ASN	conflict	UNP B8YAC7
B	189	ARG	-	expression tag	UNP B8YAC7
B	190	ALA	-	expression tag	UNP B8YAC7
E	3	THR	-	expression tag	UNP B8YAC7
E	4	ARG	-	expression tag	UNP B8YAC7
E	5	PRO	-	expression tag	UNP B8YAC7
E	77	THR	ASN	conflict	UNP B8YAC7
E	189	ARG	-	expression tag	UNP B8YAC7
E	190	ALA	-	expression tag	UNP B8YAC7

- Molecule 3 is a protein called peptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	22	174	109	37	27	1	0	0	0
			Total	C	N	O	S			
3	F	22	174	109	37	27	1	0	0	0

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	G	2	28	16	2	10	0	0	0
			Total	C	N	O			
4	H	2	28	16	2	10	0	0	0
			Total	C	N	O			
4	I	2	28	16	2	10	0	0	0
			Total	C	N	O			
4	J	2	28	16	2	10	0	0	0
			Total	C	N	O			
4	K	2	28	16	2	10	0	0	0
			Total	C	N	O			
4	L	2	28	16	2	10	0	0	0

- Molecule 5 is ISOPROPYL ALCOHOL (three-letter code: IPA) (formula: C_3H_8O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
5	A	1	4	3	1	0	0
			Total	C	O		
5	A	1	4	3	1	0	0
			Total	C	O		
5	B	1	4	3	1	0	0
			Total	C	O		
5	E	1	4	3	1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	A	104	104	104	0	0
			Total	O		
6	B	114	114	114	0	0
			Total	O		
6	C	15	15	15	0	0
			Total	O		
6	D	63	63	63	0	0
			Total	O		
6	E	58	58	58	0	0
			Total	O		
6	F	21	21	21	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: HLA class II histocompatibility antigen, DR alpha chain

Chain A:



• Molecule 1: HLA class II histocompatibility antigen, DR alpha chain

Chain D:



• Molecule 2: MHC class II antigen

Chain B:



• Molecule 2: MHC class II antigen

Chain E:



• Molecule 3: peptide

Chain C:



• Molecule 3: peptide

Chain F:



• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:

• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:



• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:



• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J:



NAG1
NAG2

• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain K: 50% 50%

NAG1
NAG2

• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain L: 50% 50%

NAG1
NAG2

4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α, β, γ	154.52Å 97.18Å 67.10Å 90.00° 105.36° 90.00°	Depositor
Resolution (Å)	40.00 – 2.20 47.42 – 2.20	Depositor EDS
% Data completeness (in resolution range)	(Not available) (40.00-2.20) 93.5 (47.42-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.50 (at 2.20Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.210 , 0.255 0.205 , 0.213	Depositor DCC
R_{free} test set	2317 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å^2)	33.8	Xtriage
Anisotropy	0.563	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.32 , 47.1	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	6910	wwPDB-VP
Average B, all atoms (Å^2)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.01% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: IPA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.40	0/1524	0.61	0/2077
1	D	0.36	0/1524	0.56	0/2077
2	B	0.43	0/1591	0.65	0/2160
2	E	0.37	0/1534	0.57	0/2080
3	C	0.31	0/177	0.59	0/237
3	F	0.36	0/177	0.62	0/237
All	All	0.38	0/6527	0.60	0/8868

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1479	0	1410	29	0
1	D	1479	0	1410	34	0
2	B	1549	0	1479	51	0
2	E	1496	0	1430	48	0
3	C	174	0	193	12	0
3	F	174	0	193	5	0
4	G	28	0	25	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	H	28	0	25	2	0
4	I	28	0	25	0	0
4	J	28	0	25	2	0
4	K	28	0	25	3	0
4	L	28	0	25	1	0
5	A	8	0	16	2	0
5	B	4	0	8	1	0
5	E	4	0	8	1	0
6	A	104	0	0	3	0
6	B	114	0	0	4	0
6	C	15	0	0	1	0
6	D	63	0	0	2	0
6	E	58	0	0	4	0
6	F	21	0	0	1	0
All	All	6910	0	6297	161	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 161 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:87:PRO:HD2	1:D:170:LEU:HD11	1.44	0.98
1:D:167:HIS:HB3	1:D:170:LEU:HD13	1.62	0.81
2:E:55:ARG:HB2	2:E:56:PRO:HD3	1.64	0.78
3:C:327:ARG:HE	3:C:327:ARG:H	1.29	0.77
2:B:52:GLU:HG2	2:B:55:ARG:HH12	1.51	0.76

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	178/180 (99%)	174 (98%)	4 (2%)	0	100	100
1	D	178/180 (99%)	170 (96%)	8 (4%)	0	100	100
2	B	186/188 (99%)	179 (96%)	5 (3%)	2 (1%)	14	12
2	E	178/188 (95%)	170 (96%)	8 (4%)	0	100	100
3	C	20/22 (91%)	20 (100%)	0	0	100	100
3	F	20/22 (91%)	20 (100%)	0	0	100	100
All	All	760/780 (97%)	733 (96%)	25 (3%)	2 (0%)	41	46

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	109	LEU
2	B	110	GLN

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	164/164 (100%)	162 (99%)	2 (1%)	71	83
1	D	164/164 (100%)	160 (98%)	4 (2%)	49	62
2	B	172/172 (100%)	170 (99%)	2 (1%)	71	83
2	E	166/172 (96%)	164 (99%)	2 (1%)	71	83
3	C	20/20 (100%)	19 (95%)	1 (5%)	24	30
3	F	20/20 (100%)	18 (90%)	2 (10%)	7	7
All	All	706/712 (99%)	693 (98%)	13 (2%)	59	72

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	171	ASP
1	D	181	ASP
3	F	325	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	E	67	LEU
3	F	324	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
2	E	34	GLN
2	E	113	ASN
2	E	150	ASN
2	E	96	HIS
2	B	156	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

12 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	NAG	G	1	1,4	14,14,15	0.59	0	17,19,21	0.75	0
4	NAG	G	2	4	14,14,15	0.50	0	17,19,21	0.69	1 (5%)
4	NAG	H	1	1,4	14,14,15	0.56	0	17,19,21	0.69	0
4	NAG	H	2	4	14,14,15	0.47	0	17,19,21	0.73	1 (5%)
4	NAG	I	1	2,4	14,14,15	0.55	0	17,19,21	0.76	1 (5%)
4	NAG	I	2	4	14,14,15	0.41	0	17,19,21	0.70	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	NAG	J	1	1,4	14,14,15	0.45	0	17,19,21	0.64	0
4	NAG	J	2	4	14,14,15	0.45	0	17,19,21	0.68	0
4	NAG	K	1	1,4	14,14,15	0.71	0	17,19,21	0.96	2 (11%)
4	NAG	K	2	4	14,14,15	0.51	0	17,19,21	0.63	0
4	NAG	L	1	2,4	14,14,15	0.61	0	17,19,21	0.80	1 (5%)
4	NAG	L	2	4	14,14,15	0.53	0	17,19,21	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	G	1	1,4	-	4/6/23/26	0/1/1/1
4	NAG	G	2	4	-	2/6/23/26	0/1/1/1
4	NAG	H	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	H	2	4	-	4/6/23/26	0/1/1/1
4	NAG	I	1	2,4	-	2/6/23/26	0/1/1/1
4	NAG	I	2	4	-	5/6/23/26	0/1/1/1
4	NAG	J	1	1,4	-	5/6/23/26	0/1/1/1
4	NAG	J	2	4	-	3/6/23/26	0/1/1/1
4	NAG	K	1	1,4	-	4/6/23/26	0/1/1/1
4	NAG	K	2	4	-	2/6/23/26	0/1/1/1
4	NAG	L	1	2,4	-	2/6/23/26	0/1/1/1
4	NAG	L	2	4	-	3/6/23/26	0/1/1/1

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
4	H	2	NAG	C2-N2-C7	-2.33	119.59	122.90
4	L	1	NAG	C2-N2-C7	-2.26	119.69	122.90
4	K	1	NAG	C4-C3-C2	2.24	114.31	111.02
4	K	1	NAG	C2-N2-C7	-2.22	119.74	122.90
4	G	2	NAG	C2-N2-C7	-2.12	119.88	122.90

There are no chirality outliers.

5 of 36 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	G	1	NAG	C8-C7-N2-C2
4	G	1	NAG	O7-C7-N2-C2
4	G	2	NAG	C8-C7-N2-C2
4	G	2	NAG	O7-C7-N2-C2
4	H	2	NAG	C8-C7-N2-C2

There are no ring outliers.

9 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	L	1	NAG	1	0
4	G	1	NAG	1	0
4	K	2	NAG	3	0
4	H	1	NAG	2	0
4	L	2	NAG	1	0
4	G	2	NAG	1	0
4	J	2	NAG	2	0
4	K	1	NAG	2	0
4	J	1	NAG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

Oligosaccharide Chain G



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain H



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain I



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain J



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain K



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain L



5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
5	IPA	E	203	-	3,3,3	0.64	0	3,3,3	0.33	0
5	IPA	A	205	-	3,3,3	0.70	0	3,3,3	0.34	0
5	IPA	A	206	-	3,3,3	0.63	0	3,3,3	0.34	0
5	IPA	B	203	-	3,3,3	0.66	0	3,3,3	0.33	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	E	203	IPA	1	0
5	A	205	IPA	2	0
5	B	203	IPA	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	180/180 (100%)	0.09	5 (2%)	53	51	22, 37, 74, 115	0
1	D	180/180 (100%)	0.39	12 (6%)	17	16	23, 48, 89, 128	0
2	B	188/188 (100%)	-0.02	3 (1%)	72	70	21, 31, 62, 93	0
2	E	182/188 (96%)	0.27	8 (4%)	34	32	27, 45, 73, 103	0
3	C	22/22 (100%)	0.24	2 (9%)	9	8	22, 43, 66, 102	0
3	F	22/22 (100%)	0.23	2 (9%)	9	8	25, 37, 72, 98	0
All	All	774/780 (99%)	0.18	32 (4%)	37	35	21, 40, 78, 128	0

The worst 5 of 32 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	182	ALA	10.9
1	D	182	ALA	8.2
1	D	181	ASP	5.6
1	D	100	ARG	4.4
2	E	165	PRO	3.9

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	NAG	G	2	14/15	0.43	0.74	179,190,193,193	0
4	NAG	K	2	14/15	0.49	0.38	143,148,149,149	0
4	NAG	K	1	14/15	0.50	0.39	117,128,130,137	0
4	NAG	L	2	14/15	0.51	0.50	134,141,142,143	0
4	NAG	I	1	14/15	0.58	0.30	100,109,112,119	0
4	NAG	G	1	14/15	0.60	0.39	129,139,148,162	0
4	NAG	I	2	14/15	0.66	0.36	127,132,134,134	0
4	NAG	L	1	14/15	0.73	0.28	103,112,116,125	0
4	NAG	J	2	14/15	0.74	0.33	99,102,103,104	0
4	NAG	H	2	14/15	0.77	0.28	75,79,81,81	0
4	NAG	J	1	14/15	0.84	0.24	81,88,90,94	0
4	NAG	H	1	14/15	0.93	0.13	51,56,61,68	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain G:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain H:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain I:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain J:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around Chain K:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain L:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
5	IPA	B	203	4/4	0.59	0.27	39,40,42,43	0
5	IPA	A	205	4/4	0.78	0.25	40,41,43,44	0
5	IPA	E	203	4/4	0.83	0.22	43,44,45,46	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
5	IPA	A	206	4/4	0.86	0.17	45,47,47,47	0

6.5 Other polymers (i)

There are no such residues in this entry.