



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 27, 2023 – 11:07 AM EDT

PDB ID : 3H9U
Title : S-adenosyl homocysteine hydrolase (SAHH) from Trypanosoma brucei
Authors : Siponen, M.I.; Schutz, P.; Arrowsmith, C.H.; Berglund, H.; Bountra, C.; Collins, R.; Edwards, A.M.; Flodin, S.; Flores, A.; Graslund, S.; Hammarstrom, M.; Johansson, A.; Johansson, I.; Karlberg, T.; Kragh Nielsen, T.; Kotenyova, T.; Kotzsch, A.; Moche, M.; Nordlund, P.; Nyman, T.; Persson, C.; Sagemark, J.; Thorsell, A.G.; Tresaugues, L.; Van Den Berg, S.; Weigelt, J.; Welin, M.; Wisniewska, M.; Schueler, H.; Structural Genomics Consortium (SGC)
Deposited on : 2009-04-30
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	436	 15% 81% 17% ..
1	B	436	 20% 82% 16% ..

Continued on next page...

CCP4 : 7.0.044 (Gargrove)
 Ideal geometry (proteins) : Engh & Huber (2001)
 Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
 Validation Pipeline (wwPDB-VP) : 2.35

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	C	436	
1	D	436	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	PG4	D	1	-	-	X	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 14413 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Adenosylhomocysteinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	430	3324	2098	573	628	25	0	0	0
1	B	430	3336	2108	578	625	25	0	0	0
1	C	435	3378	2130	584	638	26	0	2	0
1	D	434	3381	2133	583	640	25	0	3	0


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP Q383X0
A	-1	MET	-	expression tag	UNP Q383X0
B	-2	SER	-	expression tag	UNP Q383X0
B	-1	MET	-	expression tag	UNP Q383X0
C	-2	SER	-	expression tag	UNP Q383X0
C	-1	MET	-	expression tag	UNP Q383X0
D	-2	SER	-	expression tag	UNP Q383X0
D	-1	MET	-	expression tag	UNP Q383X0

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: $C_{21}H_{27}N_7O_{14}P_2$).


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	N	0	0
			10	5	5		
3	B	1	Total	C	N	0	0
			10	5	5		
3	C	1	Total	C	N	0	0
			10	5	5		
3	D	1	Total	C	N	0	0
			10	5	5		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	C	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: $C_8H_{18}O_5$).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	D	1	Total	C O	0	0
			13	8 5		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	66	Total	O	0	0
			66	66		
6	B	70	Total	O	0	0
			70	70		
6	C	308	Total	O	0	0
			308	308		
6	D	315	Total	O	0	0
			315	315		



- Molecule 1: Adenosylhomocysteinase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	91.35Å 136.05Å 191.87Å 90.00° 99.32° 90.00°	Depositor
Resolution (Å)	45.55 – 1.90 45.54 – 1.90	Depositor EDS
% Data completeness (in resolution range)	100.0 (45.55-1.90) 99.9 (45.54-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.70 (at 1.89Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.196 , 0.231 0.201 , 0.235	Depositor DCC
R_{free} test set	9087 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	20.8	Xtrriage
Anisotropy	0.047	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 50.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	14413	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 34.18 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 7.1036e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAD, PG4, GOL, ADN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.53	0/3386	0.64	0/4580
1	B	0.53	0/3398	0.62	0/4592
1	C	0.86	1/3443 (0.0%)	0.89	12/4652 (0.3%)
1	D	0.84	1/3450 (0.0%)	0.84	8/4663 (0.2%)
All	All	0.71	2/13677 (0.0%)	0.76	20/18487 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1
1	D	0	1
All	All	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	-1	MET	C-N	8.99	1.54	1.34
1	D	-1	MET	C-N	7.96	1.52	1.34

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	-1	MET	O-C-N	-13.44	101.20	122.70
1	C	237	ARG	NE-CZ-NH1	9.77	125.18	120.30
1	D	237	ARG	NE-CZ-NH1	8.59	124.60	120.30
1	C	33	ARG	NE-CZ-NH1	7.33	123.97	120.30
1	C	232	ARG	NE-CZ-NH2	-7.30	116.65	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	-1	MET	Mainchain
1	D	-1	MET	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3324	0	3286	64	0
1	B	3336	0	3321	66	0
1	C	3378	0	3358	45	0
1	D	3381	0	3359	54	0
2	A	44	0	26	0	0
2	B	44	0	26	1	0
2	C	44	0	26	0	0
2	D	44	0	26	0	0
3	A	10	0	4	0	0
3	B	10	0	4	1	0
3	C	10	0	4	0	0
3	D	10	0	4	0	0
4	C	6	0	8	0	0
5	D	13	0	18	7	0
6	A	66	0	0	2	0
6	B	70	0	0	1	0
6	C	308	0	0	2	0
6	D	315	0	0	6	0
All	All	14413	0	13470	225	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 225 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:281:ILE:HG22	1:D:286:PHE:HE1	1.06	1.10

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:299:GLY:O	1:D:343:LEU:CD1	2.05	1.05
1:D:281:ILE:HG22	1:D:286:PHE:CE1	1.97	0.99
1:C:299:GLY:O	1:C:343:LEU:CD2	2.14	0.96
1:D:327:TYR:HB3	1:D:329:MET:HE1	1.44	0.96

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	424/436 (97%)	407 (96%)	17 (4%)	0	100	100
1	B	424/436 (97%)	406 (96%)	17 (4%)	1 (0%)	47	38
1	C	432/436 (99%)	424 (98%)	7 (2%)	1 (0%)	47	38
1	D	433/436 (99%)	423 (98%)	8 (2%)	2 (0%)	29	18
All	All	1713/1744 (98%)	1660 (97%)	49 (3%)	4 (0%)	47	38

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	121	ASP
1	C	121	ASP
1	D	120	GLY
1	B	343	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	350/359 (98%)	343 (98%)	7 (2%)	55	51
1	B	351/359 (98%)	342 (97%)	9 (3%)	46	39
1	C	357/359 (99%)	350 (98%)	7 (2%)	55	51
1	D	357/359 (99%)	351 (98%)	6 (2%)	60	57
All	All	1415/1436 (98%)	1386 (98%)	29 (2%)	55	51

5 of 29 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	382	ARG
1	D	364	GLN
1	C	142[A]	CYS
1	D	156	THR
1	C	121	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	113	GLN
1	C	190	ASN
1	D	364	GLN
1	D	190	ASN
1	B	277	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAD	D	3	-	42,48,48	1.68	5 (11%)	50,73,73	1.64	10 (20%)
5	PG4	D	1	-	12,12,12	0.66	0	11,11,11	0.57	0
3	ADN	C	438	-	9,11,21	1.06	0	7,15,31	2.15	3 (42%)
2	NAD	A	438	-	42,48,48	1.68	3 (7%)	50,73,73	1.26	6 (12%)
2	NAD	C	2	-	42,48,48	1.61	4 (9%)	50,73,73	1.47	5 (10%)
3	ADN	B	438	-	9,11,21	0.82	0	7,15,31	2.30	3 (42%)
3	ADN	D	438	-	9,11,21	0.83	0	7,15,31	1.95	3 (42%)
2	NAD	B	1	-	42,48,48	1.64	3 (7%)	50,73,73	1.29	5 (10%)
3	ADN	A	439	-	9,11,21	1.05	0	7,15,31	2.26	2 (28%)
4	GOL	C	1	-	5,5,5	0.20	0	5,5,5	0.76	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	D	3	-	-	5/26/62/62	0/5/5/5
5	PG4	D	1	-	-	8/10/10/10	-
3	ADN	C	438	-	-	-	0/2/2/3
2	NAD	A	438	-	-	5/26/62/62	0/5/5/5
2	NAD	C	2	-	-	5/26/62/62	0/5/5/5
3	ADN	B	438	-	-	-	0/2/2/3
3	ADN	D	438	-	-	-	0/2/2/3
2	NAD	B	1	-	-	5/26/62/62	0/5/5/5
3	ADN	A	439	-	-	-	0/2/2/3
4	GOL	C	1	-	-	1/4/4/4	-

The worst 5 of 15 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	438	NAD	O7N-C7N	8.28	1.40	1.24
2	B	1	NAD	O7N-C7N	8.03	1.39	1.24
2	D	3	NAD	O7N-C7N	6.55	1.36	1.24
2	C	2	NAD	O7N-C7N	6.05	1.35	1.24
2	C	2	NAD	C2A-N1A	4.24	1.41	1.33

The worst 5 of 37 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	NAD	N3A-C2A-N1A	-5.56	119.98	128.68
2	C	2	NAD	N3A-C2A-N1A	-5.53	120.03	128.68
2	D	3	NAD	N3A-C2A-N1A	-5.10	120.70	128.68
2	A	438	NAD	N3A-C2A-N1A	-4.84	121.11	128.68
2	D	3	NAD	C3N-C7N-N7N	4.64	123.32	117.75

There are no chirality outliers.

5 of 29 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	438	NAD	O4D-C1D-N1N-C2N
2	A	438	NAD	O4D-C1D-N1N-C6N
2	A	438	NAD	C2D-C1D-N1N-C2N
2	A	438	NAD	C2D-C1D-N1N-C6N
2	B	1	NAD	O4D-C1D-N1N-C2N


There are no ring outliers.


3 monomers are involved in 8 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	1	PG4	7	0
3	B	438	ADN	1	0
2	B	1	NAD	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.


The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




















5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	430/436 (98%)	1.02	67 (15%) 2 2	14, 25, 34, 38	0
1	B	430/436 (98%)	1.11	87 (20%) 1 1	12, 26, 37, 41	0
1	C	435/436 (99%)	0.12	8 (1%) 68 71	2, 5, 15, 27	0
1	D	434/436 (99%)	0.12	7 (1%) 72 74	2, 6, 14, 22	0
All	All	1729/1744 (99%)	0.59	169 (9%) 7 8	2, 17, 33, 41	0

The worst 5 of 169 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	150	TYR	7.1
1	B	286	PHE	6.9
1	B	109	TRP	6.5
1	B	146	ASP	6.3
1	A	286	PHE	6.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	GOL	C	1	6/6	0.82	0.25	36,37,37,37	0
5	PG4	D	1	13/13	0.83	0.25	26,32,34,35	0
2	NAD	B	1	44/44	0.92	0.13	15,25,27,27	0
3	ADN	A	439	10/19	0.92	0.14	28,29,29,30	0
2	NAD	A	438	44/44	0.94	0.12	16,21,25,25	0
3	ADN	B	438	10/19	0.94	0.19	19,20,22,22	0
3	ADN	C	438	10/19	0.97	0.08	2,3,5,5	0
3	ADN	D	438	10/19	0.98	0.06	2,2,3,3	0
2	NAD	D	3	44/44	0.98	0.07	2,2,3,4	0
2	NAD	C	2	44/44	0.98	0.07	2,2,3,5	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around ADN A 439:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)











Electron density around ADN C 438:




$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)








Electron density around NAD D 3:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NAD C 2:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.