wwPDB X-ray Structure Validation Summary Report (i)

May 21, 2020 – 09:54 pm BST

PDB ID	:	6H9O
Title	:	Complex of the periplasmic domains of bacterial cell division proteins FtsQ and FtsB
Authors	:	Kureisaite-Ciziene, D.; Lowe, J.
Deposited on	:	2018-08-05
Resolution	:	2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	228	55% 29% • 14%
1	C	228	4% 51% 30% 5% 14%
2	B	24	67% 33%
2	D	24	71% 25% •

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 3574 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cell division protein FtsQ.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	196	1587	1000	289	292	6	0	0	0
1	C	195	1583	998	288	291	6	0	0	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	57	MET	-	initiating methionine	UNP P06136
A	277	GLY	-	expression tag	UNP P06136
A	278	SER	-	expression tag	UNP P06136
A	279	HIS	-	expression tag	UNP P06136
A	280	HIS	-	expression tag	UNP P06136
A	281	HIS	-	expression tag	UNP P06136
A	282	HIS	-	expression tag	UNP P06136
A	283	HIS	-	expression tag	UNP P06136
A	284	HIS	-	expression tag	UNP P06136
C	57	MET	-	initiating methionine	UNP P06136
C	277	GLY	-	expression tag	UNP P06136
C	278	SER	-	expression tag	UNP P06136
C	279	HIS	-	expression tag	UNP P06136
C	280	HIS	-	expression tag	UNP P06136
C	281	HIS	-	expression tag	UNP P06136
C	282	HIS	-	expression tag	UNP P06136
C	283	HIS	-	expression tag	UNP P06136
C	284	HIS	-	expression tag	UNP P06136

- Molecule 2 is a protein called Cell division protein FtsB.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	24	202	124	38	40	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	D	24	202	124	38	40	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	77	NLE	MET	conflict	UNP B7MKM2
D	77	NLE	MET	conflict	UNP B7MKM2

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cell division protein FtsQ

Chain A:



- Molecule 1: Cell division protein FtsQ

Chain C:



- Molecule 2: Cell division protein FtsB

Chain B:

- Molecule 2: Cell division protein FtsB

Chain D:


Q64
E65
E68
R72
R79
P80
G81
E82
Y85
R86
L87

4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α, β, γ	117.22Å 48.01Å 131.62Å 90.00° 112.40° 90.00°	Depositor
Resolution (Å)	29.83 – 2.80 43.90 – 2.80	Depositor EDS
% Data completeness (in resolution range)	92.4 (29.83-2.80) 98.3 (43.90-2.80)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.00 (at 2.81Å)	Xtriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, R_{free}	0.230 , 0.277 0.233 , 0.282	Depositor DCC
R_{free} test set	825 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å^2)	70.2	Xtriage
Anisotropy	0.379	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.33 , 55.2	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.93	EDS
Total number of atoms	3574	wwPDB-VP
Average B, all atoms (Å^2)	83.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.15% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NLE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.55	0/1618	0.80	2/2190 (0.1%)
1	C	0.55	2/1614 (0.1%)	1.02	9/2185 (0.4%)
2	B	0.57	0/195	0.73	0/259
2	D	0.50	0/195	0.80	1/259 (0.4%)
All	All	0.55	2/3622 (0.1%)	0.90	12/4893 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	114	GLN	C-N	5.95	1.47	1.34
1	C	189	LYS	CE-NZ	5.24	1.62	1.49

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	C	175	ARG	NE-CZ-NH1	13.56	127.08	120.30
1	C	189	LYS	CD-CE-NZ	-11.39	85.50	111.70
1	C	226	LEU	CB-CG-CD2	-11.38	91.66	111.00
1	C	175	ARG	NE-CZ-NH2	-7.73	116.43	120.30
1	C	70	ARG	CG-CD-NE	7.59	127.75	111.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1587	0	1594	58	0
1	C	1583	0	1591	94	2
2	B	202	0	195	5	0
2	D	202	0	195	7	0
All	All	3574	0	3575	160	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 22.

The worst 5 of 160 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:70:ARG:CZ	1:C:151:ARG:HH11	1.48	1.27
1:C:70:ARG:CZ	1:C:151:ARG:NH1	2.01	1.23
1:C:205:ASN:CG	1:C:207:ILE:HD13	1.63	1.19
1:C:70:ARG:NH2	1:C:151:ARG:HH11	1.50	1.09
1:A:92:VAL:HG13	1:A:111:VAL:HG23	1.34	1.05

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:115:TRP:CZ3	1:C:179:GLN:N[4_555]	1.80	0.40
1:C:115:TRP:CH2	1:C:178:GLY:CA[4_555]	2.10	0.10

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	192/228 (84%)	184 (96%)	7 (4%)	1 (0%)	29 61

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	191/228 (84%)	179 (94%)	11 (6%)	1 (0%)	29	61
2	B	21/24 (88%)	21 (100%)	0	0	100	100
2	D	21/24 (88%)	21 (100%)	0	0	100	100
All	All	425/504 (84%)	405 (95%)	18 (4%)	2 (0%)	29	61

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	116	PRO
1	C	116	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	171/199 (86%)	165 (96%)	6 (4%)	36	70
1	C	171/199 (86%)	162 (95%)	9 (5%)	22	54
2	B	20/20 (100%)	19 (95%)	1 (5%)	24	56
2	D	20/20 (100%)	20 (100%)	0	100	100
All	All	382/438 (87%)	366 (96%)	16 (4%)	30	63

5 of 16 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	66	ARG
1	C	67	HIS
1	C	166	SER
2	B	73	ASN
1	C	184	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	231	GLN
2	B	64	GLN
1	C	76	GLN
1	C	133	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	NLE	D	77	2	6,7,8	0.48	0	2,7,9	0.38	0
2	NLE	B	77	2	6,7,8	0.47	0	2,7,9	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NLE	D	77	2	-	2/5/6/8	-
2	NLE	B	77	2	-	2/5/6/8	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	77	NLE	CA-CB-CG-CD
2	D	77	NLE	CA-CB-CG-CD
2	D	77	NLE	CE-CD-CG-CB
2	B	77	NLE	CE-CD-CG-CB

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	196/228 (85%)	0.33	1 (0%)	91	88	42, 62, 111, 132	0
1	C	195/228 (85%)	0.46	8 (4%)	37	27	71, 101, 128, 137	0
2	B	23/24 (95%)	0.15	0	100	100	53, 64, 78, 93	0
2	D	23/24 (95%)	0.26	0	100	100	69, 79, 98, 111	0
All	All	437/504 (86%)	0.38	9 (2%)	63	54	42, 82, 123, 137	0

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	260	PRO	4.1
1	C	155	GLN	4.0
1	C	80	ALA	3.7
1	C	240	ARG	2.9
1	C	79	LEU	2.8

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	NLE	D	77	8/9	0.97	0.28	65,68,95,100	0
2	NLE	B	77	8/9	0.99	0.30	56,62,78,103	0

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.