



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 05:50 AM UTC

PDB ID : 7HS7 / pdb_00007hs7
Title : PanDDA analysis group deposition – Crystal Structure of FatA in complex with Z1230795916
Authors : Kot, E.; Ni, X.; Tomlinson, C.W.E.; Fearon, D.; Aschenbrenner, J.C.; Fairhead, M.; Koekemoer, L.; Marx, M.L.; Wright, N.D.; Mulholland, N.P.; Montgomery, M.G.; von Delft, F.
Deposited on : 2024-12-23
Resolution : 2.19 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.19 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	295	<div> <div>13%</div> <div>81%</div> <div>9%</div> <div>10%</div> </div>
1	B	295	<div> <div>38%</div> <div>76%</div> <div>13%</div> <div>11%</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4620 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Oleoyl-acyl carrier protein thioesterase 1, chloroplastic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	265	Total	C	N	O	S	0	3	0
			2176	1353	386	425	12			
1	B	264	Total	C	N	O	S	0	4	0
			2206	1375	392	429	10			

There are 14 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	74	MET	-	initiating methionine	UNP Q42561
A	363	HIS	-	expression tag	UNP Q42561
A	364	HIS	-	expression tag	UNP Q42561
A	365	HIS	-	expression tag	UNP Q42561
A	366	HIS	-	expression tag	UNP Q42561
A	367	HIS	-	expression tag	UNP Q42561
A	368	HIS	-	expression tag	UNP Q42561
B	74	MET	-	initiating methionine	UNP Q42561
B	363	HIS	-	expression tag	UNP Q42561
B	364	HIS	-	expression tag	UNP Q42561
B	365	HIS	-	expression tag	UNP Q42561
B	366	HIS	-	expression tag	UNP Q42561
B	367	HIS	-	expression tag	UNP Q42561
B	368	HIS	-	expression tag	UNP Q42561

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is 1-methyl-N-(2-methylpropyl)-1H-imidazole-2-carboxamide (CCD ID: A1BMO) (formula: C₉H₁₅N₃O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			13	9	3	1		
3	B	1	Total	C	N	O	0	0
			13	9	3	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	109	Total	O	0	0
			109	109		
4	B	83	Total	O	0	0
			83	83		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Oleoyl-acyl carrier protein thioesterase 1, chloroplastic



- Molecule 1: Oleoyl-acyl carrier protein thioesterase 1, chloroplastic



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	99.16Å 99.28Å 129.26Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.52 – 2.19 39.52 – 2.19	Depositor EDS
% Data completeness (in resolution range)	99.3 (39.52-2.19) 90.4 (39.52-2.19)	Depositor EDS
R_{merge}	0.40	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.09 (at 2.20Å)	Xtriage
Refinement program	BUSTER 2.10.4 (23-JAN-2024)	Depositor
R, R_{free}	0.264 , 0.308 (Not available) , 0.269	Depositor DCC
R_{free} test set	1599 reflections (4.82%)	wwPDB-VP
Wilson B-factor (Å ²)	39.4	Xtriage
Anisotropy	0.229	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 67.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	0.467 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4620	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, A1BMO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.67	0/2214	0.97	0/2997
1	B	0.69	1/2244 (0.0%)	0.98	1/3036 (0.0%)
All	All	0.68	1/4458 (0.0%)	0.98	1/6033 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	198	MET	SD-CE	-6.86	1.62	1.79

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	269	ASN	CA-CB-CG	6.69	119.29	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2176	0	2135	19	0
1	B	2206	0	2170	27	0
2	A	10	0	0	0	0
2	B	10	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	13	0	0	0	0
3	B	13	0	0	1	0
4	A	109	0	0	1	0
4	B	83	0	0	0	0
All	All	4620	0	4305	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:268:ASN:HD22	1:B:270:VAL:H	1.28	0.82
1:A:140:ILE:HG23	1:A:201:GLN:HG2	1.75	0.68
1:B:140:ILE:HG23	1:B:201:GLN:HG2	1.75	0.66
1:A:139:LEU:HD13	1:A:198:MET:HE3	1.78	0.65
1:A:280:ILE:HD11	1:A:338:HIS:HE1	1.63	0.62
1:B:280:ILE:HD11	1:B:338:HIS:HE1	1.63	0.62
1:B:281:PRO:HB2	1:B:283:GLU:HG2	1.85	0.59
1:B:146:MET:HE3	3:B:401:A1BMO:C2	2.34	0.57
1:A:95:VAL:HG11	1:A:156:TRP:HB2	1.87	0.57
1:B:234:ASN:ND2	1:B:236:SER:OG	2.37	0.56
1:B:171:ARG:NE	1:B:208:LYS:HZ1	2.05	0.55
1:B:95:VAL:HG11	1:B:156:TRP:HB2	1.88	0.54
1:A:98:ASN:HD21	1:A:359:LYS:NZ	2.06	0.54
1:A:171:ARG:HB3	1:A:208:LYS:HB3	1.90	0.53
1:B:198:MET:HE1	1:B:213:VAL:HG11	1.91	0.53
1:A:111:GLN:HE22	1:A:268:ASN:HD21	1.55	0.53
1:B:110:LEU:HD21	1:B:180[C]:LEU:HD11	1.91	0.52
1:B:110:LEU:HD21	1:B:180[A]:LEU:HD11	1.91	0.52
1:B:149[B]:GLU:HG2	1:B:292:VAL:HG13	1.92	0.51
1:B:111:GLN:HE22	1:B:268:ASN:HD21	1.57	0.51
1:A:138:HIS:HD2	1:A:202:ASP:OD2	1.95	0.49
1:B:138:HIS:HD2	1:B:202:ASP:OD2	1.95	0.49
1:A:201:GLN:NE2	4:A:505:HOH:O	2.47	0.48
1:A:98:ASN:HD21	1:A:359:LYS:HZ1	1.63	0.47
1:A:140:ILE:CG2	1:A:201:GLN:HG2	2.44	0.47
1:A:201:GLN:H	1:A:201:GLN:HG3	1.60	0.46
1:A:280:ILE:HD11	1:A:338:HIS:CE1	2.47	0.45
1:B:140:ILE:CG2	1:B:201:GLN:HG2	2.44	0.45
1:A:332:ASN:OD1	1:A:358:LYS:HD3	2.16	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:340:LEU:HB2	1:B:350:ASN:HB2	1.98	0.45
1:A:340:LEU:HB2	1:A:350:ASN:HB2	1.99	0.45
1:B:234:ASN:HD22	1:B:236:SER:H	1.65	0.45
1:B:89:VAL:HG22	1:B:159:VAL:HG22	1.97	0.45
1:A:89:VAL:HG22	1:A:159:VAL:HG22	1.98	0.45
1:B:280:ILE:HD11	1:B:338:HIS:CE1	2.47	0.44
1:B:170:GLY:O	1:B:214:ARG:NH2	2.50	0.44
1:B:145:ARG:HD3	1:B:296:ASP:OD1	2.19	0.43
1:B:211:ASP:O	1:B:214:ARG:HB2	2.19	0.43
1:B:214:ARG:NH1	1:B:217:TYR:OH	2.51	0.43
1:B:110:LEU:HD21	1:B:180[B]:LEU:HD11	1.99	0.42
1:B:201:GLN:H	1:B:201:GLN:HG3	1.60	0.42
1:A:166[B]:CYS:SG	1:A:176:ARG:CZ	3.08	0.42
1:A:166[B]:CYS:SG	1:A:176:ARG:NH1	2.93	0.41
1:B:166:CYS:SG	1:B:176:ARG:NH1	2.93	0.41
1:B:138:HIS:O	1:B:201:GLN:HG3	2.21	0.41
1:A:138:HIS:O	1:A:201:GLN:HG3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	265/295 (90%)	261 (98%)	4 (2%)	0	100	100
1	B	266/295 (90%)	260 (98%)	5 (2%)	1 (0%)	30	34
All	All	531/590 (90%)	521 (98%)	9 (2%)	1 (0%)	43	51

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	233	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	28/263 (11%)	26 (93%)	2 (7%)	13	16
1	B	93/263 (35%)	89 (96%)	4 (4%)	26	35
All	All	121/526 (23%)	115 (95%)	6 (5%)	21	28

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	302	GLN
1	A	357	ARG
1	B	97	SER
1	B	132	THR
1	B	142	VAL
1	B	145	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	302	GLN
1	A	303	GLN
1	B	138	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	402	-	4,4,4	0.30	0	6,6,6	0.28	0
3	A1BMO	B	401	-	13,13,13	0.69	1 (7%)	14,17,17	0.41	0
2	SO4	A	401	-	4,4,4	0.29	0	6,6,6	0.43	0
2	SO4	B	403	-	4,4,4	0.30	0	6,6,6	0.29	0
3	A1BMO	A	403	-	13,13,13	0.54	0	14,17,17	1.05	1 (7%)
2	SO4	B	402	-	4,4,4	0.28	0	6,6,6	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	A1BMO	A	403	-	-	2/9/9/9	0/1/1/1
3	A1BMO	B	401	-	-	2/9/9/9	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	401	A1BMO	C5-C6	-2.27	1.45	1.49

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	403	A1BMO	C4-N1-C5	3.24	126.94	122.06

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	403	A1BMO	C6-C5-N1-C4
3	A	403	A1BMO	O1-C5-N1-C4
3	B	401	A1BMO	C6-C5-N1-C4
3	B	401	A1BMO	O1-C5-N1-C4

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	401	A1BMO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	265/295 (89%)	1.25	39 (14%) 6 4	7, 46, 72, 83	10 (3%)
1	B	264/295 (89%)	1.88	112 (42%) 0 0	4, 40, 61, 82	88 (33%)
All	All	529/590 (89%)	1.56	151 (28%) 1 1	4, 44, 68, 83	98 (18%)

All (151) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	217	TYR	7.9
1	A	146	MET	7.3
1	B	219	VAL	7.1
1	B	209	VAL	6.6
1	A	129	ALA	6.5
1	B	214	ARG	6.0
1	B	155	ALA	5.9
1	A	166[A]	CYS	5.8
1	B	132	THR	5.7
1	A	123	PHE	5.6
1	A	345	ASP	5.6
1	A	130	THR	5.6
1	B	218	LEU	5.5
1	B	210	SER	5.3
1	A	120[A]	SER	5.1
1	B	204	ARG	5.1
1	A	133	THR	5.1
1	A	134	MET	5.0
1	B	135	ARG	5.0
1	B	153	TYR	4.9
1	B	213	VAL	4.9
1	A	131	THR	4.8
1	B	133	THR	4.8
1	A	124	SER	4.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	166	CYS	4.7
1	B	129	ALA	4.7
1	A	217	TYR	4.6
1	B	211	ASP	4.6
1	B	154	PRO	4.6
1	B	172	ILE	4.6
1	B	189	THR	4.4
1	B	184	VAL	4.4
1	B	357	ARG	4.4
1	B	151	TYR	4.3
1	B	156	TRP	4.2
1	B	196	TRP	4.2
1	B	130	THR	4.2
1	B	168	SER	4.1
1	A	115	CYS	4.1
1	B	144	ALA	4.1
1	B	202	ASP	4.1
1	B	185	THR	4.1
1	B	223	GLN	4.0
1	B	215	ASP	4.0
1	A	236	SER	4.0
1	B	183	SER	4.0
1	B	208	LYS	4.0
1	A	97[A]	SER	3.9
1	B	268	ASN	3.9
1	B	201	GLN	3.9
1	B	216	GLU	3.9
1	B	191[A]	ARG	3.8
1	A	332	ASN	3.7
1	B	123	PHE	3.7
1	B	181	LYS	3.7
1	A	219	VAL	3.7
1	B	336	PHE	3.7
1	B	182	ASP	3.7
1	B	164	THR	3.7
1	B	232	GLU	3.6
1	B	158	ASP	3.6
1	B	131	THR	3.6
1	A	215	ASP	3.6
1	A	205	ARG	3.5
1	B	212	ASP	3.5
1	B	206	LEU	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	237	LEU	3.4
1	A	132	THR	3.4
1	B	77	LEU	3.3
1	B	75	GLY	3.3
1	B	333	ASP	3.2
1	B	173	GLY	3.2
1	B	180[A]	LEU	3.1
1	A	235	ARG	3.1
1	B	167	GLN	3.1
1	B	79	GLU	3.1
1	B	205	ARG	3.0
1	B	124	SER	3.0
1	B	207	GLN	3.0
1	B	300[A]	GLU	2.9
1	B	160	VAL	2.9
1	B	179	ILE	2.9
1	B	142	VAL	2.9
1	B	344	GLY	2.9
1	B	139	LEU	2.8
1	B	334	SER	2.8
1	B	220	PHE	2.8
1	B	171	ARG	2.8
1	B	170	GLY	2.8
1	B	76	SER	2.8
1	B	200	ASN	2.8
1	B	194	SER	2.8
1	A	119	GLN	2.7
1	B	264	ASN	2.7
1	A	213	VAL	2.7
1	B	140	ILE	2.7
1	B	291	GLN	2.7
1	B	198	MET	2.7
1	B	161	GLU	2.7
1	B	146	MET	2.7
1	B	134	MET	2.6
1	B	195	LYS	2.6
1	B	149[A]	GLU	2.6
1	A	220	PHE	2.6
1	B	235	ARG	2.6
1	B	122	GLY	2.6
1	A	210	SER	2.6
1	B	176	ARG	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	121	VAL	2.6
1	B	193	THR	2.6
1	A	207	GLN	2.5
1	B	283	GLU	2.5
1	B	84	TYR	2.5
1	B	174	THR	2.5
1	B	356	TRP	2.5
1	B	152	LYS	2.5
1	A	233	ASN	2.5
1	B	249	TYR	2.5
1	A	172	ILE	2.5
1	B	252	ILE	2.5
1	B	165	TRP	2.4
1	B	187	GLU	2.4
1	B	225	PRO	2.4
1	B	305	ASP	2.3
1	A	247	ALA	2.3
1	B	98	ASN	2.3
1	A	218	LEU	2.3
1	B	346	GLY	2.3
1	B	169	GLU	2.3
1	A	204	ARG	2.3
1	B	310	LEU	2.3
1	B	351	ARG	2.2
1	A	206	LEU	2.2
1	B	105	THR	2.2
1	A	83	SER	2.2
1	B	119	GLN	2.2
1	A	214	ARG	2.2
1	A	234	ASN	2.2
1	B	188	VAL	2.1
1	B	186	GLY	2.1
1	B	78	THR	2.1
1	B	141	TRP	2.1
1	A	336	PHE	2.1
1	B	116	ASN	2.1
1	B	315	SER	2.1
1	B	355	LEU	2.1
1	A	173	GLY	2.1
1	B	137	LEU	2.0
1	A	135	ARG	2.0
1	B	313	THR	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	297	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	A1BMO	A	403	13/13	0.82	0.22	30,31,35,35	13
3	A1BMO	B	401	13/13	0.86	0.12	31,32,34,35	13
2	SO4	A	401	5/5	0.87	0.18	96,96,96,97	0
2	SO4	B	402	5/5	0.87	0.13	50,51,51,51	5
2	SO4	B	403	5/5	0.94	0.12	44,44,44,44	5
2	SO4	A	402	5/5	0.95	0.08	69,69,70,70	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around A1BMO A 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around A1BMO B 401:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.