



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 15, 2024 – 05:33 PM EDT

PDB ID : 2IFU  
Title : Crystal Structure of a Gamma-SNAP from Danio rerio  
Authors : Bitto, E.; Wesenberg, G.E.; Phillips Jr., G.N.; McCoy, J.G.; Bingman, C.A.;  
Center for Eukaryotic Structural Genomics (CESG)  
Deposited on : 2006-09-21  
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	307	80% 8% • 10%
1	B	307	78% 10% • 10%
1	C	307	79% 8% • 11%
1	D	307	79% 8% • 12%

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 8758 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called gamma-snap.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	275	Total	C	N	O	S	Se	0	2	0
			2181	1373	372	419	5	12			
1	B	275	Total	C	N	O	S	Se	0	1	0
			2176	1369	372	419	5	11			
1	C	272	Total	C	N	O	S	Se	0	0	0
			2152	1354	368	415	5	10			
1	D	270	Total	C	N	O	S	Se	0	0	0
			2135	1342	365	413	5	10			

There are 52 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	ALA	-	CLONING ARTIFACT	UNP Q5BJK3
A	0	ILE	-	CLONING ARTIFACT	UNP Q5BJK3
A	1	ALA	MET	CLONING ARTIFACT	UNP Q5BJK3
A	24	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
A	82	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
A	83	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
A	90	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
A	103	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
A	115	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
A	124	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
A	182	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
A	186	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
A	261	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
B	-1	ALA	-	CLONING ARTIFACT	UNP Q5BJK3
B	0	ILE	-	CLONING ARTIFACT	UNP Q5BJK3
B	1	ALA	MET	CLONING ARTIFACT	UNP Q5BJK3
B	24	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
B	82	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
B	83	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
B	90	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
B	103	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	115	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
B	124	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
B	182	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
B	186	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
B	261	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
C	-1	ALA	-	CLONING ARTIFACT	UNP Q5BJK3
C	0	ILE	-	CLONING ARTIFACT	UNP Q5BJK3
C	1	ALA	MET	CLONING ARTIFACT	UNP Q5BJK3
C	24	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
C	82	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
C	83	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
C	90	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
C	103	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
C	115	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
C	124	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
C	182	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
C	186	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
C	261	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
D	-1	ALA	-	CLONING ARTIFACT	UNP Q5BJK3
D	0	ILE	-	CLONING ARTIFACT	UNP Q5BJK3
D	1	ALA	MET	CLONING ARTIFACT	UNP Q5BJK3
D	24	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
D	82	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
D	83	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
D	90	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
D	103	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
D	115	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
D	124	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
D	182	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
D	186	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3
D	261	MSE	MET	MODIFIED RESIDUE	UNP Q5BJK3

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	41	Total	O	0	0
			41	41		
3	B	17	Total	O	0	0
			17	17		

*Continued on next page...*


*Continued from previous page...*


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	6	Total	O	0	0
			6	6		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: gamma-snap

Chain A: 




- Molecule 1: gamma-snap

Chain B: 





- Molecule 1: gamma-snap

Chain C: 



- Molecule 1: gamma-snap

Chain D: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	83.38Å 90.81Å 264.17Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.05 – 2.60 40.05 – 2.60	Depositor EDS
% Data completeness (in resolution range)	96.1 (40.05-2.60) 96.1 (40.05-2.60)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.41 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.239 , 0.261 0.363 , 0.368	Depositor DCC
$R_{free}$ test set	3047 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	61.1	Xtriage
Anisotropy	0.396	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 28.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.84	EDS
Total number of atoms	8758	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.31% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.00	6/2213 (0.3%)	0.86	3/2959 (0.1%)
1	B	0.90	6/2206 (0.3%)	0.92	4/2952 (0.1%)
1	C	0.57	1/2179 (0.0%)	0.64	2/2916 (0.1%)
1	D	0.42	0/2162	0.55	0/2894
All	All	0.76	13/8760 (0.1%)	0.76	9/11721 (0.1%)

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	241	TYR	CE2-CZ	-8.19	1.27	1.38
1	A	215	CYS	CB-SG	-7.34	1.69	1.82
1	A	103	MSE	CG-SE	-6.44	1.73	1.95
1	A	60	GLU	CB-CG	-6.21	1.40	1.52
1	B	115[A]	MSE	SE-CE	5.98	2.30	1.95
1	B	115[B]	MSE	SE-CE	5.98	2.30	1.95
1	B	241	TYR	CG-CD2	-5.83	1.31	1.39
1	C	115	MSE	SE-CE	5.78	2.29	1.95
1	B	241	TYR	CE1-CZ	-5.73	1.31	1.38
1	A	10	GLU	CD-OE2	5.52	1.31	1.25
1	A	241	TYR	CE1-CZ	-5.32	1.31	1.38
1	A	241	TYR	CG-CD2	-5.31	1.32	1.39
1	B	241	TYR	CG-CD1	-5.30	1.32	1.39

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	217	ARG	NE-CZ-NH2	15.91	128.25	120.30
1	B	217	ARG	NE-CZ-NH1	-13.78	113.41	120.30
1	A	217	ARG	NE-CZ-NH2	-8.48	116.06	120.30
1	A	217	ARG	NE-CZ-NH1	7.76	124.18	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	217	ARG	CD-NE-CZ	7.76	134.47	123.60
1	C	217	ARG	NE-CZ-NH2	-6.49	117.06	120.30
1	A	217	ARG	CD-NE-CZ	5.13	130.78	123.60
1	B	68	ARG	NE-CZ-NH2	-5.05	117.78	120.30
1	C	217	ARG	NE-CZ-NH1	5.04	122.82	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2181	0	2171	15	0
1	B	2176	0	2162	17	1
1	C	2152	0	2132	14	0
1	D	2135	0	2108	17	1
2	A	15	0	0	0	0
2	B	20	0	0	1	0
2	C	10	0	0	0	0
2	D	5	0	0	0	0
3	A	41	0	0	0	0
3	B	17	0	0	0	0
3	C	6	0	0	0	0
All	All	8758	0	8573	60	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:186:MSE:CE	1:A:186:MSE:SE	2.17	1.43
1:B:115[B]:MSE:SE	1:B:115[B]:MSE:CE	2.20	1.39
1:A:24:MSE:SE	1:A:24:MSE:CE	2.23	1.34
1:C:115:MSE:SE	1:C:115:MSE:CE	2.29	1.31

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:115[A]:MSE:CE	1:B:115[A]:MSE:SE	2.30	1.28
1:A:82[A]:MSE:SE	1:A:119:ARG:HH21	1.87	1.07
1:A:90:MSE:HE2	1:A:123:LEU:HB2	1.40	1.03
1:C:220:TYR:OH	1:C:235:GLU:OE2	1.90	0.88
1:A:85:LYS:HD2	1:A:123:LEU:HD13	1.58	0.84
1:B:244:GLN:HE21	1:B:275:PRO:HG2	1.46	0.80
1:A:82[A]:MSE:SE	1:A:119:ARG:NH2	2.69	0.71
1:B:260:TYR:OH	1:D:251:ARG:HA	1.89	0.70
1:A:248:GLN:HG2	1:A:251:ARG:NH2	2.11	0.65
1:A:244:GLN:HE21	1:A:275:PRO:HG2	1.62	0.64
1:C:248:GLN:HG2	1:C:251:ARG:HH22	1.61	0.64
1:A:248:GLN:HG2	1:A:251:ARG:HH22	1.63	0.64
1:D:248:GLN:HG2	1:D:251:ARG:NH2	2.13	0.63
1:C:248:GLN:HG2	1:C:251:ARG:NH2	2.14	0.63
1:A:90:MSE:HE2	1:A:123:LEU:CB	2.24	0.60
1:B:72:HIS:ND1	2:B:308:SO4:O3	2.33	0.60
1:D:248:GLN:HG2	1:D:251:ARG:HH22	1.66	0.60
1:A:25:LYS:HG3	1:A:25:LYS:O	2.03	0.58
1:D:197:ILE:HA	1:D:234:LEU:HD21	1.87	0.57
1:B:25:LYS:O	1:B:25:LYS:HG3	2.04	0.57
1:A:197:ILE:HA	1:A:234:LEU:HD21	1.87	0.56
1:B:197:ILE:HA	1:B:234:LEU:HD21	1.87	0.56
1:D:244:GLN:HE21	1:D:275:PRO:HG2	1.72	0.54
1:B:157:GLY:HA2	1:B:179:GLU:OE2	2.08	0.53
1:A:244:GLN:HA	1:A:274:VAL:HG23	1.92	0.51
1:B:248:GLN:HG2	1:B:251:ARG:NH2	2.28	0.49
1:C:70:LEU:HD13	1:C:106:GLU:HB3	1.95	0.48
1:B:244:GLN:HA	1:B:274:VAL:HG23	1.94	0.48
1:B:248:GLN:HG2	1:B:251:ARG:HH22	1.79	0.47
1:B:259:THR:HB	1:D:254:ARG:HG2	1.96	0.47
1:D:244:GLN:HA	1:D:274:VAL:HG23	1.96	0.46
1:D:93:ALA:O	1:D:97:ILE:HG12	2.15	0.46
1:C:157:GLY:HA2	1:C:179:GLU:OE2	2.17	0.44
1:D:24:MSE:HE3	1:D:24:MSE:HB2	1.91	0.44
1:C:244:GLN:HA	1:C:274:VAL:HG23	1.99	0.44
1:D:195:LYS:HA	1:D:195:LYS:HD3	1.89	0.44
1:D:263:ASN:HD22	1:D:263:ASN:N	2.15	0.43
1:C:143:PHE:CG	1:C:151:GLN:HB3	2.54	0.43
1:B:115[A]:MSE:HB3	1:B:115[A]:MSE:HE3	1.99	0.43
1:C:68:ARG:NH1	1:D:236:ASP:OD2	2.49	0.43
1:C:197:ILE:HA	1:C:234:LEU:HD21	2.01	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:22:SER:OG	1:D:23:PHE:N	2.51	0.43
1:C:258:VAL:O	1:C:261:MSE:HG2	2.18	0.42
1:A:85:LYS:HD2	1:A:123:LEU:CD1	2.40	0.42
1:C:99:LYS:O	1:C:103:MSE:HG3	2.19	0.42
1:B:220:TYR:OH	1:B:235:GLU:OE2	2.29	0.42
1:B:194:LYS:O	1:B:261:MSE:HE1	2.20	0.42
1:D:258:VAL:O	1:D:261:MSE:HG2	2.18	0.42
1:B:125:GLU:HB2	1:B:126:PRO:HD3	2.02	0.41
1:D:25:LYS:O	1:D:25:LYS:HG3	2.21	0.41
1:A:258:VAL:O	1:A:261:MSE:HG2	2.21	0.41
1:B:195:LYS:HA	1:B:195:LYS:HD3	1.86	0.41
1:D:125:GLU:HB2	1:D:126:PRO:HD3	2.03	0.40
1:C:246:GLU:HG3	1:C:274:VAL:HG12	2.03	0.40
1:C:22:SER:C	1:C:24:MSE:H	2.24	0.40
1:D:157:GLY:HA2	1:D:179:GLU:OE2	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:5:ILE:CG2	1:D:147:GLU:OE2[4_445]	2.17	0.03

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	275/307 (90%)	267 (97%)	8 (3%)	0	100	100
1	B	274/307 (89%)	267 (97%)	7 (3%)	0	100	100
1	C	270/307 (88%)	261 (97%)	9 (3%)	0	100	100
1	D	268/307 (87%)	262 (98%)	6 (2%)	0	100	100

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	1087/1228 (88%)	1057 (97%)	30 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	225/233 (97%)	216 (96%)	9 (4%)	31	57
1	B	224/233 (96%)	211 (94%)	13 (6%)	20	40
1	C	222/233 (95%)	216 (97%)	6 (3%)	44	71
1	D	220/233 (94%)	215 (98%)	5 (2%)	50	75
All	All	891/932 (96%)	858 (96%)	33 (4%)	34	60

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	SER
1	A	24	MSE
1	A	25	LYS
1	A	60	GLU
1	A	87	LEU
1	A	124	MSE
1	A	127	LEU
1	A	151	GLN
1	A	248	GLN
1	B	21	THR
1	B	22	SER
1	B	24	MSE
1	B	25	LYS
1	B	87	LEU
1	B	124	MSE
1	B	127	LEU
1	B	130	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	151	GLN
1	B	213	GLN
1	B	214	LYS
1	B	217	ARG
1	B	248	GLN
1	C	26	TRP
1	C	60	GLU
1	C	87	LEU
1	C	127	LEU
1	C	151	GLN
1	C	248	GLN
1	D	22	SER
1	D	60	GLU
1	D	87	LEU
1	D	127	LEU
1	D	248	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (28) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	49	GLN
1	A	52	GLN
1	A	138	GLN
1	A	213	GLN
1	A	244	GLN
1	A	263	ASN
1	B	49	GLN
1	B	52	GLN
1	B	138	GLN
1	B	151	GLN
1	B	213	GLN
1	B	244	GLN
1	B	263	ASN
1	C	49	GLN
1	C	52	GLN
1	C	138	GLN
1	C	151	GLN
1	C	166	GLN
1	C	213	GLN
1	C	244	GLN
1	C	263	ASN
1	D	49	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	52	GLN
1	D	107	ASN
1	D	138	GLN
1	D	213	GLN
1	D	244	GLN
1	D	263	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	A	308	-	4,4,4	0.27	0	6,6,6	0.38	0
2	SO4	C	306	-	4,4,4	0.31	0	6,6,6	0.34	0
2	SO4	B	306	-	4,4,4	0.27	0	6,6,6	0.25	0
2	SO4	A	306	-	4,4,4	0.31	0	6,6,6	0.89	0
2	SO4	B	308	-	4,4,4	0.44	0	6,6,6	0.95	0
2	SO4	D	306	-	4,4,4	0.22	0	6,6,6	0.17	0
2	SO4	A	307	-	4,4,4	0.32	0	6,6,6	0.76	0
2	SO4	B	307	-	4,4,4	0.31	0	6,6,6	0.82	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	C	307	-	4,4,4	0.22	0	6,6,6	0.20	0
2	SO4	B	309	-	4,4,4	0.30	0	6,6,6	0.36	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	308	SO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.