



## wwPDB EM Validation Summary Report ⓘ

Mar 7, 2026 – 11:43 PM UTC

PDB ID : 8IXN / pdb\_00008ixn  
EMDB ID : EMD-35799  
Title : Curved structure of mPIEZO1-S2472E  
Authors : Liu, S.; Yang, X.; Chen, X.; Li, X.; Xiao, B.  
Deposited on : 2023-04-01  
Resolution : 4.07 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132  
Mogul : 2022.3.0, CSD as543be (2022)  
MolProbity : 4-5-2 with Phenix2.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49



# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 4.07 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	6427 ( 3.58 - 4.57 )

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	2547	<div> <div>19%</div> <div>38%</div> <div>15%</div> <div>47%</div> </div>
1	C	2547	<div> <div>21%</div> <div>37%</div> <div>16%</div> <div>47%</div> </div>
1	D	2547	<div> <div>11%</div> <div>39%</div> <div>14%</div> <div>47%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PEE	A	2603	-	-	X	-
3	PEE	C	2603	-	-	X	-
3	PEE	D	2603	-	-	X	-
4	P5S	A	2604	-	-	X	-
4	P5S	C	2604	-	-	X	-

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 30043 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Piezo-type mechanosensitive ion channel component 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	C	1350	Total	C	N	O	S	0	0
			9804	6364	1661	1736	43		
1	A	1350	Total	C	N	O	S	0	0
			9802	6363	1660	1736	43		
1	D	1350	Total	C	N	O	S	0	0
			9810	6369	1661	1736	44		

There are 3 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	2472	GLU	SER	engineered mutation	UNP E2JF22
A	2472	GLU	SER	engineered mutation	UNP E2JF22
D	2472	GLU	SER	engineered mutation	UNP E2JF22

- Molecule 2 is (9R,11S)-9-({[(1S)-1-HYDROXYHEXADECYL]OXY}METHYL)-2,2-DIMETHYL-5,7,10-TRIOXA-2LAMBDA 5 -AZA-6LAMBDA 5 -PHOSPHAOCTACOSANE-6,6,11-TRIOX (CCD ID: PLX) (formula: C<sub>42</sub>H<sub>89</sub>NO<sub>8</sub>P) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					AltConf
2	C	1	Total	C	N	O	P	0
			52	42	1	8	1	
2	C	1	Total	C	N	O	P	0
			52	42	1	8	1	
2	A	1	Total	C	N	O	P	0
			52	42	1	8	1	
2	A	1	Total	C	N	O	P	0
			52	42	1	8	1	
2	D	1	Total	C	N	O	P	0
			52	42	1	8	1	
2	D	1	Total	C	N	O	P	0
			52	42	1	8	1	

- Molecule 3 is 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine (CCD ID: PEE) (formula:  $C_{41}H_{78}NO_8P$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					AltConf
3	C	1	Total	C	N	O	P	0
			51	41	1	8	1	
3	A	1	Total	C	N	O	P	0
			51	41	1	8	1	
3	D	1	Total	C	N	O	P	0
			51	41	1	8	1	

- Molecule 4 is O-[(R)-{[(2R)-2,3-bis(octadecanoyloxy)propyl]oxy}(hydroxy)phosphoryl]-L-serine (CCD ID: P5S) (formula:  $C_{42}H_{82}NO_{10}P$ ) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					AltConf
4	C	1	Total	C	N	O	P	0
			54	42	1	10	1	
4	A	1	Total	C	N	O	P	0
			54	42	1	10	1	
4	D	1	Total	C	N	O	P	0
			54	42	1	10	1	











- Molecule 1: Piezo-type mechanosensitive ion channel component 1








- Molecule 1: Piezo-type mechanosensitive ion channel component 1









## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	488709	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	50	Depositor
Minimum defocus (nm)	1500	Depositor
Maximum defocus (nm)	2300	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	22.183	Depositor
Minimum map value	-9.040	Depositor
Average map value	0.004	Depositor
Map value standard deviation	1.021	Depositor
Recommended contour level	8.23	Depositor
Map size ( $\text{\AA}$ )	404.02722, 404.02722, 404.02722	wwPDB
Map dimensions	368, 368, 368	wwPDB
Map angles ( $^\circ$ )	90.0, 90.0, 90.0	wwPDB
Pixel spacing ( $\text{\AA}$ )	1.0979, 1.0979, 1.0979	Depositor



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PLX, P5S, PEE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.20	0/9986	0.45	1/13618 (0.0%)
1	C	0.20	0/9990	0.46	1/13625 (0.0%)
1	D	0.19	0/9995	0.43	1/13630 (0.0%)
All	All	0.20	0/29971	0.45	3/40873 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	1233	VAL	N-CA-C	-6.31	105.19	111.88
1	A	1233	VAL	N-CA-C	-6.23	105.28	111.88
1	D	1233	VAL	N-CA-C	-6.17	105.34	111.88

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	9802	0	8926	380	0
1	C	9804	0	8928	414	0
1	D	9810	0	8942	346	0
2	A	104	0	176	14	0
2	C	104	0	176	14	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	104	0	176	14	0
3	A	51	0	82	60	0
3	C	51	0	82	72	0
3	D	51	0	82	64	0
4	A	54	0	80	24	0
4	C	54	0	80	25	0
4	D	54	0	80	17	0
All	All	30043	0	27810	1179	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 1179 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:2109:PHE:HB2	3:D:2603:PEE:C5	1.55	1.34
1:A:1981:PHE:CE2	3:A:2603:PEE:H17	1.66	1.30
1:D:1981:PHE:CE2	3:D:2603:PEE:H17	1.77	1.17
1:C:1981:PHE:CD2	3:C:2603:PEE:H17	1.78	1.17
1:A:1981:PHE:CD2	3:A:2603:PEE:H13	1.82	1.13

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	1312/2547 (52%)	1241 (95%)	69 (5%)	2 (0%)	43 75
1	C	1312/2547 (52%)	1245 (95%)	66 (5%)	1 (0%)	48 81
1	D	1312/2547 (52%)	1241 (95%)	69 (5%)	2 (0%)	43 75
All	All	3936/7641 (52%)	3727 (95%)	204 (5%)	5 (0%)	49 81

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	2182	LYS
1	A	2182	LYS
1	D	2182	LYS
1	A	2183	LYS
1	D	2183	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	869/2246 (39%)	868 (100%)	1 (0%)	88	89
1	C	870/2246 (39%)	870 (100%)	0	100	100
1	D	872/2246 (39%)	872 (100%)	0	100	100
All	All	2611/6738 (39%)	2610 (100%)	1 (0%)	100	100

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2391	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 30 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2115	ASN
1	D	2285	GLN
1	A	2324	GLN
1	D	2391	GLN
1	D	2058	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	PLX	A	2601	-	51,51,51	0.60	0	53,59,59	0.69	0
3	PEE	A	2603	-	50,50,50	1.20	7 (14%)	53,55,55	1.26	6 (11%)
2	PLX	D	2601	-	51,51,51	0.59	0	53,59,59	0.69	0
2	PLX	C	2601	-	51,51,51	0.59	0	53,59,59	0.69	0
4	P5S	A	2604	-	52,53,53	0.39	0	54,60,60	0.40	0
2	PLX	D	2602	-	51,51,51	0.64	0	53,59,59	0.71	1 (1%)
3	PEE	C	2603	-	50,50,50	1.20	7 (14%)	53,55,55	1.27	6 (11%)
3	PEE	D	2603	-	50,50,50	1.20	7 (14%)	53,55,55	1.27	7 (13%)
4	P5S	C	2604	-	52,53,53	0.38	0	54,60,60	0.40	0
4	P5S	D	2604	-	52,53,53	0.39	0	54,60,60	0.40	0
2	PLX	C	2602	-	51,51,51	0.63	0	53,59,59	0.71	1 (1%)
2	PLX	A	2602	-	51,51,51	0.64	0	53,59,59	0.71	1 (1%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLX	A	2601	-	-	3/55/55/55	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PEE	A	2603	-	-	27/54/54/54	-
2	PLX	D	2601	-	-	3/55/55/55	-
2	PLX	C	2601	-	-	3/55/55/55	-
4	P5S	A	2604	-	-	16/59/59/59	-
2	PLX	D	2602	-	-	9/55/55/55	-
3	PEE	C	2603	-	-	27/54/54/54	-
3	PEE	D	2603	-	-	27/54/54/54	-
4	P5S	C	2604	-	-	16/59/59/59	-
4	P5S	D	2604	-	-	16/59/59/59	-
2	PLX	C	2602	-	-	9/55/55/55	-
2	PLX	A	2602	-	-	9/55/55/55	-

The worst 5 of 21 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	2603	PEE	C39-C38	3.77	1.53	1.31
3	A	2603	PEE	C39-C38	3.76	1.53	1.31
3	C	2603	PEE	C39-C38	3.76	1.53	1.31
3	A	2603	PEE	C18-C19	3.55	1.51	1.31
3	C	2603	PEE	C18-C19	3.55	1.51	1.31

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	2603	PEE	O2P-P-O1P	3.46	128.56	112.44
3	D	2603	PEE	O2P-P-O1P	3.46	128.56	112.44
3	A	2603	PEE	O2P-P-O1P	3.46	128.55	112.44
3	C	2603	PEE	C2-O2-C10	2.35	123.41	117.80
3	A	2603	PEE	C2-O2-C10	2.35	123.41	117.80

There are no chirality outliers.

5 of 165 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	2602	PLX	O7-C6-O6-C4
2	C	2602	PLX	C2-O1-P1-O4
2	C	2602	PLX	C2-O1-P1-O2
2	A	2602	PLX	O7-C6-O6-C4
2	A	2602	PLX	C2-O1-P1-O4





There are no ring outliers.





12 monomers are involved in 304 short contacts:





Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	2601	PLX	1	0
3	A	2603	PEE	60	0
2	D	2601	PLX	2	0
2	C	2601	PLX	2	0
4	A	2604	P5S	24	0
2	D	2602	PLX	12	0
3	C	2603	PEE	72	0
3	D	2603	PEE	64	0
4	C	2604	P5S	25	0
4	D	2604	P5S	17	0
2	C	2602	PLX	12	0
2	A	2602	PLX	13	0





The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











Ligand PEE A 2603	
	
Bond lengths	Bond angles
	
Torsions	Rings





Ligand PLX D 2601	
	
Bond lengths	Bond angles
	
Torsions	Rings





Ligand PLX C 2601	
	
Bond lengths	Bond angles
	
Torsions	Rings

Ligand P5S A 2604	
	
Bond lengths	Bond angles
	
Torsions	Rings


Ligand PLX D 2602	
	
Bond lengths	Bond angles
	
Torsions	Rings

Ligand PEE C 2603	
	
Bond lengths	Bond angles
	
Torsions	Rings

Ligand PEE D 2603	
	
Bond lengths	Bond angles
	
Torsions	Rings

Ligand P5S C 2604	
	
Bond lengths	Bond angles
	
Torsions	Rings





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-35799. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X




Y




Z


#### 6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

## 6.2 Central slices [i](#)


### 6.2.1 Primary map



X Index: 184




Y Index: 184




Z Index: 184


### 6.2.2 Raw map



X Index: 184



Y Index: 184




Z Index: 184


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 202




Y Index: 179




Z Index: 185


### 6.3.2 Raw map



X Index: 111



Y Index: 189




Z Index: 236


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)

### 6.4.1 Primary map



### 6.4.2 Raw map



The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

## 6.5 Orthogonal surface views [i](#)

This section was not generated.


## 6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 170 nm<sup>3</sup>; this corresponds to an approximate mass of 154 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum ⓘ




\*Reported resolution corresponds to spatial frequency of 0.246  $\text{\AA}^{-1}$



## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of 0.246 Å<sup>-1</sup>

## 8.2 Resolution estimates [i](#)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	4.07	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	3.82	6.57	4.00

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.


## 9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-35799 and PDB model 8IXN. Per-residue inclusion information can be found in section 3 on page 7.

### 9.1 Map-model overlay [i](#)

This section was not generated.

### 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

### 9.3 Atom inclusion mapped to coordinate model [i](#)

This section was not generated.

## 9.4 Atom inclusion [i](#)



At the recommended contour level, 74% of all backbone atoms, 51% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (8.23) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div></div> 0.5120	<div></div> 0.3010
A	<div></div> 0.4980	<div></div> 0.2760
C	<div></div> 0.4770	<div></div> 0.2610
D	<div></div> 0.5600	<div></div> 0.3640

