wwPDB X-ray Structure Validation Summary Report (i)

Sep 10, 2023 – 06:00 PM EDT

PDB ID	:	4JKN
Title	:	Mercury Metallated Pseudomonas aeruginosa Azurin at 1.54 A
Authors	:	Zampino, A.P.; Masters, F.M.; Bladholm, E.L.; Berry, S.B.; Panzner, M.J.; Ziegler, C.J.
Deposited on	:	2013-03-10
Resolution	:	1.54 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.54 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2556 (1.56-1.52)
Clashscore	141614	2634 (1.56-1.52)
Ramachandran outliers	138981	2580 (1.56-1.52)
Sidechain outliers	138945	2577 (1.56-1.52)
RSRZ outliers	127900	2524 (1.56-1.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	128	5% 96% • •
1	B	128	7% 88% 12%
1	C	128	8% 90% 10%
1	D	128	5% 92% 7% •

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 8362 atoms, of which 3900 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Azurin.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	128	1994	627	985	170	203	9	0	4	0
1	B	128	1986	624	985	170	198	9	0	6	0
1	C	128	1951	613	968	166	195	9	0	2	0
1	D	128	1944	612	962	165	195	10	0	1	0

- Molecule 2 is MERCURY (II) ION (three-letter code: HG) (formula: Hg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Hg		
2	A	2	3	3	0	1
2	B	1	1	1	0	0
2	C	1	1	1	0	0
2	D	1	1	1	0	0

- Molecule 3 is NITRATE ION (three-letter code: NO3) (formula: NO_3).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	N	O		
3	A	1	4	1	3	0	0
			Total	N	O		
3	A	1	4	1	3	0	0
			Total	N	O		
3	B	1	4	1	3	0	0
			Total	N	O		
3	D	1	4	1	3	0	0
			Total	N	O		
3	D	1	4	1	3	0	0
			Total	N	O		
3	D	1	4	1	3	0	0
			Total	N	O		
3	D	1	4	1	3	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	D	1	Total O S 5 4 1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	117	Total O 117 117	0	0
5	B	99	Total O 99 99	0	0
5	C	103	Total O 103 103	0	0
5	D	129	Total O 129 129	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Azurin

Chain A:





- Molecule 1: Azurin

Chain B:



- Molecule 1: Azurin

Chain C:





- Molecule 1: Azurin

Chain D:





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	56.48Å 79.23Å 109.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.66 – 1.54 19.78 – 1.54	Depositor EDS
% Data completeness (in resolution range)	98.7 (19.66-1.54) 98.7 (19.78-1.54)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.14 (at 1.53Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.1_1168)	Depositor
R, R_{free}	0.211 , 0.239 0.213 , 0.240	Depositor DCC
R_{free} test set	3683 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å^2)	14.5	Xtriage
Anisotropy	0.182	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.45 , 42.1	EDS
L-test for twinning[2]	$< \|L\| > = 0.51$, $< L^2 > = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	8362	wwPDB-VP
Average B, all atoms (Å^2)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 35.20 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.0587e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, NO3, HG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.73	1/1026 (0.1%)	0.76	0/1381
1	B	0.63	0/1045	0.77	1/1405 (0.1%)
1	C	0.54	0/1009	0.70	0/1358
1	D	0.67	0/999	0.77	3/1345 (0.2%)
All	All	0.65	1/4079 (0.0%)	0.75	4/5489 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	2	GLU	CG-CD	-6.22	1.42	1.51

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	69	ASP	CB-CG-OD1	11.49	128.64	118.30
1	D	2	GLU	CA-CB-CG	5.30	125.07	113.40
1	D	38	ASN	CB-CA-C	-5.20	100.00	110.40
1	D	92	LYS	CD-CE-NZ	5.03	123.27	111.70

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	104[B]	GLU	Mainchain
1	B	1	ALA	Mainchain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1009	985	979	2	1
1	B	1001	985	965	9	1
1	C	983	968	957	9	0
1	D	982	962	960	5	0
2	A	3	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	8	0	0	0	0
3	B	4	0	0	0	0
3	D	16	0	0	0	0
4	D	5	0	0	0	0
5	A	117	0	0	0	0
5	B	99	0	0	0	0
5	C	103	0	0	0	0
5	D	129	0	0	0	0
All	All	4462	3900	3861	24	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 24 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:21:THR:HG23	1:A:128:LYS:HG2	1.90	0.52
1:B:14[B]:GLN:OE1	1:C:18:ASN:OD1	2.29	0.50
1:D:70:LYS:O	1:D:71:ASP:HB2	2.10	0.50
1:D:53:ALA:CB	1:D:107:GLN:HG3	2.43	0.49
1:D:107:GLN:O	1:D:107:GLN:HG2	2.13	0.49

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2:GLU:OE2	1:B:41:LYS:HG2[3_554]	1.36	0.24

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	130/128 (102%)	125 (96%)	5 (4%)	0	100	100
1	B	132/128 (103%)	128 (97%)	4 (3%)	0	100	100
1	C	128/128 (100%)	124 (97%)	4 (3%)	0	100	100
1	D	127/128 (99%)	124 (98%)	3 (2%)	0	100	100
All	All	517/512 (101%)	501 (97%)	16 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	114/110 (104%)	112 (98%)	2 (2%)	59	29
1	B	116/110 (106%)	114 (98%)	2 (2%)	60	31
1	C	112/110 (102%)	112 (100%)	0	100	100
1	D	111/110 (101%)	109 (98%)	2 (2%)	59	29

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	453/440 (103%)	447 (99%)	6 (1%)	78 42

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	28[B]	GLN
1	D	2	GLU
1	D	122	LYS
1	A	122[B]	LYS
1	A	122[A]	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 14 ligands modelled in this entry, 6 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	NO3	D	205	-	1,3,3	3.36	1 (100%)	0,3,3	-	-
3	NO3	A	203	-	1,3,3	3.25	1 (100%)	0,3,3	-	-
3	NO3	D	203	-	1,3,3	3.41	1 (100%)	0,3,3	-	-
3	NO3	B	202	-	1,3,3	3.86	1 (100%)	0,3,3	-	-
3	NO3	A	204	-	1,3,3	3.40	1 (100%)	0,3,3	-	-
3	NO3	D	204	-	1,3,3	3.49	1 (100%)	0,3,3	-	-
4	SO4	D	206	-	4,4,4	0.22	0	6,6,6	0.50	0
3	NO3	D	202	-	1,3,3	3.61	1 (100%)	0,3,3	-	-

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	202	NO3	O1-N	3.86	1.41	1.24
3	D	202	NO3	O1-N	3.61	1.40	1.24
3	D	204	NO3	O1-N	3.49	1.40	1.24
3	D	203	NO3	O1-N	3.41	1.39	1.24
3	A	204	NO3	O1-N	3.40	1.39	1.24

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	128/128 (100%)	0.26	6 (4%)	31	36	11, 17, 32, 54	0
1	B	128/128 (100%)	0.44	9 (7%)	16	18	13, 21, 38, 68	0
1	C	128/128 (100%)	0.59	10 (7%)	13	14	12, 21, 35, 64	0
1	D	128/128 (100%)	0.30	7 (5%)	25	28	9, 16, 30, 55	0
All	All	512/512 (100%)	0.40	32 (6%)	20	23	9, 18, 36, 68	0

The worst 5 of 32 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1	ALA	7.3
1	C	1	ALA	7.0
1	D	1	ALA	6.9
1	B	128	LYS	4.5
1	C	128	LYS	4.4

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	NO3	D	204	4/4	0.55	0.18	26,33,33,34	0
3	NO3	D	202	4/4	0.90	0.12	24,26,27,27	0
3	NO3	A	203	4/4	0.91	0.10	17,18,19,19	0
3	NO3	D	205	4/4	0.91	0.12	28,30,30,31	0
3	NO3	B	202	4/4	0.92	0.11	25,26,26,33	0
3	NO3	D	203	4/4	0.92	0.15	22,24,26,32	0
4	SO4	D	206	5/5	0.93	0.11	10,17,19,24	5
3	NO3	A	204	4/4	0.95	0.10	30,33,34,39	0
2	HG	A	202[A]	1/1	0.97	0.53	52,52,52,52	1
2	HG	A	202[B]	1/1	0.97	0.53	65,65,65,65	1
2	HG	C	201	1/1	0.99	0.05	16,16,16,16	1
2	HG	D	201	1/1	1.00	0.05	12,12,12,12	1
2	HG	B	201	1/1	1.00	0.03	15,15,15,15	1
2	HG	A	201	1/1	1.00	0.04	15,15,15,15	0

6.5 Other polymers (i)

There are no such residues in this entry.