



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 12:04 PM UTC

PDB ID : 9JMX / pdb\_00009jmx  
Title : Crystal structure of IRED-M235A in complex with NADP<sup>+</sup> and substrate  
Authors : Fu, X.L.; Zhang, M.T.; Yi, S.M.; Dai, K.X.; Yang, Y.Y.; Chen, C.-C.; Guo, R.-T.  
Deposited on : 2024-09-21  
Resolution : 2.04 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.04 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	2260 (2.04-2.04)
Clashscore	190562	2333 (2.04-2.04)
Ramachandran outliers	187476	2318 (2.04-2.04)
Sidechain outliers	187428	2318 (2.04-2.04)
RSRZ outliers	180081	2260 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	286	<div> <div>2%</div> <div> <div></div> <div>92%</div> <div>7%</div> </div> <div>.</div> </div>
1	B	286	<div> <div>8%</div> <div> <div></div> <div>92%</div> <div>7%</div> </div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	A1ECB	A	302	-	X	-	-

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4515 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 3-hydroxyisobutyrate dehydrogenase-like beta-hydroxyacid dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	283	Total	C	N	O	S	0	0	0
			2092	1320	366	400	6			
1	B	284	Total	C	N	O	S	0	0	0
			2100	1325	367	401	7			

There are 10 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	59	GLY	GLU	conflict	UNP A0A3N2GPT5
A	83	ALA	THR	conflict	UNP A0A3N2GPT5
A	106	ALA	THR	conflict	UNP A0A3N2GPT5
A	235	ALA	MET	engineered mutation	UNP A0A3N2GPT5
A	264	ASP	GLU	conflict	UNP A0A3N2GPT5
B	59	GLY	GLU	conflict	UNP A0A3N2GPT5
B	83	ALA	THR	conflict	UNP A0A3N2GPT5
B	106	ALA	THR	conflict	UNP A0A3N2GPT5
B	235	ALA	MET	engineered mutation	UNP A0A3N2GPT5
B	264	ASP	GLU	conflict	UNP A0A3N2GPT5

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NAP) (formula: C<sub>21</sub>H<sub>28</sub>N<sub>7</sub>O<sub>17</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 3 is 1- {tert}-butyl-4,9-dihydro-3 {H}-pyrido[3,4-b]indole (CCD ID: A1ECB) (formula: C<sub>15</sub>H<sub>18</sub>N<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	N	0	0
			17	15	2		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	N	0	0
			17	15	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	88	Total	O	0	0
			88	88		
4	B	105	Total	O	0	0
			105	105		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 3-hydroxyisobutyrate dehydrogenase-like beta-hydroxyacid dehydrogenase

Chain A: 



- Molecule 1: 3-hydroxyisobutyrate dehydrogenase-like beta-hydroxyacid dehydrogenase

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	78.75Å 78.75Å 160.41Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	34.59 – 2.04 34.59 – 2.04	Depositor EDS
% Data completeness (in resolution range)	99.5 (34.59-2.04) 99.5 (34.59-2.04)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.79 (at 2.05Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, $R_{free}$	0.206 , 0.260 0.214 , 0.267	Depositor DCC
$R_{free}$ test set	1854 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	30.0	Xtriage
Anisotropy	0.298	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 36.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.011 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4515	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.02% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A1ECB, NAP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.06	0/2133	1.45	0/2912
1	B	1.10	0/2141	1.39	3/2922 (0.1%)
All	All	1.08	0/4274	1.42	3/5834 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	143	HIS	CA-C-N	5.71	127.23	120.09
1	B	143	HIS	C-N-CA	5.71	127.23	120.09
1	B	284	ALA	CA-C-O	-5.06	112.20	120.80

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2092	0	2073	11	0
1	B	2100	0	2085	13	0
2	A	48	0	25	0	0
2	B	48	0	25	0	0
3	A	17	0	0	0	0
3	B	17	0	0	3	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	88	0	0	2	0
4	B	105	0	0	3	0
All	All	4515	0	4208	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:111:THR:HG21	1:A:139:VAL:HG11	1.85	0.58
4:A:420:HOH:O	3:B:302:A1ECB:CAL	2.53	0.56
1:B:71:MET:HE1	1:B:101:ALA:HB1	1.91	0.51
3:B:302:A1ECB:NAG	3:B:302:A1ECB:CAP	2.76	0.49
1:A:86:VAL:HG21	1:A:143:HIS:NE2	2.29	0.48
1:A:16:MET:HE1	1:A:64:SER:HB2	1.96	0.48
1:B:39:ASP:HB2	4:B:464:HOH:O	2.12	0.48
1:B:5:THR:HA	1:B:28:THR:O	2.15	0.46
1:A:17:VAL:HG11	1:A:40:VAL:HB	1.98	0.46
4:A:420:HOH:O	3:B:302:A1ECB:CAM	2.63	0.46
1:B:36:ARG:HD2	4:B:485:HOH:O	2.15	0.45
1:A:240:GLU:OE2	1:A:260:LYS:HE3	2.17	0.44
1:B:144:GLU:N	1:B:145:PRO:CD	2.81	0.44
1:A:144:GLU:N	1:A:145:PRO:CD	2.82	0.43
1:A:165:LEU:HG	1:B:188:LEU:HD22	2.00	0.43
1:B:135:GLY:O	1:B:155:TYR:OH	2.34	0.43
1:A:276:TRP:CZ3	1:B:178:LEU:HB3	2.54	0.42
1:A:71:MET:HE1	1:A:101:ALA:HB1	2.02	0.42
1:A:144:GLU:OE2	1:A:148:ARG:NH1	2.52	0.41
1:B:8:GLY:O	1:B:13:GLY:HA3	2.21	0.41
1:A:174:PHE:HB2	1:B:205:VAL:HG22	2.02	0.41
1:B:137:ARG:NH2	4:B:412:HOH:O	2.53	0.40
1:B:179:SER:HB3	1:B:255:LEU:HD22	2.03	0.40
1:B:51:PRO:O	1:B:55:LEU:HG	2.20	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	281/286 (98%)	274 (98%)	7 (2%)	0	100	100
1	B	282/286 (99%)	274 (97%)	8 (3%)	0	100	100
All	All	563/572 (98%)	548 (97%)	15 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	206/209 (99%)	204 (99%)	2 (1%)	68	72
1	B	207/209 (99%)	205 (99%)	2 (1%)	68	72
All	All	413/418 (99%)	409 (99%)	4 (1%)	68	72

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	41	VAL
1	A	228	ASP
1	B	188	LEU
1	B	205	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	160	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	A1ECB	B	302	-	18,19,19	4.65	10 (55%)	26,29,29	2.52	8 (30%)
3	A1ECB	A	302	-	18,19,19	4.76	9 (50%)	26,29,29	3.32	14 (53%)
2	NAP	B	301	-	50,52,52	1.30	6 (12%)	71,80,80	1.82	19 (26%)
2	NAP	A	301	-	50,52,52	1.49	9 (18%)	71,80,80	1.97	21 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	A1ECB	B	302	-	-	6/6/16/16	0/3/3/3

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	A1ECB	A	302	-	-	6/6/16/16	0/3/3/3
2	NAP	B	301	-	-	7/35/67/67	0/5/5/5
2	NAP	A	301	-	-	9/35/67/67	0/5/5/5

All (34) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	302	A1ECB	CAM-CAI	-13.84	1.32	1.50
3	B	302	A1ECB	CAM-CAI	-13.15	1.33	1.50
3	A	302	A1ECB	CAE-CAI	-7.89	1.29	1.44
3	B	302	A1ECB	CAE-CAI	-6.56	1.32	1.44
3	B	302	A1ECB	CAE-CAD	-6.37	1.33	1.41
3	A	302	A1ECB	CAC-CAD	-6.04	1.30	1.39
3	A	302	A1ECB	CAE-CAD	-5.62	1.34	1.41
3	B	302	A1ECB	CAC-CAD	-5.51	1.30	1.39
2	A	301	NAP	PN-O3	5.38	1.65	1.59
3	B	302	A1ECB	CAN-CAJ	-5.37	1.42	1.52
2	B	301	NAP	O4D-C1D	4.31	1.46	1.40
3	A	302	A1ECB	CAF-CAE	-4.24	1.33	1.39
3	B	302	A1ECB	CAF-CAE	-4.19	1.33	1.39
3	B	302	A1ECB	CAH-NAG	-4.11	1.32	1.38
3	A	302	A1ECB	CAJ-NAK	4.02	1.32	1.28
3	A	302	A1ECB	CAH-NAG	-3.96	1.32	1.38
2	A	301	NAP	C5A-C4A	3.69	1.45	1.39
3	A	302	A1ECB	CAD-NAG	-3.67	1.32	1.38
2	B	301	NAP	C5A-C4A	3.53	1.45	1.39
3	B	302	A1ECB	CAJ-NAK	3.49	1.32	1.28
3	A	302	A1ECB	CAN-CAJ	-3.38	1.46	1.52
3	B	302	A1ECB	CAD-NAG	-3.33	1.32	1.38
2	B	301	NAP	C5A-N7A	-3.23	1.33	1.39
2	A	301	NAP	O4D-C1D	3.20	1.45	1.40
3	B	302	A1ECB	CAH-CAI	-2.89	1.34	1.38
2	B	301	NAP	PA-O3	2.36	1.62	1.59
2	A	301	NAP	C3N-C7N	-2.30	1.47	1.50
2	A	301	NAP	C4A-N9A	-2.30	1.32	1.37
2	B	301	NAP	C4A-N9A	-2.29	1.32	1.37
2	A	301	NAP	PA-O3	2.16	1.61	1.59
2	A	301	NAP	C8A-N7A	2.13	1.35	1.31
2	A	301	NAP	C2N-N1N	-2.05	1.32	1.35
2	A	301	NAP	O4B-C4B	-2.05	1.40	1.45
2	B	301	NAP	C8A-N7A	2.04	1.35	1.31

All (62) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	302	A1ECB	CAL-CAM-CAI	-10.29	95.64	109.14
3	B	302	A1ECB	CAN-CAJ-CAH	6.59	125.06	119.46
3	A	302	A1ECB	CAF-CAE-CAD	6.23	125.30	118.84
3	A	302	A1ECB	CAN-CAJ-CAH	6.06	124.61	119.46
3	B	302	A1ECB	CAL-CAM-CAI	-5.40	102.05	109.14
2	B	301	NAP	C5A-C4A-N3A	-4.99	119.84	126.72
2	A	301	NAP	C4A-N9A-C8A	4.86	110.84	105.74
3	B	302	A1ECB	CAQ-CAN-CAJ	4.57	117.78	109.97
2	A	301	NAP	N3A-C4A-N9A	4.52	134.85	127.17
2	B	301	NAP	N3A-C4A-N9A	4.39	134.63	127.17
2	A	301	NAP	C5A-C4A-N3A	-4.15	121.00	126.72
3	A	302	A1ECB	CAQ-CAN-CAJ	4.14	117.04	109.97
3	B	302	A1ECB	CAI-CAH-NAG	-4.13	105.86	109.59
2	A	301	NAP	N3A-C2A-N1A	-4.00	122.53	128.58
2	B	301	NAP	C2B-C1B-N9A	3.90	120.17	113.75
2	A	301	NAP	O7N-C7N-C3N	-3.68	115.10	119.60
2	A	301	NAP	C5N-C4N-C3N	-3.62	116.81	120.36
2	A	301	NAP	O2A-PA-O1A	3.61	129.25	112.44
2	B	301	NAP	C3N-C7N-N7N	3.58	122.16	117.74
3	B	302	A1ECB	CAF-CAE-CAD	3.53	122.50	118.84
2	B	301	NAP	O2A-PA-O3	3.44	116.56	107.27
2	B	301	NAP	C2A-N3A-C4A	3.36	120.03	111.83
2	B	301	NAP	N3A-C2A-N1A	-3.33	123.53	128.58
2	B	301	NAP	C4A-N9A-C8A	3.21	109.11	105.74
2	A	301	NAP	N6A-C6A-N1A	3.12	125.34	118.38
3	B	302	A1ECB	CAE-CAI-CAH	3.06	110.33	106.79
2	A	301	NAP	N9A-C8A-N7A	-3.04	109.63	113.94
2	B	301	NAP	C4D-O4D-C1D	3.03	112.70	109.92
3	A	302	A1ECB	CAF-CAE-CAI	-3.01	128.59	133.73
2	A	301	NAP	C2N-C3N-C4N	3.00	121.74	118.26
2	A	301	NAP	C2A-N3A-C4A	2.98	119.12	111.83
2	B	301	NAP	O2A-PA-O1A	2.96	126.19	112.44
3	A	302	A1ECB	CAN-CAJ-NAK	2.90	121.54	117.16
3	A	302	A1ECB	CAI-CAH-NAG	-2.88	106.99	109.59
3	B	302	A1ECB	CAB-CAA-CAF	-2.87	116.70	120.24
3	A	302	A1ECB	CAQ-CAN-CAO	-2.85	101.29	108.95
2	A	301	NAP	C3N-C7N-N7N	2.85	121.25	117.74
2	B	301	NAP	N9A-C8A-N7A	-2.79	109.97	113.94
2	A	301	NAP	C5A-C6A-N6A	-2.75	116.48	123.29
3	A	302	A1ECB	CAE-CAI-CAH	2.62	109.83	106.79
3	A	302	A1ECB	CAM-CAI-CAE	-2.60	125.82	130.72
3	A	302	A1ECB	CAA-CAB-CAC	2.59	123.44	120.24

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	NAP	C2B-C1B-N9A	2.59	118.01	113.75
2	A	301	NAP	C6N-N1N-C2N	-2.58	119.68	121.88
2	B	301	NAP	O7N-C7N-C3N	-2.54	116.50	119.60
3	B	302	A1ECB	CAP-CAN-CAJ	-2.52	105.66	109.97
2	A	301	NAP	C2A-N1A-C6A	2.49	122.83	118.73
2	B	301	NAP	C5A-N7A-C8A	2.49	107.36	103.45
2	A	301	NAP	O3X-P2B-O2B	2.43	115.31	105.85
2	A	301	NAP	C2N-N1N-C1D	2.32	124.25	119.13
2	B	301	NAP	C4A-C5A-N7A	-2.23	108.03	110.58
2	B	301	NAP	O3B-C3B-C4B	-2.13	104.95	111.08
2	A	301	NAP	O2B-P2B-O1X	-2.13	101.74	109.33
2	A	301	NAP	C4A-N9A-C1B	-2.12	121.68	126.63
2	A	301	NAP	C4N-C3N-C7N	-2.08	115.39	121.06
3	A	302	A1ECB	CAJ-CAH-NAG	2.08	133.23	129.07
2	B	301	NAP	O2N-PN-O3	2.06	112.83	107.27
2	B	301	NAP	C3B-C2B-C1B	-2.05	98.88	102.81
3	A	302	A1ECB	CAB-CAA-CAF	-2.03	117.74	120.24
3	A	302	A1ECB	CAA-CAF-CAE	-2.02	116.14	119.80
2	B	301	NAP	C6N-N1N-C1D	-2.02	115.76	119.73
2	B	301	NAP	C5A-C6A-N6A	-2.01	118.30	123.29

There are no chirality outliers.

All (28) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	NAP	PA-O3-PN-O5D
2	A	301	NAP	O4D-C1D-N1N-C2N
2	A	301	NAP	O4D-C1D-N1N-C6N
2	A	301	NAP	C2D-C1D-N1N-C2N
2	A	301	NAP	C2D-C1D-N1N-C6N
2	B	301	NAP	PA-O3-PN-O5D
2	B	301	NAP	O4D-C1D-N1N-C2N
2	B	301	NAP	O4D-C1D-N1N-C6N
2	B	301	NAP	C2D-C1D-N1N-C2N
2	B	301	NAP	C2D-C1D-N1N-C6N
2	A	301	NAP	C1B-C2B-O2B-P2B
2	A	301	NAP	C3B-C2B-O2B-P2B
3	B	302	A1ECB	CAH-CAJ-CAN-CAQ
3	A	302	A1ECB	CAH-CAJ-CAN-CAQ
3	B	302	A1ECB	NAK-CAJ-CAN-CAO
3	A	302	A1ECB	NAK-CAJ-CAN-CAQ
3	B	302	A1ECB	NAK-CAJ-CAN-CAQ

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
3	B	302	A1ECB	CAH-CAJ-CAN-CAO
2	A	301	NAP	C3B-C4B-C5B-O5B
3	A	302	A1ECB	NAK-CAJ-CAN-CAO
3	A	302	A1ECB	NAK-CAJ-CAN-CAP
2	A	301	NAP	O4B-C4B-C5B-O5B
3	B	302	A1ECB	NAK-CAJ-CAN-CAP
3	A	302	A1ECB	CAH-CAJ-CAN-CAO
3	A	302	A1ECB	CAH-CAJ-CAN-CAP
2	B	301	NAP	PN-O3-PA-O1A
2	B	301	NAP	C1B-C2B-O2B-P2B
3	B	302	A1ECB	CAH-CAJ-CAN-CAP

There are no ring outliers.


1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	302	A1ECB	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




## Ligand A1ECB B 302




Bond lengths



Bond angles





Torsions




Rings

## Ligand A1ECB A 302







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	283/286 (98%)	0.43	7 (2%) 58 60	20, 35, 55, 68	0
1	B	284/286 (99%)	0.49	23 (8%) 18 18	21, 35, 58, 84	0
All	All	567/572 (99%)	0.46	30 (5%) 32 32	20, 35, 57, 84	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	46	VAL	3.7
1	B	74	VAL	3.1
1	B	103	ALA	3.0
1	B	47	LEU	3.0
1	B	48	ALA	2.9
1	B	52	ALA	2.8
1	B	82	LEU	2.7
1	B	81	ALA	2.6
1	B	110	ALA	2.6
1	B	30	TRP	2.5
1	A	2	THR	2.5
1	B	101	ALA	2.5
1	A	81	ALA	2.5
1	B	1	MET	2.4
1	B	79	GLY	2.4
1	B	106	ALA	2.3
1	B	46	VAL	2.3
1	B	78	ALA	2.3
1	A	150	ILE	2.3
1	A	106	ALA	2.3
1	B	57	ALA	2.3
1	B	221	ARG	2.2
1	B	44	GLY	2.2
1	A	47	LEU	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	34	LYS	2.2
1	B	4	ALA	2.1
1	B	35	SER	2.1
1	B	22	GLU	2.1
1	B	40	VAL	2.1
1	A	45	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	A1ECB	A	302	17/17	0.84	0.15	32,42,47,48	0
3	A1ECB	B	302	17/17	0.87	0.12	33,39,46,47	0
2	NAP	B	301	48/48	0.96	0.07	26,32,44,48	0
2	NAP	A	301	48/48	0.97	0.06	23,29,34,36	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around A1ECB A 302:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around A1ECB B 302:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around NAP B 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around NAP A 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.