



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 17, 2020 – 12:53 PM BST

PDB ID : 5JPN
Title : Structure of human complement C4 rebuilt using iMDFP
Authors : Croll, T.I.; Andersen, G.R.
Deposited on : 2016-05-03
Resolution : 3.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1257 (3.70-3.50)
Clashscore	141614	1353 (3.70-3.50)
Ramachandran outliers	138981	1307 (3.70-3.50)
Sidechain outliers	138945	1307 (3.70-3.50)
RSRZ outliers	127900	1161 (3.70-3.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	656	 4% 93% 6%
2	B	767	 3% 87% 9%
3	C	290	 % 92% 8%
4	D	3	 100%
5	E	2	 50% 50%
6	F	5	 40% 60%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
6	G	5	 A horizontal bar chart showing the quality of chain. The bar is divided into two segments: a green segment on the left labeled '40%' and a yellow segment on the right labeled '60%'. The total length of the bar represents 100%.

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	MAN	F	4	-	-	-	X
6	MAN	F	5	-	-	-	X
6	MAN	G	5	-	-	-	X

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 12891 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Complement C4-A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	651	5012	3185	872	939	16	0	0	0

- Molecule 2 is a protein called Complement C4-A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	697	5361	3371	943	1023	24	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	1201	SER	THR	variant	UNP P0C0L4

- Molecule 3 is a protein called Complement C4-A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	290	2311	1451	411	432	17	0	0	0

- Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	D	3	39	22	2	15	0	0	0

- Molecule 5 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
5	E	2	28	16	2	10	0	0	0

- Molecule 6 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
6	F	5	61	34	2	25	0	0	0
6	G	5	61	34	2	25	0	0	0

- Molecule 7 is TRIMETHYL LEAD ION (three-letter code: PBM) (formula: C₃H₉Pb).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total Pb 1 1	0	0
7	B	1	Total Pb 1 1	0	0
7	B	1	Total Pb 1 1	0	0
7	C	1	Total Pb 1 1	0	0

- Molecule 8 is 2-acetamido-2-deoxy-alpha-D-galactopyranose (three-letter code: A2G) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	B	1	Total C N O 14 8 1 5	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Complement C4-A



- Molecule 2: Complement C4-A



- Molecule 3: Complement C4-A



- Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




MAG1
MAG2
BMA3

- Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  50% 50%

MAG1
MAG2

- Molecule 6: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  40% 60%

MAG1
MAG2
BMA3
MAN4
MAN5

- Molecule 6: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  40% 60%

MAG1
MAG2
BMA3
MAN4
MAN5

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	85.45Å 103.31Å 256.03Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.90 – 3.60 45.90 – 3.60	Depositor EDS
% Data completeness (in resolution range)	99.6 (45.90-3.60) 99.7 (45.90-3.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.48 (at 3.57Å)	Xtrriage
Refinement program	PHENIX dev_2376	Depositor
R, R_{free}	0.215 , 0.263 0.215 , 0.263	Depositor DCC
R_{free} test set	1348 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	113.2	Xtrriage
Anisotropy	0.501	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 85.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	12891	wwPDB-VP
Average B, all atoms (Å ²)	130.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PBM, BMA, NAG, A2G, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/5128	0.46	0/6961
2	B	0.24	0/5472	0.44	1/7436 (0.0%)
3	C	0.24	0/2359	0.48	0/3188
All	All	0.25	0/12959	0.45	1/17585 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	783	LEU	CA-CB-CG	6.06	129.23	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	582	ARG	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5012	0	5025	16	0
2	B	5361	0	5310	12	0
3	C	2311	0	2253	7	0
4	D	39	0	34	0	0
5	E	28	0	25	0	0
6	F	61	0	52	0	0
6	G	61	0	52	0	0
7	A	1	0	0	0	0
7	B	2	0	0	0	0
7	C	1	0	0	0	0
8	B	14	0	12	0	0
All	All	12891	0	12763	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:476:GLU:HB3	1:A:492:ASN:HB2	1.71	0.72
1:A:100:HIS:CG	1:A:101:GLN:H	2.08	0.71
2:B:1325:ARG:HB3	2:B:1374:SER:HB2	1.89	0.55
1:A:81:ASP:N	1:A:81:ASP:OD1	2.39	0.53
2:B:1013:GLN:N	2:B:1013:GLN:OE1	2.38	0.49
3:C:1667:PHE:HB3	3:C:1703:LEU:HD21	1.94	0.49
2:B:1191:HIS:CE1	2:B:1195:ILE:HD11	2.47	0.49
1:A:100:HIS:CG	1:A:101:GLN:N	2.79	0.49
3:C:1685:ILE:HA	3:C:1708:TRP:O	2.13	0.48
2:B:978:VAL:HG12	2:B:1380:VAL:HG22	1.95	0.48
3:C:1547:GLY:HA3	3:C:1582:LEU:HD22	1.93	0.48
2:B:831:ARG:HH11	2:B:831:ARG:HA	1.79	0.48
3:C:1583:CYS:HB2	3:C:1587:VAL:HG23	1.97	0.46
1:A:137:ARG:HG3	1:A:225:SER:HA	1.97	0.46
2:B:847:GLU:O	2:B:903:PRO:HD2	2.16	0.46
2:B:1352:ARG:HA	2:B:1355:GLU:OE2	2.16	0.46
1:A:255:VAL:HG12	1:A:257:GLY:H	1.81	0.45
2:B:1399:ILE:HD12	3:C:1556:LEU:HB2	1.99	0.45
2:B:832:GLU:H	2:B:832:GLU:CD	2.20	0.44
1:A:206:ASP:N	1:A:206:ASP:OD2	2.45	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:1685:ILE:HG22	3:C:1709:ILE:HG12	2.00	0.44
2:B:948:LEU:HD22	2:B:1370:VAL:HG21	2.00	0.44
2:B:947:GLU:OE2	2:B:1291:ARG:NH1	2.51	0.43
3:C:1623:VAL:HG21	3:C:1686:MET:HB2	2.01	0.43
1:A:338:LEU:N	1:A:360:TRP:O	2.48	0.43
1:A:425:THR:HG22	1:A:431:VAL:HG23	2.01	0.43
1:A:247:PRO:HB3	1:A:360:TRP:CZ2	2.54	0.42
1:A:377:HIS:CG	1:A:640:GLY:HA2	2.54	0.42
1:A:315:SER:HB2	1:A:318:GLU:HG3	2.02	0.42
2:B:1302:THR:O	2:B:1306:VAL:HG12	2.20	0.41
1:A:374:THR:O	1:A:376:ARG:NH1	2.47	0.40
1:A:173:THR:HG22	1:A:192:TYR:HA	2.03	0.40
1:A:582:ARG:HA	1:A:582:ARG:HD2	1.77	0.40
1:A:59:ARG:HD2	1:A:65:ASN:HB3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	649/656 (99%)	618 (95%)	30 (5%)	1 (0%)	47 79
2	B	693/767 (90%)	661 (95%)	30 (4%)	2 (0%)	41 75
3	C	288/290 (99%)	272 (94%)	15 (5%)	1 (0%)	41 75
All	All	1630/1713 (95%)	1551 (95%)	75 (5%)	4 (0%)	47 79

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	1390	LYS
1	A	101	GLN
3	C	1622	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	1244	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	557/562 (99%)	542 (97%)	15 (3%)	44 73
2	B	581/641 (91%)	568 (98%)	13 (2%)	52 77
3	C	248/248 (100%)	237 (96%)	11 (4%)	28 63
All	All	1386/1451 (96%)	1347 (97%)	39 (3%)	43 72

All (39) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	87	LEU
1	A	88	GLN
1	A	130	ILE
1	A	156	VAL
1	A	254	THR
1	A	260	ASP
1	A	314	LEU
1	A	320	GLN
1	A	345	ILE
1	A	360	TRP
1	A	422	GLN
1	A	524	ARG
1	A	571	LEU
1	A	580	GLN
1	A	616	HIS
2	B	745	LYS
2	B	796	THR
2	B	828	ARG
2	B	831	ARG
2	B	837	LEU
2	B	1010	CYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	1031	THR
2	B	1244	THR
2	B	1304	ASP
2	B	1328	ASN
2	B	1346	LEU
2	B	1355	GLU
2	B	1390	LYS
3	C	1460	GLU
3	C	1465	ARG
3	C	1477	LYS
3	C	1582	LEU
3	C	1588	CYS
3	C	1592	GLU
3	C	1630	LYS
3	C	1658	LYS
3	C	1665	ARG
3	C	1719	ARG
3	C	1734	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

15 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	D	1	1,4	14,14,15	0.30	0	17,19,21	0.44	0
4	NAG	D	2	4	14,14,15	0.22	0	17,19,21	0.58	0
4	BMA	D	3	4	11,11,12	0.61	0	15,15,17	0.74	0
5	NAG	E	1	2,5	14,14,15	0.62	1 (7%)	17,19,21	0.68	0
5	NAG	E	2	5	14,14,15	0.30	0	17,19,21	0.45	0
6	NAG	F	1	2,6	14,14,15	0.29	0	17,19,21	0.53	0
6	NAG	F	2	6	14,14,15	0.39	0	17,19,21	0.56	0
6	BMA	F	3	6	11,11,12	0.70	0	15,15,17	0.92	1 (6%)
6	MAN	F	4	6	11,11,12	0.66	0	15,15,17	0.95	2 (13%)
6	MAN	F	5	6	11,11,12	0.79	0	15,15,17	0.84	1 (6%)
6	NAG	G	1	2,6	14,14,15	0.43	0	17,19,21	0.53	0
6	NAG	G	2	6	14,14,15	0.21	0	17,19,21	0.65	1 (5%)
6	BMA	G	3	6	11,11,12	0.55	0	15,15,17	0.77	0
6	MAN	G	4	6	11,11,12	0.73	0	15,15,17	0.89	1 (6%)
6	MAN	G	5	6	11,11,12	0.72	0	15,15,17	0.89	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	D	1	1,4	-	1/6/23/26	0/1/1/1
4	NAG	D	2	4	-	0/6/23/26	0/1/1/1
4	BMA	D	3	4	-	0/2/19/22	0/1/1/1
5	NAG	E	1	2,5	-	0/6/23/26	0/1/1/1
5	NAG	E	2	5	-	0/6/23/26	0/1/1/1
6	NAG	F	1	2,6	-	0/6/23/26	0/1/1/1
6	NAG	F	2	6	-	0/6/23/26	0/1/1/1
6	BMA	F	3	6	-	0/2/19/22	0/1/1/1
6	MAN	F	4	6	-	1/2/19/22	0/1/1/1
6	MAN	F	5	6	-	0/2/19/22	0/1/1/1
6	NAG	G	1	2,6	-	0/6/23/26	0/1/1/1
6	NAG	G	2	6	-	0/6/23/26	0/1/1/1
6	BMA	G	3	6	-	0/2/19/22	0/1/1/1
6	MAN	G	4	6	-	1/2/19/22	0/1/1/1
6	MAN	G	5	6	-	0/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	E	1	NAG	O5-C1	-2.07	1.40	1.43

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	F	4	MAN	C1-O5-C5	2.27	115.26	112.19
6	F	5	MAN	O2-C2-C3	-2.26	105.61	110.14
6	G	4	MAN	O2-C2-C3	-2.25	105.64	110.14
6	G	5	MAN	O2-C2-C3	-2.23	105.67	110.14
6	F	4	MAN	O2-C2-C3	-2.18	105.77	110.14
6	G	2	NAG	C1-O5-C5	2.16	115.12	112.19
6	F	3	BMA	C1-O5-C5	2.03	114.94	112.19

There are no chirality outliers.


All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	F	4	MAN	O5-C5-C6-O6
6	G	4	MAN	O5-C5-C6-O6
4	D	1	NAG	O5-C5-C6-O6


There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 4 are modelled with single atom - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
8	A2G	B	1508	2	14,14,15	1.78	5 (35%)	17,19,21	1.12	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	A2G	B	1508	2	-	1/6/23/26	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	B	1508	A2G	C7-N2	3.29	1.45	1.34
8	B	1508	A2G	O5-C1	2.33	1.47	1.43
8	B	1508	A2G	C3-C2	-2.24	1.47	1.52
8	B	1508	A2G	O3-C3	2.12	1.48	1.43
8	B	1508	A2G	C2-N2	2.07	1.49	1.46

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	B	1508	A2G	C8-C7-N2	2.21	119.85	116.10
8	B	1508	A2G	C2-N2-C7	-2.18	119.80	122.90

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	B	1508	A2G	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	651/656 (99%)	0.14	24 (3%) 41 27	77, 131, 198, 234	0
2	B	697/767 (90%)	0.04	21 (3%) 50 34	76, 118, 190, 231	0
3	C	290/290 (100%)	-0.05	2 (0%) 87 78	79, 118, 162, 191	0
All	All	1638/1713 (95%)	0.07	47 (2%) 51 35	76, 123, 191, 234	0

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	918	SER	6.0
2	B	743	ARG	5.2
2	B	1237	GLN	4.0
1	A	339	TYR	3.5
2	B	739	ALA	3.4
2	B	1326	GLY	3.2
2	B	1236	SER	3.2
3	C	1744	VAL	3.2
1	A	615	SER	3.2
1	A	296	PHE	3.2
1	A	326	LEU	3.1
1	A	282	VAL	3.0
1	A	286	LEU	2.9
2	B	1159	GLU	2.9
1	A	346	GLU	2.8
2	B	920	GLU	2.8
1	A	288	ASP	2.8
1	A	329	GLY	2.7
1	A	352	MET	2.6
2	B	1129	GLN	2.6
1	A	333	LEU	2.6
1	A	276	VAL	2.6
1	A	287	LEU	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	707	VAL	2.5
2	B	736	CYS	2.4
1	A	283	ARG	2.4
2	B	742	LEU	2.3
1	A	314	LEU	2.3
3	C	1477	LYS	2.3
1	A	280	ALA	2.3
2	B	1135	ASN	2.3
1	A	328	MET	2.2
2	B	1240	ALA	2.2
2	B	827	LEU	2.2
1	A	292	LYS	2.2
1	A	325	LYS	2.2
2	B	1130	GLY	2.2
2	B	744	LYS	2.2
1	A	260	ASP	2.1
2	B	897	VAL	2.1
1	A	338	LEU	2.1
2	B	1234	THR	2.1
2	B	1247	PRO	2.1
1	A	398	ALA	2.1
1	A	284	PHE	2.0
1	A	242	GLU	2.0
2	B	830	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	MAN	G	5	11/12	0.55	0.59	206,211,223,225	0
6	BMA	F	3	11/12	0.67	0.26	190,197,204,208	0
5	NAG	E	2	14/15	0.69	0.37	218,233,237,240	0
6	MAN	G	4	11/12	0.71	0.30	192,205,209,211	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	MAN	F	4	11/12	0.72	0.75	176,188,194,195	0
6	MAN	F	5	11/12	0.72	0.52	174,183,191,194	0
6	BMA	G	3	11/12	0.77	0.37	189,196,206,206	0
5	NAG	E	1	14/15	0.83	0.19	210,214,220,228	0
4	BMA	D	3	11/12	0.84	0.21	171,178,184,185	0
4	NAG	D	2	14/15	0.84	0.27	181,187,192,199	0
6	NAG	F	2	14/15	0.84	0.26	162,194,205,210	0
6	NAG	G	1	14/15	0.86	0.35	171,182,193,197	0
6	NAG	F	1	14/15	0.90	0.26	153,155,163,175	0
6	NAG	G	2	14/15	0.92	0.38	178,183,185,188	0
4	NAG	D	1	14/15	0.94	0.18	138,151,167,172	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain F:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain G:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
7	PBM	B	1515	1/4	0.50	0.33	137,137,137,137	1
8	A2G	B	1508	14/15	0.76	0.38	157,178,183,184	14
7	PBM	C	1801	1/4	0.88	0.21	123,123,123,123	1
7	PBM	A	704	1/4	0.94	0.19	134,134,134,134	1
7	PBM	B	1514	1/4	0.95	0.07	142,142,142,142	1

6.5 Other polymers [i](#)

There are no such residues in this entry.