# wwPDB NMR Structure Validation Summary Report (i)

May 28, 2020 – 10:40 pm BST

| | | |
|---|---|---|
| PDB ID | : | 2KRL |
| Title | : | The ensemble of the solution global structures of the 102-nt ribosome binding structure element of the turnip crinkle virus 3' UTR RNA |
| Authors | : | Zuo, X.; Wang, J.; Yu, P.; Eyler, D.; Xu, H.; Starich, M.; Tiede, D.; Simon, A.; Kasprzak, W.; Schwieters, C.; Shapiro, B. |
| Deposited on | : | 2009-12-18 |

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

| | | |
|---|---|---|
| Cyrange | : | Kirchner and Güntert (2011) |
| NmrClust | : | Kelley et al. (1996) |
| MolProbity | : | 4.02b-467 |
| Percentile statistics | : | 20191225.v01 (using entries in the PDB archive December 25th 2019) |
| RCI | : | v_1n_11_5_13_A (Berjanski et al., 2005) |
| PANAV | : | Wang et al. (2010) |
| ShiftChecker | : | 2.11 |
| Ideal geometry (proteins) | : | Engh & Huber (2001) |
| Ideal geometry (DNA, RNA) | : | Parkinson et al. (1996) |
| Validation Pipeline (wwPDB-VP) | : | 2.11 |

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
*SOLUTION NMR, SOLUTION SCATTERING*

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




| Metric | Whole archive (#Entries) | NMR archive (#Entries) |
|---|---|---|
| Clashscore | 158937 | 12864 |
| RNA backbone | 4643 | 676 |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

| Mol | Chain | Length | Quality of chain |
|---|---|---|---|
| 1 | A | 102 | |

## 2 Ensemble composition and analysis (i)

This entry contains 10 models. This entry does not contain polypeptide chains, therefore identification of well-defined residues and clustering analysis are not possible. All residues are included in the validation scores.

# 3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 3284 atoms, of which 1102 are hydrogens and 0 are deuteriums.

- Molecule 1 is a RNA chain called RNA (102-MER).

| Mol | Chain | Residues | Atoms | | | | | | Trace |
|---|---|---|---|---|---|---|---|---|---|
| | | | Total | C | H | N | O | P | |
| 1 | A | 102 | 3284 | 974 | 1102 | 396 | 711 | 101 | 0 |

There are 3 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|---|---|---|---|---|---|
| A | 1 | G | - | INSERTION | GB M22445.2 |
| A | 2 | G | - | INSERTION | GB M22445.2 |
| A | 98 | C | U | CONFLICT | GB M22445.2 |

# 4 Residue-property plots (i)

## 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: RNA (102-MER)

Chain A: 25% 65% 11%




## 4.2 Residue scores for the representative (author defined) model from the NMR ensemble

The representative model is number 1. Colouring as in section 4.1 above.

- Molecule 1: RNA (102-MER)

Chain A: 24% 61% 15%



# 5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *simulated annealing, duplex-orientation restrained.*

Of the 100 calculated structures, 10 were deposited, based on the following criterion: *structures with the lowest energy.*

The following table shows the software used for structure solution, optimisation and refinement.

| Software name | Classification | Version |
|---|---|---|
| X-PLOR NIH | refinement | 2.22 |

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

COVALENT-GEOMETRY INFOmissingINFO

## 5.1 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes |
|---|---|---|---|---|---|
| 1 | A | 2182 | 1102 | 1098 | 407±12 |
| All | All | 21820 | 11020 | 11004 | 4070 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 124.

5 of 1342 unique clashes are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Clash(Å) | Distance(Å) | Models | |
|---|---|---|---|---|---|
| | | | | Worst | Total |
| 1:A:81:G:N3 | 1:A:82:G:N7 | 1.22 | 1.84 | 3 | 1 |
| 1:A:99:U:H2' | 1:A:100:C:O4' | 1.02 | 1.54 | 1 | 10 |
| 1:A:23:C:N3 | 1:A:28:G:O5' | 1.01 | 1.93 | 7 | 10 |
| 1:A:12:G:O2' | 1:A:13:C:O5' | 1.01 | 1.76 | 9 | 1 |
| 1:A:53:C:O2' | 1:A:54:U:H5" | 0.99 | 1.56 | 8 | 10 |

## 5.2 Torsion angles (i)

### 5.2.1 Protein backbone (i)

There are no protein molecules in this entry.

### 5.2.2 Protein sidechains (i)

There are no protein molecules in this entry.

### 5.2.3 RNA (i)

| Mol | Chain | Analysed | Backbone Outliers | Pucker Outliers | Suiteness |
|---|---|---|---|---|---|
| 1 | A | 101/102 (99%) | 69±3 (68±3%) | 31±2 (30±2%) | 0.10±0.01 |
| All | All | 1014/1020 (99%) | 686 (68%) | 305 (30%) | 0.10 |

The overall RNA backbone suiteness is 0.10.

5 of 86 unique RNA backbone outliers are listed below:

| Mol | Chain | Res | Type | Models (Total) |
|---|---|---|---|---|
| 1 | A | 91 | G | 10 |
| 1 | A | 69 | A | 10 |
| 1 | A | 13 | C | 10 |
| 1 | A | 86 | G | 10 |
| 1 | A | 88 | U | 10 |

5 of 60 unique RNA pucker outliers are listed below:

| Mol | Chain | Res | Type | Models (Total) |
|---|---|---|---|---|
| 1 | A | 26 | C | 10 |
| 1 | A | 87 | C | 10 |
| 1 | A | 53 | C | 10 |
| 1 | A | 90 | U | 10 |
| 1 | A | 81 | G | 10 |

## 5.3 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.4 Carbohydrates (i)

There are no carbohydrates in this entry.

## 5.5 Ligand geometry (i)

There are no ligands in this entry.

## 5.6 Other polymers (i)

There are no such molecules in this entry.

## 5.7 Polymer linkage issues (i)

There are no chain breaks in this entry.

# 6 Chemical shift validation (i)

No chemical shift data were provided