Full wwPDB NMR Structure Validation Report (i)

Apr 16, 2023 – 06:48 AM EDT

PDB ID	:	2L7H
BMRB ID	:	6795
Title	:	The solution structure of the HAMP domain of the hypothetical transmembrane receptor Af1503
Authors	:	Coles, M.; Hulko, M.; Martin, J.; Lupas, A.N.
Deposited on	:	2010-12-09

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.32.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment is 38%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	58	71% 14% 16%
1	B	58	69% 14% 17%

2 Ensemble composition and analysis (i)

This entry contains 18 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:282-A:330, B:282-B:329 (97)	0.17	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 2 single-model clusters were found.

Cluster number	Models
1	1, 3, 8, 10, 12, 13, 15
2	2, 5, 6, 16, 17
3	4, 11, 14, 18
Single-model clusters	7; 9

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1822 atoms, of which 926 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Uncharacterized protein.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	58	911	273	463	84	89	2	0
1	B	58	911	273	463	84	89	2	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	274	GLY	-	expression tag	UNP O28769
A	275	SER	-	expression tag	UNP O28769
A	276	HIS	-	expression tag	UNP O28769
A	277	MET	-	expression tag	UNP O28769
B	274	GLY	-	expression tag	UNP O28769
B	275	SER	-	expression tag	UNP O28769
B	276	HIS	-	expression tag	UNP O28769
B	277	MET	-	expression tag	UNP O28769

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: Uncharacterized protein

Chain A:

• Molecule 1: Uncharacterized protein

Chain B:

4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1 (medoid)

• Molecule 1: Uncharacterized protein

Chain A:



• Molecule 1: Uncharacterized protein

Chain B:

4.2.2 Score per residue for model 2

- Molecule 1: Uncharacterized protein

Chain A:

- Molecule 1: Uncharacterized protein

Chain B:

4.2.3 Score per residue for model 3

- Molecule 1: Uncharacterized protein

Chain A:



- Molecule 1: Uncharacterized protein

Chain B:

4.2.4 Score per residue for model 4

- Molecule 1: Uncharacterized protein

Chain A:



- Molecule 1: Uncharacterized protein

Chain B:



4.2.5 Score per residue for model 5

• Molecule 1: Uncharacterized protein

Chain A: 74% 10% 16%

G274 S275 H276 M277 S278 T279 I280 T281 T290 V303 E311 R324 V328 A329 M330 E331

• Molecule 1: Uncharacterized protein

Chain B: 79% • 17%

G274 S275 H276 M277 S278 T279 I280 T281 I285 A329 M330 E331

4.2.6 Score per residue for model 6

• Molecule 1: Uncharacterized protein

Chain A: 71% 14% 16%

G274 S275 H276 M277 S278 T279 I280 T281 I285 T290 V303 E311 L315 R324 V328 A329 M330 E331

• Molecule 1: Uncharacterized protein

Chain B: 72% 10% 17%

G274 S275 H276 M277 S278 T279 I280 T281 I285 E311 L315 R324 V328 A329 M330 E331

4.2.7 Score per residue for model 7

• Molecule 1: Uncharacterized protein

Chain A: 69% 16% 16%

G274 S275 H276 M277 S278 T279 I280 T281 L287 T290 V303 L315 R324 S325 L326 K327 V328 A329 M330 E331

• Molecule 1: Uncharacterized protein

Chain B: 67% 16% 17%

G274 S275 H276 M277 S278 T279 I280 T281 L287 T290 V303 L315 R324 S325 L326 K327 V328 A329 M330 E331

4.2.8 Score per residue for model 8

- Molecule 1: Uncharacterized protein

Chain A: 67% 17% 16%

- Molecule 1: Uncharacterized protein

Chain B: 71% 12% 17%

4.2.9 Score per residue for model 9

- Molecule 1: Uncharacterized protein

Chain A: 71% 14% 16%

- Molecule 1: Uncharacterized protein

Chain B: 69% 14% 17%

4.2.10 Score per residue for model 10

- Molecule 1: Uncharacterized protein

Chain A: 74% 10% 16%



- Molecule 1: Uncharacterized protein

Chain B: 76% 7% 17%



4.2.11 Score per residue for model 11

- Molecule 1: Uncharacterized protein

Chain A: 74% 10% 16%

G274 S275 H276 M277 S278 T279 I280 T281 I285 T290 V303 E311 R324 V328 E331

- Molecule 1: Uncharacterized protein

Chain B: 72% 10% 17%

G274 S275 H276 M277 S278 T279 I280 T281 I285 T290 V303 E311 R324 V328 A329 M330 E331

4.2.12 Score per residue for model 12

- Molecule 1: Uncharacterized protein

Chain A: 76% 9% 16%

G274 S275 H276 M277 S278 T279 I280 T281 T290 V303 E311 I314 A329 M330 E331

- Molecule 1: Uncharacterized protein

Chain B: 72% 10% 17%

G274 S275 H276 M277 S278 T279 I280 T281 I285 T290 V303 R324 V328 A329 M330 E331

4.2.13 Score per residue for model 13

- Molecule 1: Uncharacterized protein

Chain A: 76% 9% 16%

G274 S275 H276 M277 S278 T279 I280 T281 T290 V303 E311 R324 V328 E331

- Molecule 1: Uncharacterized protein

Chain B: 81% • 17%

G274 S275 H276 M277 S278 T279 I280 T281 I285 M330 E331

4.2.14 Score per residue for model 14

- Molecule 1: Uncharacterized protein

Chain A: 79% 5% 16%

G274 S275 H276 M277 S278 T279 I280 T281 E311 R324 V328 E331

- Molecule 1: Uncharacterized protein

Chain B: 74% 9% 17%

G274 S275 H276 M277 S278 T279 I280 T281 I285 T290 V303 R324 V328 A329 M330 E331

4.2.15 Score per residue for model 15

- Molecule 1: Uncharacterized protein

Chain A: 76% 9% 16%

G274 S275 H276 M277 S278 T279 I280 T281 I285 R324 S325 L326 K327 V328 A329 M330 E331

- Molecule 1: Uncharacterized protein

Chain B: 67% 16% 17%

G274 S275 H276 M277 S278 T279 I280 T281 L287 T290 V303 E311 L315 R324 S325 V328 A329 M330 E331

4.2.16 Score per residue for model 16

- Molecule 1: Uncharacterized protein

Chain A: 76% 9% 16%

G274 S275 H276 M277 S278 T279 I280 T281 I285 E311 L315 R324 V328 E331

- Molecule 1: Uncharacterized protein

Chain B: 67% 16% 17%

G274 S275 H276 M277 S278 T279 I280 T281 I285 E286 L287 S288 N289 T290 V303 E311 L315 R324 V328 A329 M330 E331

4.2.17 Score per residue for model 17

- Molecule 1: Uncharacterized protein

Chain A:



- Molecule 1: Uncharacterized protein

Chain B:



4.2.18 Score per residue for model 18

- Molecule 1: Uncharacterized protein

Chain A:



- Molecule 1: Uncharacterized protein

Chain B:



5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *simulated annealing*.

Of the 50 calculated structures, 18 were deposited, based on the following criterion: *structures with the least restraint violations*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR NIH	structure solution	2.9.7
X-PLOR NIH	refinement	2.9.7

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	577
Number of shifts mapped to atoms	577
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	38%

6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	382	401	401	5±2
1	B	374	392	392	5±2
All	All	13608	14274	14274	120

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:324:ARG:O	1:A:328:VAL:HG23	0.72	1.85	15	17
1:A:285:ILE:HD11	1:B:311:GLU:OE2	0.65	1.91	1	9
1:A:311:GLU:OE2	1:B:285:ILE:HD11	0.65	1.90	12	8
1:A:287:LEU:HD13	1:A:315:LEU:HD23	0.65	1.68	9	4
1:A:329:ALA:HB1	1:B:329:ALA:HB1	0.64	1.70	2	10
1:B:287:LEU:HD13	1:B:315:LEU:HD23	0.62	1.70	9	7
1:B:290:THR:HG21	1:B:303:VAL:HA	0.60	1.72	1	14
1:B:324:ARG:O	1:B:328:VAL:HG23	0.59	1.98	8	16
1:A:290:THR:HG21	1:A:303:VAL:HA	0.58	1.74	4	11
1:A:315:LEU:HD21	1:B:315:LEU:HD21	0.55	1.78	6	6
1:A:326:LEU:HD21	1:B:325:SER:HB2	0.52	1.80	18	1
1:A:285:ILE:HD11	1:B:311:GLU:CD	0.46	2.31	8	1
1:A:311:GLU:OE1	1:B:285:ILE:HD11	0.46	2.10	11	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:329:ALA:CB	1:B:329:ALA:HB1	0.46	2.41	7	1
1:A:326:LEU:HD21	1:B:325:SER:CB	0.46	2.41	7	3
1:A:315:LEU:HD21	1:B:315:LEU:CD2	0.45	2.41	6	1
1:A:311:GLU:HA	1:A:314:ILE:HD12	0.44	1.88	10	3
1:A:325:SER:HB2	1:B:326:LEU:HD21	0.44	1.89	7	1
1:A:285:ILE:HD11	1:B:311:GLU:OE1	0.44	2.13	8	2
1:B:285:ILE:HG22	1:B:289:ASN:ND2	0.43	2.27	16	1
1:A:315:LEU:CD2	1:B:315:LEU:HD21	0.43	2.44	8	2
1:A:329:ALA:HB1	1:B:329:ALA:CB	0.41	2.43	2	1

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	49/58 (84%)	48±0 (97±1%)	1±0 (3±1%)	0±0 (0±0%)	100	100
1	B	48/58 (83%)	46±0 (97±1%)	2±0 (3±1%)	0±0 (0±0%)	100	100
All	All	1746/2088 (84%)	1693 (97%)	53 (3%)	0 (0%)	100	100

There are no Ramachandran outliers.

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	41/49 (84%)	41±0 (100±0%)	0±0 (0±0%)	100	100
1	B	40/49 (82%)	40±0 (100±0%)	0±0 (0±0%)	100	100
All	All	1458/1764 (83%)	1458 (100%)	0 (0%)	100	100

There are no protein residues with a non-rotameric sidechain to report.

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 38% for the well-defined parts and 36% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *hamp_renumbered.bmrb*

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	577
Number of shifts mapped to atoms	577
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	1

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction ± precision, *ppm*	Suggested action
$^{13}C_{\alpha}$	56	-0.33 ± 0.10	None needed (< 0.5 ppm)
$^{13}C_{\beta}$	53	0.63 ± 0.14	Should be checked
$^{13}C'$	46	-0.29 ± 0.15	None needed (< 0.5 ppm)
^{15}N	49	0.35 ± 0.34	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 38%, i.e. 524 atoms were assigned a chemical shift out of a possible 1371. 0 out of 14 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	235/481 (49%)	97/194 (50%)	92/194 (47%)	46/93 (49%)
Sidechain	286/876 (33%)	163/567 (29%)	117/265 (44%)	6/44 (14%)

Continued on next page...

Continued from previous page...

	Total	^{1}H	^{13}C	^{15}N
Aromatic	3/14 (21%)	2/8 (25%)	1/4 (25%)	0/2 (0%)
Overall	524/1371 (38%)	262/769 (34%)	210/463 (45%)	52/139 (37%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 36%, i.e. 573 atoms were assigned a chemical shift out of a possible 1594. 0 out of 14 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	259/578 (45%)	108/234 (46%)	102/232 (44%)	49/112 (44%)
Sidechain	311/988 (31%)	176/642 (27%)	129/302 (43%)	6/44 (14%)
Aromatic	3/28 (11%)	2/16 (12%)	1/8 (12%)	0/4 (0%)
Overall	573/1594 (36%)	286/892 (32%)	232/542 (43%)	55/160 (34%)

7.1.4 Statistically unusual chemical shifts (i)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, *ppm*	Expected range, *ppm*	Z-score
1	A	290	THR	HG1	5.02	0.08 – 2.19	18.4

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:


RCI
0.40
0.35
0.30
0.25
0.20
0.15
0.10
0.05
0.00
274:GLY
278:SER
280:ILE
282:ARG
284:ILE
286:GLU
288:SER
290:THR
292:ASP
294:ILE
296:GLU
298:ASN
300:GLU
302:GLU
304:PRO
306:GLN
308:ARG
310:ASP
312:ILE
314:ILE
316:ALA
318:SER
320:GLU
322:LEU
324:ARG
326:LEU
328:VAL
330:MET
Auth_seq_ID