



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:41 AM UTC

PDB ID : 9LMG / pdb_00009lmg
Title : Crystal structure of LooH
Authors : Zhang, F.; Li, Y.C.; Zhu, Y.; Wang, K.; Chang, C.Y.; Xu, Z.
Deposited on : 2025-01-19
Resolution : 1.63 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.63 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1141 (1.64-1.64)
Clashscore	190562	1171 (1.64-1.64)
Ramachandran outliers	187476	1151 (1.64-1.64)
Sidechain outliers	187428	1150 (1.64-1.64)
RSRZ outliers	180081	1141 (1.64-1.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	550	6% 93%
1	B	550	5% 92%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 9380 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LooH.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	527	Total	C	N	O	S	0	0	0
			4214	2675	734	784	21			
1	B	528	Total	C	N	O	S	0	0	0
			4221	2680	735	785	21			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	initiating methionine	UNP A0A6M5K9A4
A	-18	GLY	-	expression tag	UNP A0A6M5K9A4
A	-17	SER	-	expression tag	UNP A0A6M5K9A4
A	-16	SER	-	expression tag	UNP A0A6M5K9A4
A	-15	HIS	-	expression tag	UNP A0A6M5K9A4
A	-14	HIS	-	expression tag	UNP A0A6M5K9A4
A	-13	HIS	-	expression tag	UNP A0A6M5K9A4
A	-12	HIS	-	expression tag	UNP A0A6M5K9A4
A	-11	HIS	-	expression tag	UNP A0A6M5K9A4
A	-10	HIS	-	expression tag	UNP A0A6M5K9A4
A	-9	SER	-	expression tag	UNP A0A6M5K9A4
A	-8	SER	-	expression tag	UNP A0A6M5K9A4
A	-7	GLY	-	expression tag	UNP A0A6M5K9A4
A	-6	LEU	-	expression tag	UNP A0A6M5K9A4
A	-5	VAL	-	expression tag	UNP A0A6M5K9A4
A	-4	PRO	-	expression tag	UNP A0A6M5K9A4
A	-3	ARG	-	expression tag	UNP A0A6M5K9A4
A	-2	GLY	-	expression tag	UNP A0A6M5K9A4
A	-1	SER	-	expression tag	UNP A0A6M5K9A4
A	0	HIS	-	expression tag	UNP A0A6M5K9A4
B	-19	MET	-	initiating methionine	UNP A0A6M5K9A4
B	-18	GLY	-	expression tag	UNP A0A6M5K9A4
B	-17	SER	-	expression tag	UNP A0A6M5K9A4
B	-16	SER	-	expression tag	UNP A0A6M5K9A4
B	-15	HIS	-	expression tag	UNP A0A6M5K9A4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-14	HIS	-	expression tag	UNP A0A6M5K9A4
B	-13	HIS	-	expression tag	UNP A0A6M5K9A4
B	-12	HIS	-	expression tag	UNP A0A6M5K9A4
B	-11	HIS	-	expression tag	UNP A0A6M5K9A4
B	-10	HIS	-	expression tag	UNP A0A6M5K9A4
B	-9	SER	-	expression tag	UNP A0A6M5K9A4
B	-8	SER	-	expression tag	UNP A0A6M5K9A4
B	-7	GLY	-	expression tag	UNP A0A6M5K9A4
B	-6	LEU	-	expression tag	UNP A0A6M5K9A4
B	-5	VAL	-	expression tag	UNP A0A6M5K9A4
B	-4	PRO	-	expression tag	UNP A0A6M5K9A4
B	-3	ARG	-	expression tag	UNP A0A6M5K9A4
B	-2	GLY	-	expression tag	UNP A0A6M5K9A4
B	-1	SER	-	expression tag	UNP A0A6M5K9A4
B	0	HIS	-	expression tag	UNP A0A6M5K9A4


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	465	Total O 465 465	0	0
2	B	480	Total O 480 480	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: LooH



• Molecule 1: LooH



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	52.29Å 130.49Å 168.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	26.20 – 1.63 26.20 – 1.63	Depositor EDS
% Data completeness (in resolution range)	98.0 (26.20-1.63) 98.0 (26.20-1.63)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.25 (at 1.63Å)	Xtriage
Refinement program	REFMAC 5.8.0430	Depositor
R, R_{free}	0.192 , 0.223 0.201 , 0.229	Depositor DCC
R_{free} test set	7148 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	12.5	Xtriage
Anisotropy	0.060	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 33.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9380	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	0/4329	0.91	0/5872
1	B	0.51	0/4336	0.91	1/5882 (0.0%)
All	All	0.51	0/8665	0.91	1/11754 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	140	ASP	CA-CB-CG	5.73	118.33	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4214	0	4020	12	0
1	B	4221	0	4029	18	0
2	A	465	0	0	4	0
2	B	480	0	0	8	0
All	All	9380	0	8049	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:74:CYS:HB3	2:B:893:HOH:O	1.93	0.69
1:A:83:ARG:HE	1:A:106:HIS:HD2	1.41	0.68
1:B:83:ARG:HE	1:B:106:HIS:HD2	1.43	0.67
1:A:19:MET:CE	2:A:731:HOH:O	2.48	0.61
1:B:353:VAL:HG13	1:B:396:PHE:CE1	2.40	0.56
1:A:243:PHE:CZ	1:A:331:ARG:HG2	2.40	0.56
1:B:254:SER:OG	1:B:329:ARG:NH2	2.39	0.55
1:B:83:ARG:HE	1:B:106:HIS:CD2	2.27	0.53
1:B:113:LEU:HD21	1:B:451:ALA:HB2	1.92	0.51
1:A:83:ARG:HE	1:A:106:HIS:CD2	2.26	0.51
1:A:106:HIS:HE1	2:A:973:HOH:O	1.94	0.51
1:A:437:TYR:HB2	1:A:443:ILE:HD11	1.95	0.48
1:A:49:GLU:HG2	1:A:361:ILE:CD1	2.44	0.47
1:B:255:ALA:HA	1:B:296:TYR:O	2.17	0.45
1:A:19:MET:HE1	2:A:731:HOH:O	2.13	0.45
1:A:199:ARG:NH2	2:A:604:HOH:O	2.48	0.45
1:B:231:ARG:NH1	2:B:612:HOH:O	2.47	0.45
1:B:113:LEU:HD11	1:B:451:ALA:H	1.82	0.45
1:B:389:ASN:O	1:B:393:GLU:HG3	2.18	0.43
1:B:71:MET:CE	2:B:989:HOH:O	2.65	0.43
1:B:113:LEU:CD2	1:B:451:ALA:HB2	2.48	0.42
1:B:106:HIS:HE1	2:B:1002:HOH:O	2.02	0.42
1:A:243:PHE:CE2	1:A:331:ARG:HG2	2.55	0.42
1:A:401:ASP:HB3	1:A:430:ILE:HD12	2.02	0.42
1:A:255:ALA:HA	1:A:296:TYR:O	2.20	0.42
1:B:74:CYS:CB	2:B:893:HOH:O	2.63	0.42
1:B:437:TYR:HB2	1:B:443:ILE:HD11	2.01	0.42
1:B:71:MET:CB	2:B:989:HOH:O	2.68	0.41
1:B:71:MET:HG2	2:B:989:HOH:O	2.20	0.40
1:B:455:TYR:HB3	2:B:626:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	525/550 (96%)	514 (98%)	11 (2%)	0	100	100
1	B	526/550 (96%)	516 (98%)	10 (2%)	0	100	100
All	All	1051/1100 (96%)	1030 (98%)	21 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	439/459 (96%)	437 (100%)	2 (0%)	81	71
1	B	440/459 (96%)	438 (100%)	2 (0%)	81	71
All	All	879/918 (96%)	875 (100%)	4 (0%)	81	71

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	106	HIS
1	A	136	LYS
1	B	45	LEU
1	B	106	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	29	ASN
1	A	106	HIS
1	A	174	HIS
1	A	194	GLN
1	A	259	GLN
1	A	305	GLN
1	B	106	HIS
1	B	166	ASN
1	B	194	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	305	GLN
1	B	448	ASN
1	B	504	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	527/550 (95%)	0.40	33 (6%) 26 29	6, 13, 28, 50	0
1	B	528/550 (96%)	0.32	27 (5%) 33 38	5, 14, 27, 54	0
All	All	1055/1100 (95%)	0.36	60 (5%) 29 33	5, 14, 28, 54	0

All (60) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	453	ILE	6.7
1	B	451	ALA	5.6
1	B	455	TYR	5.5
1	B	452	GLY	5.0
1	A	42	ILE	4.8
1	A	113	LEU	4.8
1	A	452	GLY	4.5
1	B	113	LEU	4.3
1	A	455	TYR	4.2
1	B	456	GLY	3.9
1	B	530	VAL	3.7
1	A	204	THR	3.5
1	A	451	ALA	3.5
1	A	230	PHE	3.3
1	B	74	CYS	3.3
1	A	448	ASN	3.3
1	A	453	ILE	3.3
1	A	43	PRO	3.3
1	B	529	GLY	3.2
1	B	329	ARG	3.2
1	B	454	TYR	3.2
1	A	205	ASP	3.1
1	A	529	GLY	3.1
1	B	320	PRO	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	448	ASN	3.0
1	B	321	GLN	3.0
1	A	463	ARG	3.0
1	A	330	PHE	3.0
1	A	328	ILE	2.9
1	A	321	GLN	2.9
1	B	463	ARG	2.8
1	A	112	GLY	2.8
1	B	45	LEU	2.8
1	B	447	PRO	2.7
1	A	237	LYS	2.7
1	A	454	TYR	2.6
1	A	329	ARG	2.5
1	A	201	GLU	2.5
1	A	449	ASP	2.4
1	A	174	HIS	2.4
1	B	449	ASP	2.4
1	B	413	GLU	2.3
1	A	44	THR	2.3
1	A	413	GLU	2.2
1	A	203	ASN	2.2
1	A	450	ASP	2.2
1	A	71	MET	2.2
1	B	136	LYS	2.2
1	A	182	ARG	2.2
1	B	182	ARG	2.2
1	A	245	ASP	2.2
1	B	71	MET	2.2
1	A	41	GLN	2.2
1	B	289	LEU	2.1
1	B	266	ALA	2.1
1	B	102	PRO	2.1
1	B	216	GLY	2.0
1	A	3	ASP	2.0
1	A	515	ASP	2.0
1	B	301	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.