



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 12:21 PM UTC

PDB ID : 9LWV / pdb\_00009lwv  
Title : Arginine bis-prenyltransferase DciF  
Authors : Mori, T.; Taniguchi, T.; Matsuda, K.; Wakimoto, T.; Abe, I.  
Deposited on : 2025-02-17  
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	309	<div> <div>4%</div> <div>72%</div> <div>19%</div> <div>8%</div> </div>
1	B	309	<div> <div>4%</div> <div>78%</div> <div>13%</div> <div>9%</div> </div>
1	C	309	<div> <div>5%</div> <div>75%</div> <div>15%</div> <div>10%</div> </div>
1	D	309	<div> <div>22%</div> <div>70%</div> <div>11%</div> <div>19%</div> </div>

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 8594 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DciF.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	285	Total	C	N	O	S	0	0	0
			2277	1480	372	415	10			
1	B	281	Total	C	N	O	S	0	0	0
			2223	1446	356	412	9			
1	C	277	Total	C	N	O	S	0	0	0
			2224	1448	357	409	10			
1	D	251	Total	C	N	O	S	0	0	0
			1826	1176	303	341	6			

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	14	Total 14	O 14	0	0
3	B	8	Total 8	O 8	0	0
3	C	13	Total 13	O 13	0	0
3	D	3	Total 3	O 3	0	0





● Molecule 1: DciF



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	69.98Å 94.24Å 92.64Å 90.00° 92.19° 90.00°	Depositor
Resolution (Å)	47.37 – 2.60 47.37 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.6 (47.37-2.60) 99.7 (47.37-2.60)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.92 (at 2.61Å)	Xtriage
Refinement program	PHENIX 1.21.1_5286	Depositor
R, $R_{free}$	0.254 , 0.286 0.255 , 0.286	Depositor DCC
$R_{free}$ test set	2011 reflections (5.42%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	46.2	Xtriage
Anisotropy	1.008	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 59.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.000 for -h,-l,-k 0.000 for -h,l,k 0.019 for h,-k,-l	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	8594	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.40% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.12	0/2323	0.25	0/3144
1	B	0.10	0/2269	0.24	0/3076
1	C	0.10	0/2267	0.24	0/3067
1	D	0.11	0/1850	0.26	0/2506
All	All	0.11	0/8709	0.25	0/11793

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2277	0	2234	37	0
1	B	2223	0	2143	23	0
1	C	2224	0	2159	25	0
1	D	1826	0	1605	20	0
2	A	6	0	8	1	0
3	A	14	0	0	1	0
3	B	8	0	0	1	0
3	C	13	0	0	0	0
3	D	3	0	0	0	0
All	All	8594	0	8149	105	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (105) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:53:LYS:HB3	1:D:60:TYR:HB2	1.70	0.73
1:B:219:CYS:HB2	1:B:232:TYR:HB2	1.71	0.73
1:C:136:ILE:O	1:C:166:LEU:N	2.23	0.71
1:D:269:ALA:HB3	1:D:284:ASN:HB2	1.71	0.71
1:D:267:CYS:HB3	1:D:286:TYR:HB2	1.72	0.70
1:C:10:LYS:HG3	1:C:48:LEU:HB2	1.75	0.68
1:A:99:ILE:HD13	1:A:132:ILE:HD12	1.76	0.68
1:D:132:ILE:HB	1:D:170:PHE:HB2	1.79	0.64
1:B:99:ILE:HD13	1:B:132:ILE:HD12	1.79	0.63
1:C:219:CYS:HB2	1:C:232:TYR:HB2	1.79	0.63
1:A:219:CYS:HB2	1:A:232:TYR:HB2	1.81	0.63
1:A:7:SER:O	1:A:13:TYR:OH	2.17	0.63
1:B:12:TYR:O	1:B:16:ILE:HG12	2.00	0.62
1:C:132:ILE:HB	1:C:170:PHE:HB2	1.82	0.61
1:C:249:ASP:OD1	1:C:252:ARG:NH1	2.33	0.60
1:C:269:ALA:HB3	1:C:284:ASN:HB2	1.82	0.59
1:A:53:LYS:NZ	2:A:301:GOL:O1	2.35	0.59
1:D:99:ILE:HD11	1:D:172:LEU:HD11	1.85	0.59
1:B:156:ASN:HB3	1:B:203:VAL:HG22	1.86	0.58
1:A:271:GLN:NE2	3:A:401:HOH:O	2.39	0.55
1:A:10:LYS:N	1:A:40:GLU:OE2	2.35	0.54
1:B:10:LYS:HB3	1:B:48:LEU:HD22	1.88	0.54
1:D:108:ASP:OD1	1:D:111:LYS:N	2.41	0.54
1:B:269:ALA:HB3	1:B:284:ASN:HB2	1.90	0.54
1:B:114:GLU:O	1:B:134:LEU:HA	2.09	0.53
1:B:159:ASN:O	1:B:199:ARG:NH1	2.42	0.52
1:C:249:ASP:HA	1:C:252:ARG:HD2	1.92	0.52
1:C:156:ASN:HB3	1:C:203:VAL:HG13	1.90	0.52
1:A:223:SER:OG	1:A:226:ASN:OD1	2.24	0.52
1:C:114:GLU:O	1:C:134:LEU:HA	2.10	0.51
1:C:36:VAL:HG13	1:C:66:ILE:HD12	1.92	0.51
1:D:51:SER:HB3	1:D:63:ARG:HB3	1.93	0.51
1:A:153:LEU:HD12	1:A:154:ASP:H	1.76	0.51
1:C:190:GLU:N	1:C:190:GLU:OE1	2.44	0.51
1:C:99:ILE:HD11	1:C:172:LEU:HD11	1.92	0.50
1:D:92:VAL:HG13	1:D:174:LEU:HB3	1.94	0.50
1:B:79:VAL:HG11	1:B:132:ILE:HD11	1.93	0.50

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:10:LYS:HB3	1:A:48:LEU:HD22	1.94	0.49
1:A:267:CYS:HB3	1:A:286:TYR:HB2	1.94	0.49
1:A:79:VAL:HG11	1:A:132:ILE:HD11	1.95	0.49
1:D:87:GLU:HG3	1:D:94:ILE:H	1.78	0.49
1:B:238:LEU:H	1:B:238:LEU:HD23	1.78	0.48
1:A:264:ARG:HD3	1:A:289:ILE:C	2.38	0.48
1:C:143:ILE:HD13	1:C:161:LEU:HD21	1.96	0.48
1:A:121:ALA:HA	1:A:128:THR:HG23	1.95	0.47
1:C:241:PHE:CG	1:C:266:MET:HE1	2.50	0.47
1:A:91:ASP:OD1	1:A:91:ASP:N	2.40	0.47
1:A:265:GLU:HB2	1:A:288:LEU:HB3	1.97	0.47
1:B:189:GLN:O	3:B:301:HOH:O	2.20	0.47
1:B:216:SER:HB3	1:B:236:LYS:HG2	1.96	0.46
1:A:241:PHE:CG	1:A:266:MET:HE1	2.51	0.46
1:A:46:CYS:SG	1:A:66:ILE:HD11	2.55	0.46
1:A:167:HIS:HB3	1:A:183:TYR:HB2	1.97	0.46
1:A:87:GLU:HG3	1:A:94:ILE:H	1.80	0.46
1:D:75:GLN:HG3	1:D:76:LEU:H	1.81	0.46
1:B:223:SER:OG	1:B:226:ASN:OD1	2.33	0.46
1:A:269:ALA:HB3	1:A:284:ASN:HB2	1.98	0.45
1:B:121:ALA:HA	1:B:128:THR:HG23	1.99	0.45
1:A:240:ASP:O	1:A:244:TYR:HD1	2.00	0.45
1:B:111:LYS:NZ	1:B:138:ASN:O	2.47	0.45
1:B:26:PHE:CE1	1:B:123:GLN:HG2	2.51	0.45
1:A:154:ASP:HB2	1:A:222:LEU:HD12	1.98	0.45
1:B:267:CYS:HB3	1:B:286:TYR:HB2	1.99	0.45
1:A:184:PRO:HD2	1:A:220:VAL:O	2.17	0.44
1:C:70:LEU:HD21	1:C:76:LEU:HD13	1.99	0.44
1:A:10:LYS:HA	1:A:13:TYR:HD2	1.82	0.44
1:C:189:GLN:OE1	1:C:189:GLN:N	2.40	0.44
1:A:53:LYS:HB3	1:A:60:TYR:HB2	1.99	0.44
1:D:103:PHE:CE1	1:D:112:MET:HE1	2.53	0.44
1:A:12:TYR:O	1:A:16:ILE:HG12	2.18	0.43
1:A:160:LEU:HD22	1:A:200:LEU:HD23	2.00	0.43
1:B:142:LYS:HA	1:B:142:LYS:HD3	1.82	0.43
1:D:112:MET:HE3	1:D:112:MET:HB2	1.72	0.43
1:D:114:GLU:O	1:D:134:LEU:HA	2.19	0.43
1:D:129:LYS:HG3	1:D:173:SER:HA	2.00	0.43
1:C:255:HIS:O	1:C:259:GLN:HB2	2.19	0.43
1:C:27:TYR:HA	1:C:28:PRO:HA	1.84	0.43
1:C:274:GLN:HB3	1:C:280:ILE:HG23	2.01	0.43

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:10:LYS:HE3	1:A:289:ILE:HG22	2.00	0.43
1:C:188:ASN:HA	1:C:191:PHE:CD2	2.53	0.43
1:A:258:TYR:HD1	1:A:263:MET:HE3	1.84	0.43
1:B:10:LYS:HD2	1:B:48:LEU:HB2	2.00	0.43
1:D:102:PHE:CE2	1:D:134:LEU:HD11	2.54	0.43
1:C:11:ILE:HD11	1:C:37:ASN:HB2	2.01	0.42
1:A:189:GLN:OE1	1:A:189:GLN:N	2.51	0.42
1:C:87:GLU:CD	1:C:94:ILE:H	2.28	0.42
1:A:140:PRO:HB3	1:A:164:ASN:ND2	2.34	0.42
1:A:209:LEU:HA	1:A:212:LEU:HD13	2.02	0.42
1:A:116:MET:HE3	1:A:116:MET:HB2	1.87	0.42
1:A:107:PHE:HD1	1:A:142:LYS:HE3	1.84	0.42
1:D:26:PHE:HB2	1:D:121:ALA:HB1	2.02	0.41
1:A:27:TYR:HA	1:A:28:PRO:HA	1.85	0.41
1:C:103:PHE:O	1:C:107:PHE:HB2	2.20	0.41
1:B:273:LYS:HA	1:B:276:LEU:HD12	2.01	0.41
1:C:107:PHE:HD1	1:C:142:LYS:HE3	1.85	0.41
1:B:29:LEU:HD23	1:B:119:ILE:HD13	2.02	0.41
1:A:114:GLU:O	1:A:134:LEU:HA	2.20	0.41
1:A:228:ASP:OD1	1:A:229:LYS:N	2.54	0.41
1:D:35:PHE:HE1	1:D:78:VAL:HG12	1.86	0.41
1:D:132:ILE:O	1:D:170:PHE:N	2.42	0.41
1:B:212:LEU:HD11	1:B:218:ILE:HD13	2.02	0.41
1:C:120:ASP:HB2	1:C:131:LYS:HE2	2.02	0.41
1:D:268:VAL:HG12	1:D:270:VAL:HG23	2.03	0.41
1:D:12:TYR:O	1:D:16:ILE:HG12	2.21	0.40
1:B:154:ASP:OD1	1:B:156:ASN:N	2.55	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	283/309 (92%)	275 (97%)	8 (3%)	0	100	100
1	B	279/309 (90%)	272 (98%)	7 (2%)	0	100	100
1	C	269/309 (87%)	263 (98%)	6 (2%)	0	100	100
1	D	227/309 (74%)	222 (98%)	4 (2%)	1 (0%)	30	51
All	All	1058/1236 (86%)	1032 (98%)	25 (2%)	1 (0%)	48	70

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	107	PHE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	242/281 (86%)	241 (100%)	1 (0%)	84	93
1	B	233/281 (83%)	233 (100%)	0	100	100
1	C	236/281 (84%)	235 (100%)	1 (0%)	84	93
1	D	166/281 (59%)	165 (99%)	1 (1%)	78	91
All	All	877/1124 (78%)	874 (100%)	3 (0%)	86	94

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	288	LEU
1	C	171	ASP
1	D	53	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	37	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	237	ASN
1	A	255	HIS
1	B	75	GLN
1	B	95	ASN
1	B	106	ASN
1	B	284	ASN
1	C	37	ASN
1	C	38	GLN
1	C	188	ASN
1	D	150	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GOL	A	301	-	5,5,5	0.37	0	5,5,5	0.47	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	301	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	285/309 (92%)	0.52	11 (3%)	43 38	36, 54, 80, 141	0
1	B	281/309 (90%)	0.65	12 (4%)	40 34	40, 59, 85, 112	0
1	C	277/309 (89%)	0.58	15 (5%)	31 26	37, 56, 76, 120	0
1	D	251/309 (81%)	1.64	69 (27%)	1 1	56, 91, 124, 151	0
All	All	1094/1236 (88%)	0.83	107 (9%)	13 10	36, 61, 106, 151	0

All (107) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	247	VAL	4.5
1	D	241	PHE	4.4
1	D	153	LEU	4.3
1	D	109	PHE	4.3
1	D	233	PHE	4.0
1	D	13	TYR	3.9
1	C	227	ALA	3.7
1	D	106	ASN	3.7
1	D	239	ASN	3.6
1	C	124	GLU	3.6
1	D	54	ILE	3.6
1	D	90	ILE	3.5
1	D	285	LEU	3.5
1	A	277	GLY	3.5
1	D	70	LEU	3.5
1	D	163	SER	3.4
1	D	26	PHE	3.3
1	D	215	CYS	3.3
1	D	258	TYR	3.3
1	B	45	ASN	3.2
1	C	45	ASN	3.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	289	ILE	3.2
1	D	245	PHE	3.1
1	D	11	ILE	3.1
1	D	256	ALA	3.0
1	D	139	TYR	3.0
1	D	93	GLN	2.9
1	D	59	LEU	2.9
1	D	234	TYR	2.9
1	D	268	VAL	2.9
1	B	44	GLU	2.9
1	D	198	GLN	2.9
1	D	149	LEU	2.9
1	D	203	VAL	2.9
1	D	211	PHE	2.9
1	D	257	TYR	2.8
1	C	222	LEU	2.8
1	D	251	ALA	2.8
1	B	13	TYR	2.8
1	D	280	ILE	2.8
1	D	146	ALA	2.7
1	D	154	ASP	2.7
1	D	196	ILE	2.7
1	D	41	THR	2.7
1	D	107	PHE	2.7
1	C	25	ASN	2.7
1	D	270	VAL	2.7
1	A	46	CYS	2.6
1	A	250	THR	2.6
1	D	246	THR	2.6
1	D	267	CYS	2.6
1	D	111	LYS	2.6
1	D	32	PHE	2.6
1	B	12	TYR	2.6
1	D	235	LEU	2.6
1	D	114	GLU	2.6
1	D	140	PRO	2.6
1	A	64	PHE	2.5
1	D	262	PRO	2.5
1	D	269	ALA	2.5
1	B	289	ILE	2.5
1	C	4	LEU	2.5
1	D	64	PHE	2.5

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	142	LYS	2.5
1	C	125	LEU	2.5
1	D	230	VAL	2.5
1	D	266	MET	2.4
1	C	238	LEU	2.4
1	B	46	CYS	2.4
1	D	66	ILE	2.4
1	A	43	SER	2.4
1	B	112	MET	2.4
1	D	283	MET	2.4
1	A	258	TYR	2.3
1	C	46	CYS	2.3
1	D	143	ILE	2.3
1	D	180	ILE	2.3
1	D	21	PHE	2.3
1	A	289	ILE	2.3
1	A	17	HIS	2.3
1	A	5	LEU	2.3
1	D	164	ASN	2.3
1	D	242	LEU	2.2
1	D	156	ASN	2.2
1	B	163	SER	2.2
1	D	46	CYS	2.2
1	D	287	TYR	2.2
1	B	151	GLY	2.2
1	D	16	ILE	2.2
1	D	220	VAL	2.2
1	B	62	ALA	2.1
1	B	76	LEU	2.1
1	D	179	GLU	2.1
1	C	10	LYS	2.1
1	D	282	LYS	2.1
1	A	238	LEU	2.1
1	D	104	GLY	2.1
1	C	104	GLY	2.1
1	C	11	ILE	2.1
1	D	18	LYS	2.1
1	C	157	ILE	2.1
1	D	22	GLU	2.0
1	C	185	TYR	2.0
1	D	83	PHE	2.0
1	A	198	GLN	2.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	261	GLN	2.0
1	C	96	TYR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GOL	A	301	6/6	0.83	0.17	49,51,55,56	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.