wwPDB X-ray Structure Validation Summary Report (i)

Sep 12, 2023 – 03:06 AM EDT

PDB ID	:	4MCX
Title	:	P. vulgaris HIGBA structure, crystal form 2
Authors	:	Schureck, M.A.; Maehigashi, T.; Dunham, C.M.
Deposited on	:	2013-08-21
Resolution	:	2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.35.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	104	2%, 78%, 12%, 11%
1	C	104	2%, 85%, •, 12%
1	E	104	87%, •, 12%
2	B	113	4%, 73%, 10%, 18%
2	D	113	4%, 68%, 13%, 19%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	113	2% 73% 8% 19%

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 4606 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Antidote protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	93	719	451	131	134	3	0	0	0
1	C	92	710	446	130	131	3	0	0	0
1	E	92	710	446	130	131	3	0	0	0

- Molecule 2 is a protein called Killer protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	93	762	486	135	139	2	0	0	0
2	D	92	756	483	134	137	2	0	0	0
2	F	91	746	477	131	136	2	0	0	0

There are 63 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-20	MET	-	expression tag	UNP Q7A225
B	-19	GLY	-	expression tag	UNP Q7A225
B	-18	SER	-	expression tag	UNP Q7A225
B	-17	SER	-	expression tag	UNP Q7A225
B	-16	HIS	-	expression tag	UNP Q7A225
B	-15	HIS	-	expression tag	UNP Q7A225
B	-14	HIS	-	expression tag	UNP Q7A225
B	-13	HIS	-	expression tag	UNP Q7A225
B	-12	HIS	-	expression tag	UNP Q7A225
B	-11	HIS	-	expression tag	UNP Q7A225
B	-10	SER	-	expression tag	UNP Q7A225
B	-9	SER	-	expression tag	UNP Q7A225

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-8	GLY	-	expression tag	UNP Q7A225
B	-7	LEU	-	expression tag	UNP Q7A225
B	-6	VAL	-	expression tag	UNP Q7A225
B	-5	PRO	-	expression tag	UNP Q7A225
B	-4	ARG	-	expression tag	UNP Q7A225
B	-3	GLY	-	expression tag	UNP Q7A225
B	-2	SER	-	expression tag	UNP Q7A225
B	-1	HIS	-	expression tag	UNP Q7A225
B	0	MET	-	expression tag	UNP Q7A225
D	-20	MET	-	expression tag	UNP Q7A225
D	-19	GLY	-	expression tag	UNP Q7A225
D	-18	SER	-	expression tag	UNP Q7A225
D	-17	SER	-	expression tag	UNP Q7A225
D	-16	HIS	-	expression tag	UNP Q7A225
D	-15	HIS	-	expression tag	UNP Q7A225
D	-14	HIS	-	expression tag	UNP Q7A225
D	-13	HIS	-	expression tag	UNP Q7A225
D	-12	HIS	-	expression tag	UNP Q7A225
D	-11	HIS	-	expression tag	UNP Q7A225
D	-10	SER	-	expression tag	UNP Q7A225
D	-9	SER	-	expression tag	UNP Q7A225
D	-8	GLY	-	expression tag	UNP Q7A225
D	-7	LEU	-	expression tag	UNP Q7A225
D	-6	VAL	-	expression tag	UNP Q7A225
D	-5	PRO	-	expression tag	UNP Q7A225
D	-4	ARG	-	expression tag	UNP Q7A225
D	-3	GLY	-	expression tag	UNP Q7A225
D	-2	SER	-	expression tag	UNP Q7A225
D	-1	HIS	-	expression tag	UNP Q7A225
D	0	MET	-	expression tag	UNP Q7A225
F	-20	MET	-	expression tag	UNP Q7A225
F	-19	GLY	-	expression tag	UNP Q7A225
F	-18	SER	-	expression tag	UNP Q7A225
F	-17	SER	-	expression tag	UNP Q7A225
F	-16	HIS	-	expression tag	UNP Q7A225
F	-15	HIS	-	expression tag	UNP Q7A225
F	-14	HIS	-	expression tag	UNP Q7A225
F	-13	HIS	-	expression tag	UNP Q7A225
F	-12	HIS	-	expression tag	UNP Q7A225
F	-11	HIS	-	expression tag	UNP Q7A225
F	-10	SER	-	expression tag	UNP Q7A225
F	-9	SER	-	expression tag	UNP Q7A225

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	-8	GLY	-	expression tag	UNP Q7A225
F	-7	LEU	-	expression tag	UNP Q7A225
F	-6	VAL	-	expression tag	UNP Q7A225
F	-5	PRO	-	expression tag	UNP Q7A225
F	-4	ARG	-	expression tag	UNP Q7A225
F	-3	GLY	-	expression tag	UNP Q7A225
F	-2	SER	-	expression tag	UNP Q7A225
F	-1	HIS	-	expression tag	UNP Q7A225
F	0	MET	-	expression tag	UNP Q7A225

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	26	26	26	0	0
			Total	O		
3	B	44	44	44	0	0
			Total	O		
3	C	22	22	22	0	0
			Total	O		
3	D	33	33	33	0	0
			Total	O		
3	E	46	46	46	0	0
			Total	O		
3	F	32	32	32	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Antidote protein

Chain A:



- Molecule 1: Antidote protein

Chain C:



- Molecule 1: Antidote protein

Chain E:



- Molecule 2: Killer protein

Chain B:



- Molecule 2: Killer protein

Chain D:



- Molecule 2: Killer protein

Chain F:

2%
73%
8%
19%


MET
GLY
SER
SER
HIS
HIS
HIS
HIS
HIS
HIS
SER
SER
GLY
LEU
VAL
PRO
ARG
GLY
SER
HIS
M0
M1
I2
V18
R29
L56
K57
G58
D59
R60
S65
V68
L86
E89
D90
TYR
HIS

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 62	Depositor
Cell constants a, b, c, α, β, γ	120.50Å 120.50Å 64.45Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.79 – 2.10 30.79 – 2.11	Depositor EDS
% Data completeness (in resolution range)	99.8 (30.79-2.10) 92.8 (30.79-2.11)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.68 (at 2.12Å)	Xtriage
Refinement program	PHENIX 1.8.4_1496	Depositor
R, R_{free}	0.173 , 0.211 0.182 , 0.212	Depositor DCC
R_{free} test set	1406 reflections (4.56%)	wwPDB-VP
Wilson B-factor (Å^2)	31.9	Xtriage
Anisotropy	0.278	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.36 , 47.5	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.055 for h,-h-k,-l	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	4606	wwPDB-VP
Average B, all atoms (Å^2)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 40.76 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.6076e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.48	0/731	0.56	0/988
1	C	0.52	0/722	0.64	0/976
1	E	0.55	0/722	0.56	0/976
2	B	0.51	0/778	0.60	0/1048
2	D	0.57	0/772	0.60	0/1040
2	F	0.54	0/761	0.63	0/1025
All	All	0.53	0/4486	0.60	0/6053

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	719	0	730	9	0
1	C	710	0	724	5	0
1	E	710	0	724	3	0
2	B	762	0	760	6	0
2	D	756	0	755	13	0
2	F	746	0	748	5	0
3	A	26	0	0	0	0
3	B	44	0	0	0	0
3	C	22	0	0	0	0
3	D	33	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	E	46	0	0	0	0
3	F	32	0	0	0	0
All	All	4606	0	4441	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 34 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:47:ASN:ND2	1:E:23:SER:HB3	1.96	0.79
2:B:58:GLY:O	2:B:61:GLU:HG3	1.87	0.74
2:D:47:ASN:HD21	1:E:23:SER:HB3	1.58	0.66
1:C:69:ARG:NH2	2:D:45:GLU:OE1	2.29	0.66
2:B:72:TRP:CE3	2:B:88:TYR:CE2	2.88	0.61

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	91/104 (88%)	89 (98%)	2 (2%)	0	100	100
1	C	90/104 (86%)	89 (99%)	1 (1%)	0	100	100
1	E	90/104 (86%)	89 (99%)	1 (1%)	0	100	100
2	B	91/113 (80%)	86 (94%)	5 (6%)	0	100	100
2	D	90/113 (80%)	85 (94%)	5 (6%)	0	100	100
2	F	89/113 (79%)	86 (97%)	3 (3%)	0	100	100
All	All	541/651 (83%)	524 (97%)	17 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	77/87 (88%)	77 (100%)	0	100	100
1	C	76/87 (87%)	76 (100%)	0	100	100
1	E	76/87 (87%)	76 (100%)	0	100	100
2	B	81/98 (83%)	81 (100%)	0	100	100
2	D	80/98 (82%)	80 (100%)	0	100	100
2	F	79/98 (81%)	79 (100%)	0	100	100
All	All	469/555 (84%)	469 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	D	47	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	93/104 (89%)	-0.18	2 (2%)	62	66	29, 38, 65, 83	0
1	C	92/104 (88%)	-0.12	2 (2%)	62	66	27, 39, 62, 70	0
1	E	92/104 (88%)	-0.30	0	100	100	24, 34, 53, 61	0
2	B	93/113 (82%)	-0.08	4 (4%)	35	41	25, 39, 78, 92	0
2	D	92/113 (81%)	-0.04	4 (4%)	35	41	27, 40, 75, 80	0
2	F	91/113 (80%)	-0.13	2 (2%)	62	66	24, 36, 76, 94	0
All	All	553/651 (84%)	-0.14	14 (2%)	57	62	24, 38, 67, 94	0

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	57	LYS	4.8
2	F	59	ASP	4.4
1	C	25	ARG	3.3
2	B	57	LYS	3.3
1	C	29	HIS	3.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.