wwPDB NMR Structure Validation Summary Report (i)

May 8, 2024 – 06:05 PM EDT

PDB ID	:	2MGN
BMRB ID	:	19594
Title	:	Solution structure of a G-quadruplex bound to the bisquinolinium compound Phen-DC3
Authors	:	Chung, W.J.; Heddi, B.; Hamon, F.; Teulade-Fichou, M.P.; Phan, A.T.
Deposited on	:	2013-11-01

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
BMRB Restraints Analysis	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment is 32%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	24	25% 67% 8%

2 Ensemble composition and analysis (i)

This entry contains 10 models. This entry does not contain polypeptide chains, therefore identification of well-defined residues and clustering analysis are not possible. All residues are included in the validation scores.

3 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 851 atoms, of which 296 are hydrogens and 0 are deuteriums.

- Molecule 1 is a DNA chain called 5'-D(*TP*GP*AP*GP*GP*GP*TP*GP*GP*TP*GP*AP*GP*GP*GP*TP*GP*GP*GP*GP*AP*AP*GP*G)-3'.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	P	
1	A	24	783	240	270	108	142	23	0

- Molecule 2 is N2,N9-bis(1-methylquinolin-3-yl)-1,10-phenanthroline-2,9-dicarboxamide (three-letter code: PQ3) (formula: $C_{34}H_{26}N_6O_2$).



Mol	Chain	Residues	Atoms				
			Total	C	H	N	O
2	A	1	68	34	26	6	2

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: 5'-D(*TP*GP*AP*GP*GP*GP*TP*GP*GP*TP*GP*AP*GP*GP*GP*TP*GP*GP*GP*GP*AP*AP*GP*G)-3'

Chain A: 25% 67% 8%



4.2 Residue scores for the representative (author defined) model from the NMR ensemble

The representative model is number 1. Colouring as in section 4.1 above.

• Molecule 1: 5'-D(*TP*GP*AP*GP*GP*GP*TP*GP*GP*TP*GP*AP*GP*GP*GP*TP*GP*GP*GP*GP*AP*AP*GP*G)-3'

Chain A: 29% 67% •



5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *DGSA-distance geometry simulated annealing, simulated annealing, molecular dynamics.*

Of the 100 calculated structures, 10 were deposited, based on the following criterion: *structures with the lowest energy.*

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
Amber	geometry optimization	
Amber	refinement	
Amber	structure solution	
X-PLOR NIH	geometry optimization	
X-PLOR NIH	refinement	
X-PLOR NIH	structure solution	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	194
Number of shifts mapped to atoms	194
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	32%

6 Model quality (i)

6.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PQ3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.14±0.02	3±1/580 (0.5± 0.1%)	1.32±0.02	6±1/900 (0.7± 0.1%)
All	All	1.14	31/5800 (0.5%)	1.32	63/9000 (0.7%)

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	10	DT	N1-C6	-6.46	1.33	1.38	10	2
1	A	7	DT	C5-C7	5.64	1.53	1.50	5	8
1	A	16	DT	C5-C7	5.40	1.53	1.50	2	10
1	A	1	DT	C5-C7	5.36	1.53	1.50	8	9
1	A	10	DT	C5-C7	5.21	1.53	1.50	7	2

5 of 12 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	10	DT	C6-C5-C7	-6.35	119.09	122.90	4	10
1	A	7	DT	C6-C5-C7	-6.14	119.21	122.90	1	10
1	A	1	DT	C6-C5-C7	-6.03	119.28	122.90	2	10
1	A	16	DT	C6-C5-C7	-5.90	119.36	122.90	8	10
1	A	17	DG	C1'-O4'-C4'	-5.70	104.40	110.10	2	3

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	513	270	270	30±4
2	A	42	26	26	0±1
All	All	5550	2960	2960	296

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 35.

5 of 109 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:23:DG:H1'	1:A:24:DG:OP2	0.88	1.67	5	10
1:A:24:DG:N3	1:A:24:DG:H3'	0.88	1.84	3	10
1:A:6:DG:H1'	1:A:7:DT:OP2	0.83	1.74	5	7
1:A:18:DG:H4'	1:A:19:DG:OP1	0.83	1.71	2	6
1:A:22:DA:H2"	1:A:23:DG:OP1	0.78	1.76	2	7

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

There are no protein molecules in this entry.

6.3.2 Protein sidechains (i)

There are no protein molecules in this entry.

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
2	PQ3	A	25	-	48,48,48	1.03±0.01	2±0 (4±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
2	PQ3	A	25	-	66,70,70	1.15±0.04	4±2 (6±2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PQ3	A	25	-	-	0±0,16,16,16	0±0,7,7,7

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
2	A	25	PQ3	C8'-N3'	3.20	1.37	1.33	4	10
2	A	25	PQ3	C8-N3	3.17	1.37	1.33	7	10

5 of 8 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})	Models	
								Worst	Total
2	A	25	PQ3	C8'-N3'-C10'	3.78	118.27	121.59	8	10
2	A	25	PQ3	C8-N3-C10	3.65	118.39	121.59	2	10
2	A	25	PQ3	C17-C17'-N1'	2.60	122.09	118.57	5	5
2	A	25	PQ3	C17'-C17-N1	2.38	121.80	118.57	5	4
2	A	25	PQ3	C1-C2-C3	2.38	117.62	123.19	5	3

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 32% for the well-defined parts and 32% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	194
Number of shifts mapped to atoms	194
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	1

7.1.2 Chemical shift referencing (i)

No chemical shift referencing corrections were calculated (not enough data).

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 32%, i.e. 156 atoms were assigned a chemical shift out of a possible 488. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Sugar	119/288 (41%)	119/168 (71%)	0/120 (0%)	0/0 (—%)
Base	37/200 (18%)	37/128 (29%)	0/32 (0%)	0/40 (0%)
Overall	156/488 (32%)	156/296 (53%)	0/152 (0%)	0/40 (0%)

7.1.4 Statistically unusual chemical shifts (i)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules con-

taining paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, *ppm*	Expected range, *ppm*	Z-score
1	A	8	DG	H3'	4.11	4.11 – 5.76	-5.0

7.1.5 Random Coil Index (RCI) plots (i)

No *random coil index*(RCI) plot could be generated from the current chemical shift list. RCI is only applicable to proteins

8 NMR restraints analysis (i)

8.1 Conformationally restricting restraints (i)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	188
Intra-residue (|i-j|=0)	43
Sequential (|i-j|=1)	37
Medium range (|i-j|>1 and |i-j|<5)	2
Long range (|i-j|≥5)	11
Inter-chain	39
Hydrogen bond restraints	56
Disulfide bond restraints	0
Total dihedral-angle restraints	24
Number of unmapped restraints	0
Number of restraints per residue	8.5
Number of long range restraints per residue[1]	1.6

[1]Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations (i)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model (i)

Distance violations less than 0.1 Å
are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	3.6	0.2
0.2-0.5 (Medium)	3.0	0.48
>0.5 (Large)	0.9	0.68

8.2.2 Average number of dihedral-angle violations per model (i)

Dihedral-angle violations less than 1° are not included in the calculation.

Bins (°)	Average number of violations per model	Max (°)
1.0-10.0 (Small)	0.3	9.5
10.0-20.0 (Medium)	0.4	19.67
>20.0 (Large)	4.6	127.44

9 Distance violation analysis (i)

9.1 Summary of distance violations (i)

The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Restraints type	Count	%[1]	Violated[3]			Consistently Violated[4]		
			Count	%[2]	%[1]	Count	%[2]	%[1]
Intra-residue (|i-j|=0)	43	22.9	6	14.0	3.2	2	4.7	1.1
Backbone-Backbone	10	5.3	3	30.0	1.6	2	20.0	1.1
Backbone-Sidechain	31	16.5	3	9.7	1.6	0	0.0	0.0
Sidechain-Sidechain	2	1.1	0	0.0	0.0	0	0.0	0.0
Sequential (|i-j|=1)	37	19.7	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	16	8.5	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	20	10.6	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	1	0.5	0	0.0	0.0	0	0.0	0.0
Medium range (|i-j|>1 & |i-j|<5)	2	1.1	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	2	1.1	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Long range (|i-j|≥5)	11	5.9	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	2	1.1	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	7	3.7	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	2	1.1	0	0.0	0.0	0	0.0	0.0
Inter-chain	39	20.7	5	12.8	2.7	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	21	11.2	3	14.3	1.6	0	0.0	0.0
Sidechain-Sidechain	18	9.6	2	11.1	1.1	0	0.0	0.0
Hydrogen bond	56	29.8	12	21.4	6.4	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	188	100.0	23	12.2	12.2	2	1.1	1.1
Backbone-Backbone	28	14.9	3	10.7	1.6	2	7.1	1.1
Backbone-Sidechain	81	43.1	6	7.4	3.2	0	0.0	0.0
Sidechain-Sidechain	79	42.0	14	17.7	7.4	0	0.0	0.0

[1] percentage calculated with respect to the total number of distance restraints, [2] percentage calculated with respect to the number of restraints in a particular restraint category, [3] violated in at least one model, [4] violated in all the models

9.1.1 Bar chart : Distribution of distance restraints and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model (i)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å
are not included in the statistics.

Model ID	Number of violations						Mean (Å)	Max (Å)	SD[6] (Å)	Median (Å)
	IR[1]	SQ[2]	MR[3]	LR[4]	IC[5]	Total				
1	2	0	0	3	0	5	0.25	0.46	0.14	0.2
2	3	0	0	3	2	8	0.3	0.6	0.16	0.28
3	3	0	0	3	0	6	0.31	0.58	0.17	0.29
4	2	0	0	3	3	8	0.3	0.68	0.19	0.25
5	4	0	0	4	2	10	0.27	0.61	0.16	0.21
6	3	0	0	4	0	7	0.29	0.54	0.16	0.24
7	4	0	1	2	2	9	0.19	0.37	0.08	0.16
8	5	0	0	3	3	11	0.28	0.64	0.18	0.17
9	2	0	1	5	1	9	0.24	0.54	0.15	0.18
10	4	0	0	2	1	7	0.33	0.59	0.15	0.3

[1]Intra-residue restraints, [2]Sequential restraints, [3]Medium range restraints, [4]Long range restraints, [5]Inter-chain restraints, [6]Standard deviation

9.2.1 Bar graph : Distance Violation statistics for each model (i)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right

9.3 Distance violation statistics for the ensemble (i)

Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 121(IR:37, SQ:37, MR:2, LR:11, IC:34) restraints are not violated in the ensemble.

Number of violated restraints						Fraction of the ensemble	
IR[1]	SQ[2]	MR[3]	LR[4]	IC[5]	Total	Count[6]	%
1	0	0	0	1	2	1	10.0
1	0	0	0	2	3	2	20.0
1	0	0	0	1	2	3	30.0

Continued on next page...

Continued from previous page...

Number of violated restraints						Fraction of the ensemble	
IR[1]	SQ[2]	MR[3]	LR[4]	IC[5]	Total	Count[6]	%
0	0	0	0	0	0	4	40.0
0	0	0	0	0	0	5	50.0
1	0	0	0	1	2	6	60.0
0	0	0	0	0	0	7	70.0
0	0	0	0	0	0	8	80.0
0	0	0	0	0	0	9	90.0
2	0	0	0	0	2	10	100.0

[1]Intra-residue restraints, [2]Sequential restraints, [3]Medium range restraints, [4]Long range restraints, [5]Inter-chain restraints, [6] Number of models with violations

9.3.1 Bar graph : Distance violation statistics for the ensemble (i)



9.4 Most violated distance restraints in the ensemble (i)

9.4.1 Histogram : Distribution of mean distance violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models

in the ensemble



9.4.2 Table: Most violated distance restraints (i)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Models[1]	Mean (Å)	SD[1] (Å)	Median (Å)
(2,16)	1:1:A:DT:H2'	1:1:A:DT:H5'	10	0.43	0.06	0.46
(2,16)	1:1:A:DT:H2'	1:1:A:DT:H5"	10	0.43	0.06	0.46
(2,15)	1:1:A:DT:H1'	1:1:A:DT:H5'	10	0.36	0.02	0.37
(2,15)	1:1:A:DT:H1'	1:1:A:DT:H5"	10	0.36	0.02	0.37
(1,14)	1:17:A:DG:N2	1:4:A:DG:N7	9	0.16	0.04	0.15
(2,24)	1:1:A:DT:H6	1:1:A:DT:H5'	6	0.58	0.03	0.58
(2,24)	1:1:A:DT:H6	1:1:A:DT:H5"	6	0.58	0.03	0.58
(2,108)	1:13:A:DG:H1'	2:25:A:PQ3:HG'	6	0.32	0.1	0.28
(2,108)	1:13:A:DG:H1'	2:25:A:PQ3:HG	6	0.32	0.1	0.28
(1,8)	1:8:A:DG:N1	1:13:A:DG:O6	6	0.15	0.02	0.15
(1,13)	1:17:A:DG:H21	1:4:A:DG:N7	4	0.19	0.02	0.18
(2,98)	1:3:A:DA:H4'	2:25:A:PQ3:HJ	3	0.56	0.15	0.64
(1,21)	1:9:A:DG:H21	1:14:A:DG:N7	3	0.15	0.01	0.16
(1,38)	1:24:A:DG:N2	1:15:A:DG:N7	3	0.12	0.01	0.11

[1]Number of violated models, [2]Standard deviation

9.5 All violated distance restraints (i)

9.5.1 Histogram : Distribution of distance violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



9.5.2 Table : All distance violations (i)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(2,98)	1:3:A:DA:H4'	2:25:A:PQ3:HJ	4	0.68
(2,98)	1:3:A:DA:H4'	2:25:A:PQ3:HJ	8	0.64
(2,24)	1:1:A:DT:H6	1:1:A:DT:H5'	5	0.61
(2,24)	1:1:A:DT:H6	1:1:A:DT:H5"	5	0.61
(2,24)	1:1:A:DT:H6	1:1:A:DT:H5'	2	0.6
(2,24)	1:1:A:DT:H6	1:1:A:DT:H5"	2	0.6
(2,24)	1:1:A:DT:H6	1:1:A:DT:H5'	10	0.59
(2,24)	1:1:A:DT:H6	1:1:A:DT:H5"	10	0.59

Continued on next page...

Continued from previous page...

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(2,24)	1:1:A:DT:H6	1:1:A:DT:H5'	3	0.58
(2,24)	1:1:A:DT:H6	1:1:A:DT:H5"	3	0.58
(2,24)	1:1:A:DT:H6	1:1:A:DT:H5'	8	0.55
(2,24)	1:1:A:DT:H6	1:1:A:DT:H5"	8	0.55
(2,108)	1:13:A:DG:H1'	2:25:A:PQ3:HG'	9	0.54
(2,24)	1:1:A:DT:H6	1:1:A:DT:H5'	6	0.54
(2,24)	1:1:A:DT:H6	1:1:A:DT:H5"	6	0.54
(2,16)	1:1:A:DT:H2'	1:1:A:DT:H5'	6	0.48

10 Dihedral-angle violation analysis (i)

10.1 Summary of dihedral-angle violations (i)

The following table provides the summary of dihedral-angle violations in different dihedral-angle types. Violations less than 1° are not included in the calculation.

Angle type	Count	%[1]	Violated[3]			Consistently Violated[4]		
			Count	%[2]	%[1]	Count	%[2]	%[1]
CHI	20	83.3	7	35.0	29.2	1	5.0	4.2
UNNAMED	4	16.7	4	100.0	16.7	2	50.0	8.3
Total	24	100.0	11	45.8	45.8	3	12.5	12.5

[1] percentage calculated with respect to total number of dihedral-angle restraints, [2] percentage calculated with respect to number of restraints in a particular dihedral-angle type, [3] violated in at least one model, [4] violated in all the models

10.1.1 Bar chart : Distribution of dihedral-angles and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories

10.2 Dihedral-angle violation statistics for each model (i)

The following table provides the dihedral-angle violation statistics for each model in the ensemble. Violations less than 1° are not included in the statistics.

Model ID	Number of violations			Mean (°)	Max (°)	SD (°)	Median (°)
	CHI	UNNAMED	Total				
1	2	4	6	75.69	125.48	40.52	83.73
2	2	3	5	73.38	123.57	47.25	103.17
3	1	4	5	83.53	113.45	31.42	104.46
4	2	4	6	66.25	111.68	31.77	56.82
5	1	3	4	84.68	118.05	41.98	103.97
6	2	4	6	75.42	120.06	38.35	80.66
7	3	2	5	61.66	127.44	45.96	34.59
8	1	3	4	84.5	106.43	32.95	102.02
9	3	3	6	54.19	108.61	41.85	47.1
10	2	4	6	60.22	106.13	45.46	64.73

10.2.1 Bar graph : Dihedral violation statistics for each model (i)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right

10.3 Dihedral-angle violation statistics for the ensemble (i)

Violation analysis may find that some restraints are violated in very few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of ensemble.

Number of violated restraints			Fraction of the ensemble	
CHI	UNNAMED	Total	Count[1]	%
4	0	4	1	10.0
1	0	1	2	20.0
1	0	1	3	30.0
0	0	0	4	40.0
0	0	0	5	50.0
0	1	1	6	60.0
0	0	0	7	70.0
0	1	1	8	80.0
0	0	0	9	90.0
1	2	3	10	100.0

[1] Number of models with violations

10.3.1 Bar graph : Dihedral-angle Violation statistics for the ensemble (i)



10.4 Most violated dihedral-angle restraints in the ensemble (i)

10.4.1 Histogram : Distribution of mean dihedral-angle violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble



10.4.2 Table: Most violated dihedral-angle restraints (i)

The following table provides the mean and the standard deviation of the violation for each restraint sorted by number of violated models and the mean value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	Models[1]	Mean	SD[2]	Median
(1,7)	1:7:A:DT:O4'	1:7:A:DT:C1'	1:7:A:DT:N1	1:7:A:DT:C6	10	113.29	8.43	112.56
(1,16)	1:16:A:DT:O4'	1:16:A:DT:C1'	1:16:A:DT:N1	1:16:A:DT:C6	10	106.65	2.06	106.63
(1,22)	1:22:A:DA:O4'	1:22:A:DA:C1'	1:22:A:DA:N9	1:22:A:DA:C4	10	28.96	7.12	29.8
(1,1)	1:1:A:DT:O4'	1:1:A:DT:C1'	1:1:A:DT:N1	1:1:A:DT:C6	8	94.62	18.21	103.82
(1,10)	1:10:A:DT:O4'	1:10:A:DT:C1'	1:10:A:DT:N1	1:10:A:DT:C6	6	60.07	34.12	55.18
(1,23)	1:23:A:DG:O4'	1:23:A:DG:C1'	1:23:A:DG:N9	1:23:A:DG:C4	3	17.56	4.25	20.09
(1,11)	1:11:A:DG:O4'	1:11:A:DG:C1'	1:11:A:DG:N9	1:11:A:DG:C4	2	15.34	4.33	15.34

[1] Number of violated models, [2]Standard deviation, All angle values are in degree (°)

10.5 All violated dihedral-angle restraints (i)

10.5.1 Histogram : Distribution of violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



10.5.2 Table: All violated dihedral-angle restraints (i)

The following table provides the list of violations for the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	Model ID	Violation (°)
(1,7)	1:7:A:DT:O4'	1:7:A:DT:C1'	1:7:A:DT:N1	1:7:A:DT:C6	7	127.44
(1,10)	1:10:A:DT:O4'	1:10:A:DT:C1'	1:10:A:DT:N1	1:10:A:DT:C6	1	125.48
(1,7)	1:7:A:DT:O4'	1:7:A:DT:C1'	1:7:A:DT:N1	1:7:A:DT:C6	2	123.57
(1,7)	1:7:A:DT:O4'	1:7:A:DT:C1'	1:7:A:DT:N1	1:7:A:DT:C6	6	120.06
(1,7)	1:7:A:DT:O4'	1:7:A:DT:C1'	1:7:A:DT:N1	1:7:A:DT:C6	5	118.05
(1,7)	1:7:A:DT:O4'	1:7:A:DT:C1'	1:7:A:DT:N1	1:7:A:DT:C6	3	113.45
(1,7)	1:7:A:DT:O4'	1:7:A:DT:C1'	1:7:A:DT:N1	1:7:A:DT:C6	4	111.68
(1,16)	1:16:A:DT:O4'	1:16:A:DT:C1'	1:16:A:DT:N1	1:16:A:DT:C6	1	111.48
(1,1)	1:1:A:DT:O4'	1:1:A:DT:C1'	1:1:A:DT:N1	1:1:A:DT:C6	6	110.08
(1,7)	1:7:A:DT:O4'	1:7:A:DT:C1'	1:7:A:DT:N1	1:7:A:DT:C6	9	108.61