



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 2, 2025 – 02:16 PM EDT

PDB ID : 9MSO / pdb_00009mso
Title : Crystal structure of MPXV A35R in complex with neutralizing antibody EV35-6
Authors : Yuan, M.; Zhu, X.; Wilson, I.A.
Deposited on : 2025-01-10
Resolution : 3.18 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0rc1
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.006 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.45.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.18 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1851 (3.20-3.16)
Clashscore	180529	1999 (3.20-3.16)
Ramachandran outliers	177936	1961 (3.20-3.16)
Sidechain outliers	177891	1960 (3.20-3.16)
RSRZ outliers	164620	1852 (3.20-3.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	228	
1	H	228	
2	B	214	
2	L	214	
3	E	100	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	100	<div><div><div></div><div></div><div></div></div><div>6%68%10%22%</div></div>
3	G	100	<div><div><div></div><div></div><div></div></div><div>%75%8%17%</div></div>
3	I	100	<div><div><div></div><div></div><div></div></div><div>3%77%•21%</div></div>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 9073 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called EV35-6 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	H	221	Total	C	N	O	S	0	0	0
			1678	1076	279	319	4			
1	A	217	Total	C	N	O	S	0	0	0
			1653	1062	274	313	4			

- Molecule 2 is a protein called EV35-6 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	211	Total	C	N	O	S	0	0	0
			1623	1014	278	327	4			
2	B	211	Total	C	N	O	S	0	0	0
			1623	1014	278	327	4			

- Molecule 3 is a protein called Protein OPG161.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	G	83	Total	C	N	O	S	0	0	0
			650	403	103	140	4			
3	I	79	Total	C	N	O	S	0	0	0
			621	387	97	133	4			
3	E	78	Total	C	N	O	S	0	0	0
			615	384	96	131	4			
3	F	78	Total	C	N	O	S	0	0	0
			610	381	96	129	4			

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	182	GLY	-	expression tag	UNP A0A7H0DND2
G	183	SER	-	expression tag	UNP A0A7H0DND2
G	184	HIS	-	expression tag	UNP A0A7H0DND2
G	185	HIS	-	expression tag	UNP A0A7H0DND2

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	186	HIS	-	expression tag	UNP A0A7H0DND2
G	187	HIS	-	expression tag	UNP A0A7H0DND2
G	188	HIS	-	expression tag	UNP A0A7H0DND2
G	189	HIS	-	expression tag	UNP A0A7H0DND2
I	182	GLY	-	expression tag	UNP A0A7H0DND2
I	183	SER	-	expression tag	UNP A0A7H0DND2
I	184	HIS	-	expression tag	UNP A0A7H0DND2
I	185	HIS	-	expression tag	UNP A0A7H0DND2
I	186	HIS	-	expression tag	UNP A0A7H0DND2
I	187	HIS	-	expression tag	UNP A0A7H0DND2
I	188	HIS	-	expression tag	UNP A0A7H0DND2
I	189	HIS	-	expression tag	UNP A0A7H0DND2
E	182	GLY	-	expression tag	UNP A0A7H0DND2
E	183	SER	-	expression tag	UNP A0A7H0DND2
E	184	HIS	-	expression tag	UNP A0A7H0DND2
E	185	HIS	-	expression tag	UNP A0A7H0DND2
E	186	HIS	-	expression tag	UNP A0A7H0DND2
E	187	HIS	-	expression tag	UNP A0A7H0DND2
E	188	HIS	-	expression tag	UNP A0A7H0DND2
E	189	HIS	-	expression tag	UNP A0A7H0DND2
F	182	GLY	-	expression tag	UNP A0A7H0DND2
F	183	SER	-	expression tag	UNP A0A7H0DND2
F	184	HIS	-	expression tag	UNP A0A7H0DND2
F	185	HIS	-	expression tag	UNP A0A7H0DND2
F	186	HIS	-	expression tag	UNP A0A7H0DND2
F	187	HIS	-	expression tag	UNP A0A7H0DND2
F	188	HIS	-	expression tag	UNP A0A7H0DND2
F	189	HIS	-	expression tag	UNP A0A7H0DND2


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: EV35-6 heavy chain




- Molecule 1: EV35-6 heavy chain




- Molecule 2: EV35-6 light chain




- Molecule 2: EV35-6 light chain




- Molecule 3: Protein OPG161




• Molecule 3: Protein OPG161



• Molecule 3: Protein OPG161



• Molecule 3: Protein OPG161



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	72.50Å 165.50Å 118.45Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.90 – 3.18 43.90 – 3.18	Depositor EDS
% Data completeness (in resolution range)	87.2 (43.90-3.18) 87.3 (43.90-3.18)	Depositor EDS
R_{merge}	0.39	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.49 (at 3.19Å)	Xtriage
Refinement program	PHENIX (1.21.2_5419: ???)	Depositor
R, R_{free}	0.240 , 0.251 0.239 , 0.249	Depositor DCC
R_{free} test set	1131 reflections (4.58%)	wwPDB-VP
Wilson B-factor (Å ²)	69.4	Xtriage
Anisotropy	0.215	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 17.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	9073	wwPDB-VP
Average B, all atoms (Å ²)	69.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.08	0/1704	0.28	0/2333
1	H	0.08	0/1729	0.28	0/2365
2	B	0.08	0/1659	0.29	0/2256
2	L	0.08	0/1659	0.26	0/2256
3	E	0.08	0/628	0.26	0/853
3	F	0.15	0/623	0.29	0/846
3	G	0.10	0/663	0.30	0/898
3	I	0.09	0/634	0.29	0/861
All	All	0.09	0/9299	0.28	0/12668

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1653	0	1615	18	0
1	H	1678	0	1641	16	0
2	B	1623	0	1574	14	0
2	L	1623	0	1574	8	0
3	E	615	0	560	3	0
3	F	610	0	557	5	0
3	G	650	0	591	5	0
3	I	621	0	565	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	9073	0	8677	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:I:163:THR:HG22	3:I:164:SER:H	1.42	0.82
1:A:100(C):GLY:HA2	3:F:149:GLU:HB3	1.77	0.67
2:L:29:VAL:HG11	2:L:90:GLN:HB2	1.78	0.66
2:L:120:PRO:HD3	2:L:132:VAL:HG22	1.78	0.66
1:H:18:LEU:HB3	1:H:82:LEU:HB3	1.80	0.63
1:A:18:LEU:HD12	1:A:109:VAL:HG11	1.83	0.61
1:H:2:VAL:HG22	1:H:27:ALA:HB2	1.86	0.58
1:H:40:PRO:HB2	1:H:43:LYS:HE2	1.87	0.57
1:A:2:VAL:HG22	1:A:27:ALA:HB2	1.86	0.56
2:B:120:PRO:HD3	2:B:132:VAL:HG22	1.86	0.56
2:B:37:GLN:HB2	2:B:47:LEU:HD11	1.90	0.54
1:H:39:GLN:HB2	1:H:45:LEU:HD23	1.89	0.54
1:H:38:ARG:HB3	1:H:48:ILE:HD11	1.88	0.54
3:G:97:LYS:HG3	3:G:98:GLU:HG2	1.91	0.52
3:I:163:THR:HG22	3:I:164:SER:N	2.20	0.52
1:H:18:LEU:HD12	1:H:109:VAL:HG11	1.90	0.52
3:G:134:PRO:HD3	3:G:179:PHE:O	2.10	0.52
1:A:35:THR:HG22	1:A:50:GLU:HB3	1.91	0.52
1:A:94:ARG:HG2	1:A:102:PRO:HD2	1.91	0.52
1:A:39:GLN:HB2	1:A:45:LEU:HD23	1.92	0.51
1:H:35:THR:HG22	1:H:50:GLU:HB3	1.94	0.50
1:H:33:TYR:CZ	1:H:52:HIS:HB2	2.47	0.50
1:H:8:GLY:HA3	1:H:20:LEU:HD23	1.94	0.49
3:F:134:PRO:HD3	3:F:179:PHE:O	2.12	0.48
3:G:145:ILE:O	3:G:149:GLU:HB2	2.14	0.47
1:A:100(D):LYS:HG2	3:E:116:TYR:HE1	1.79	0.47
1:A:18:LEU:HB3	1:A:82:LEU:HB3	1.96	0.47
1:A:1:GLN:HG2	1:A:2:VAL:H	1.80	0.47
3:F:131:SER:HB2	3:F:180:CYS:HB3	1.97	0.47
1:H:17:THR:HG22	1:H:82(A):THR:HA	1.97	0.46
2:B:119:PRO:HB3	2:B:209:PHE:CE1	2.51	0.46
1:A:119:PRO:HB3	1:A:145:TYR:HB3	1.97	0.46
1:A:34:TRP:CZ3	1:A:94:ARG:HB3	2.51	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:38:ARG:HB3	1:A:48:ILE:HD11	1.97	0.45
2:L:21:LEU:HD23	2:L:102:THR:HB	1.98	0.45
2:L:33:LEU:HD13	2:L:71:PHE:CD1	2.51	0.45
2:B:33:LEU:HD13	2:B:71:PHE:CD1	2.51	0.45
1:H:95:VAL:HG12	1:H:100(F):PRO:HB2	1.98	0.45
2:B:120:PRO:HG3	2:B:130:ALA:HB1	1.99	0.45
2:B:142:ARG:HH12	2:B:163:VAL:HG21	1.82	0.45
1:A:17:THR:HG22	1:A:82(A):THR:HA	1.99	0.44
1:A:6:GLN:HG2	1:A:92:CYS:SG	2.58	0.44
1:A:47:TRP:CZ3	2:B:95:GLY:HA2	2.52	0.44
2:L:37:GLN:HB2	2:L:47:LEU:HD11	1.99	0.44
1:A:100(D):LYS:HG2	3:E:116:TYR:CE1	2.53	0.43
2:B:33:LEU:HD11	2:B:88:CYS:SG	2.59	0.43
1:A:200:HIS:CD2	1:A:202:PRO:HD2	2.54	0.43
1:H:34:TRP:CZ3	1:H:94:ARG:HB2	2.53	0.43
2:L:33:LEU:HD11	2:L:88:CYS:SG	2.59	0.43
2:B:21:LEU:HD23	2:B:102:THR:HB	2.01	0.43
3:F:119:PHE:CE1	3:F:156:GLY:HA2	2.53	0.43
1:H:6:GLN:HG2	1:H:92:CYS:SG	2.58	0.42
1:H:100(E):LYS:HA	1:H:100(F):PRO:HD3	1.92	0.42
2:B:2:ILE:HB	2:B:90:GLN:HE22	1.84	0.42
2:L:69:THR:HG22	2:B:70:ASP:HB2	2.02	0.42
1:A:24:VAL:HG11	1:A:34:TRP:CH2	2.55	0.41
2:B:134:CYS:HB2	2:B:148:TRP:CH2	2.56	0.41
3:F:140:LEU:HD11	3:F:161:LYS:HA	2.01	0.41
2:L:56:THR:HG21	3:G:173:GLN:O	2.19	0.41
3:E:131:SER:HB2	3:E:180:CYS:HB3	2.02	0.41
1:H:100(B):TRP:HZ3	3:G:112:LEU:HD22	1.86	0.41
2:B:140:TYR:CG	2:B:141:PRO:HA	2.56	0.41
2:B:128:GLY:HA2	2:B:183:LYS:HB2	2.02	0.40
1:H:200:HIS:CD2	1:H:202:PRO:HD2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	213/228 (93%)	207 (97%)	5 (2%)	1 (0%)	25	58
1	H	217/228 (95%)	213 (98%)	4 (2%)	0	100	100
2	B	209/214 (98%)	202 (97%)	7 (3%)	0	100	100
2	L	209/214 (98%)	203 (97%)	6 (3%)	0	100	100
3	E	74/100 (74%)	71 (96%)	3 (4%)	0	100	100
3	F	74/100 (74%)	71 (96%)	3 (4%)	0	100	100
3	G	79/100 (79%)	72 (91%)	7 (9%)	0	100	100
3	I	75/100 (75%)	70 (93%)	5 (7%)	0	100	100
All	All	1150/1284 (90%)	1109 (96%)	40 (4%)	1 (0%)	48	78

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	100(D)	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	184/193 (95%)	183 (100%)	1 (0%)	86	93
1	H	187/193 (97%)	187 (100%)	0	100	100
2	B	183/185 (99%)	183 (100%)	0	100	100
2	L	183/185 (99%)	183 (100%)	0	100	100
3	E	70/91 (77%)	70 (100%)	0	100	100
3	F	69/91 (76%)	68 (99%)	1 (1%)	62	81
3	G	74/91 (81%)	74 (100%)	0	100	100
3	I	71/91 (78%)	71 (100%)	0	100	100
All	All	1021/1120 (91%)	1019 (100%)	2 (0%)	92	97

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	GLN
3	F	150	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
3	G	173	GLN
1	A	52	HIS
2	B	199	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	217/228 (95%)	0.21	4 (1%) 67 52	52, 65, 97, 108	0
1	H	221/228 (96%)	0.17	2 (0%) 81 67	49, 63, 86, 94	0
2	B	211/214 (98%)	0.21	4 (1%) 66 50	49, 71, 92, 104	0
2	L	211/214 (98%)	0.23	4 (1%) 66 50	48, 61, 96, 109	0
3	E	78/100 (78%)	0.33	0 100 100	63, 71, 93, 116	0
3	F	78/100 (78%)	0.41	6 (7%) 21 14	64, 75, 92, 108	0
3	G	83/100 (83%)	0.37	1 (1%) 76 60	53, 64, 98, 119	0
3	I	79/100 (79%)	0.34	3 (3%) 44 29	54, 65, 100, 124	0
All	All	1178/1284 (91%)	0.25	24 (2%) 64 48	48, 66, 95, 124	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	94	TRP	3.7
3	F	163	THR	3.0
1	A	100(E)	LYS	2.9
2	L	194	CYS	2.9
2	L	208	SER	2.9
3	F	162	THR	2.9
1	A	100(D)	LYS	2.8
2	B	56	THR	2.6
3	I	98	GLU	2.6
1	A	213	PRO	2.6
1	A	207	VAL	2.6
2	L	94	TRP	2.5
3	I	171	VAL	2.4
3	I	164	SER	2.3
3	F	182	GLY	2.2
3	F	171	VAL	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	H	100(D)	LYS	2.1
3	F	99	SER	2.1
3	F	160	THR	2.1
1	H	214	LYS	2.1
2	B	88	CYS	2.1
2	L	206	THR	2.1
3	G	162	THR	2.1
2	B	187	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.