wwPDB X-ray Structure Validation Summary Report (i)

Aug 10, 2020 – 05:51 AM BST

PDB ID	:	1MUU
Title	:	2.0 A crystal structure of GDP-mannose dehydrogenase
Authors	:	Snook, C.F.; Tipton, P.A.; Beamer, L.J.
Deposited on	:	2002-09-24
Resolution	:	2.02 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.13.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.02 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	11643 (2.04-2.00)
Ramachandran outliers	138981	11493 (2.04-2.00)
Sidechain outliers	138945	11492 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	436	85% 14% •
1	B	436	85% 13% •
1	C	436	83% 14% •
1	D	436	85% 13% •
2	E	2	50% 50%

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 14082 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GDP-mannose 6-dehydrogenase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	436	3193	2020	555	600	9	9	0	0	0
1	B	436	3210	2027	556	609	9	9	0	0	0
1	C	436	3227	2039	558	612	9	9	0	0	0
1	D	436	3226	2038	558	612	9	9	0	0	0

- Molecule 2 is an oligosaccharide called beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	E	2	23	12	11	0	0	0

- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: $C_{21}H_{27}N_7O_{14}P_2$).

NAD



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	35	15	5	13	2	0	0
			Total	C	N	O	P		
3	B	1	35	15	5	13	2	0	0
			Total	C	N	O	P		
3	C	1	35	15	5	13	2	0	0
			Total	C	N	O	P		
3	D	1	35	15	5	13	2	0	0

- Molecule 4 is GUANOSINE 5'-(TRIHYDROGEN DIPHOSPHATE), P'-D-MANNOPYRANOSYL ESTER (three-letter code: GDX) (formula: $C_{16}H_{23}N_5O_{17}P_2$).

GDX



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
4	B	1	40	16	5	17	2	0	0
			Total	C	N	O	P		
4	B	1	40	16	5	17	2	0	0
			Total	C	N	O	P		
4	D	1	40	16	5	17	2	0	0
			Total	C	N	O	P		
4	D	1	40	16	5	17	2	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	183	183	183	0	0
			Total	O		
5	B	222	222	222	0	0
			Total	O		
5	C	241	241	241	0	0
			Total	O		
5	D	257	257	257	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: GDP-mannose 6-dehydrogenase

Chain A:



- Molecule 1: GDP-mannose 6-dehydrogenase

Chain B:



- Molecule 1: GDP-mannose 6-dehydrogenase

Chain C:



- Molecule 1: GDP-mannose 6-dehydrogenase

Chain D:



- Molecule 2: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose

Chain E:

4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, α, β, γ	82.50Å 82.50Å 310.17Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.02	Depositor
% Data completeness (in resolution range)	99.1 (50.00-2.02)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	REFMAC 5.0	Depositor
R, R_{free}	0.209 , 0.234	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	14082	wwPDB-VP
Average B, all atoms (Å^2)	20.0	wwPDB-VP

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GDX, GLC, FRU, NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.49	1/3238 (0.0%)	0.77	8/4391 (0.2%)
1	B	0.50	0/3254	0.77	9/4407 (0.2%)
1	C	0.51	0/3272	0.79	7/4430 (0.2%)
1	D	0.52	0/3272	0.79	9/4432 (0.2%)
All	All	0.50	1/13036 (0.0%)	0.78	33/17660 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	289	MSE	SE-CE	-5.14	1.65	1.95

The worst 5 of 33 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	76	ASP	CB-CG-OD2	6.68	124.31	118.30
1	A	284	ASP	CB-CG-OD2	6.46	124.11	118.30
1	D	306	ASP	CB-CG-OD2	6.17	123.85	118.30
1	C	284	ASP	CB-CG-OD2	6.08	123.78	118.30
1	D	385	ASP	CB-CG-OD2	5.98	123.68	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3193	0	3111	53	0
1	B	3210	0	3155	44	0
1	C	3227	0	3182	67	1
1	D	3226	0	3173	47	1
2	E	23	0	21	2	0
3	A	35	0	19	1	0
3	B	35	0	19	0	0
3	C	35	0	19	1	0
3	D	35	0	19	0	0
4	B	80	0	32	0	0
4	D	80	0	32	0	0
5	A	183	0	0	2	2
5	B	222	0	0	3	6
5	C	241	0	0	9	3
5	D	257	0	0	1	5
All	All	14082	0	12782	180	9

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 180 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:315:LYS:NZ	5:C:1232:HOH:O	1.81	1.11
1:C:392:ASP:OD1	2:E:1:GLC:H62	1.52	1.07
1:A:289:MSE:HE3	1:B:293:LEU:CD2	1.91	1.01
1:A:116:ARG:NH2	1:A:147:GLY:O	1.99	0.94
1:A:1:MSE:HE1	1:A:186:LEU:HD21	1.52	0.90

The worst 5 of 9 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:1163:HOH:O	5:C:1045:HOH:O[3_464]	1.88	0.32
5:B:1090:HOH:O	5:D:1157:HOH:O[1_565]	1.92	0.28
1:D:104:ARG:NE	5:B:1100:HOH:O[3_554]	2.03	0.17
5:B:1090:HOH:O	5:D:1184:HOH:O[1_565]	2.03	0.17
5:B:1091:HOH:O	5:D:1044:HOH:O[1_565]	2.06	0.14

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	434/436 (100%)	416 (96%)	18 (4%)	0	100	100
1	B	434/436 (100%)	422 (97%)	12 (3%)	0	100	100
1	C	434/436 (100%)	422 (97%)	12 (3%)	0	100	100
1	D	434/436 (100%)	422 (97%)	12 (3%)	0	100	100
All	All	1736/1744 (100%)	1682 (97%)	54 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	327/361 (91%)	313 (96%)	14 (4%)	29	25
1	B	336/361 (93%)	318 (95%)	18 (5%)	22	17
1	C	340/361 (94%)	325 (96%)	15 (4%)	28	24
1	D	340/361 (94%)	321 (94%)	19 (6%)	21	16
All	All	1343/1444 (93%)	1277 (95%)	66 (5%)	25	20

5 of 66 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	395	VAL
1	C	114	SER
1	D	396	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	396	LEU
1	C	96	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	182	GLN
1	D	365	ASN
1	D	247	GLN
1	B	182	GLN
1	D	362	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	GLC	E	1	2	11,11,12	0.56	0	15,15,17	1.36	1 (6%)
2	FRU	E	2	2	11,12,12	0.64	0	10,18,18	0.59	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	E	1	2	-	0/2/19/22	0/1/1/1
2	FRU	E	2	2	-	2/5/24/24	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	E	1	GLC	C1-O5-C5	4.13	117.79	112.19

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	E	2	FRU	O1-C1-C2-O2
2	E	2	FRU	O1-C1-C2-C3

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	1	GLC	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

Oligosaccharide Chain E



5.6 Ligand geometry (i)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	NAD	D	1004	-	33,38,48	1.12	2 (6%)	37,58,73	1.49	5 (13%)
4	GDX	B	1005	-	33,43,43	4.51	14 (42%)	45,67,67	2.09	10 (22%)
4	GDX	D	1008	-	33,43,43	4.33	15 (45%)	45,67,67	1.91	10 (22%)
3	NAD	A	1001	-	33,38,48	1.08	2 (6%)	37,58,73	1.48	4 (10%)
4	GDX	D	1007	-	33,43,43	4.45	14 (42%)	45,67,67	2.03	9 (20%)
3	NAD	B	1002	-	33,38,48	1.13	2 (6%)	37,58,73	1.44	4 (10%)
3	NAD	C	1003	-	33,38,48	1.11	2 (6%)	37,58,73	1.41	4 (10%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	GDX	B	1006	-	33,43,43	4.37	12 (36%)	45,67,67	1.97	10 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAD	D	1004	-	-	3/18/51/62	0/4/4/5
4	GDX	B	1005	-	-	2/17/61/61	0/4/4/4
4	GDX	D	1008	-	-	1/17/61/61	0/4/4/4
3	NAD	A	1001	-	-	2/18/51/62	0/4/4/5
4	GDX	D	1007	-	-	2/17/61/61	0/4/4/4
3	NAD	B	1002	-	-	2/18/51/62	0/4/4/5
3	NAD	C	1003	-	-	1/18/51/62	0/4/4/5
4	GDX	B	1006	-	-	2/17/61/61	0/4/4/4

The worst 5 of 63 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	1005	GDX	C2D-C1D	-11.42	1.36	1.53
4	D	1008	GDX	C2D-C1D	-11.35	1.36	1.53
4	D	1007	GDX	C2D-C1D	-11.24	1.36	1.53
4	B	1006	GDX	C2D-C1D	-11.09	1.36	1.53
4	D	1007	GDX	C4'-C5'	-8.35	1.35	1.53

The worst 5 of 56 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	D	1004	NAD	N3A-C2A-N1A	-5.91	119.43	128.68
4	B	1005	GDX	C3D-C2D-C1D	5.82	109.74	100.98
3	C	1003	NAD	N3A-C2A-N1A	-5.70	119.77	128.68
3	B	1002	NAD	N3A-C2A-N1A	-5.58	119.96	128.68
3	A	1001	NAD	N3A-C2A-N1A	-5.55	120.00	128.68

There are no chirality outliers.

5 of 15 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	1005	GDX	C1'-O1B-PB-O3A

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	D	1008	GDX	C1'-O1B-PB-O3A
4	D	1007	GDX	C1'-O1B-PB-O3A
3	B	1002	NAD	O4B-C4B-C5B-O5B
3	D	1004	NAD	PN-O3-PA-O1A

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1001	NAD	1	0
3	C	1003	NAD	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand NAD D 1004



Bond lengths



Bond angles



Torsions



Rings

Ligand GDX B 1005



Ligand GDX D 1008



Ligand NAD A 1001



Bond lengths



Bond angles



Torsions



Rings

Ligand GDX D 1007



Ligand NAD B 1002



Bond lengths



Bond angles



Torsions



Rings

Ligand NAD C 1003



Bond lengths



Bond angles



Torsions



Rings

Ligand GDX B 1006



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.