wwPDB X-ray Structure Validation Summary Report (i)

Oct 27, 2023 – 05:49 AM EDT

PDB ID	:	3N2X
Title	:	Crystal structure of YagE, a prophage protein belonging to the dihydrodipicolinic acid synthase family from E. coli K12 in complex with pyruvate
Authors	:	Bhaskar, V.; Kumar, P.M.; Manicka, S.; Krishnaswamy, S.
Deposited on	:	2010-05-19
Resolution	:	2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	298	90% 7% • •
1	B	298	92% 5% • •
1	C	298	86% 11% • •
1	D	298	88% 9% • •

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 9576 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uncharacterized protein yagE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	298	2263	1449	385	423	6	0	0	0
1	B	298	2265	1450	388	421	6	0	0	0
1	C	298	2269	1454	388	421	6	0	1	0
1	D	298	2268	1451	388	423	6	0	0	0

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	4	2	2	0	0
2	B	1	4	2	2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	C	1	4	2	2	0	0
			Total	C	O		
2	D	1	4	2	2	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	123	123	123	0	0
			Total	O		
3	B	136	136	136	0	0
			Total	O		
3	C	117	117	117	0	0
			Total	O		
3	D	119	119	119	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Uncharacterized protein yagE

Chain A:



• Molecule 1: Uncharacterized protein yagE

Chain B:



• Molecule 1: Uncharacterized protein yagE

Chain C:



• Molecule 1: Uncharacterized protein yagE

Chain D:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α, β, γ	141.13Å 153.55Å 55.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.98 – 2.20 48.98 – 2.20	Depositor EDS
% Data completeness (in resolution range)	100.0 (48.98-2.20) 99.3 (48.98-2.20)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	5.37 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.179 , 0.222 0.177 , 0.220	Depositor DCC
R_{free} test set	3106 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å^2)	15.4	Xtriage
Anisotropy	0.634	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.36 , 33.8	EDS
L-test for twinning[2]	$< \|L\| > = 0.47$, $< L^2 > = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	9576	wwPDB-VP
Average B, all atoms (Å^2)	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 41.91 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.2019e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, KPI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	1.03	0/2296	1.00	9/3129 (0.3%)
1	B	0.96	0/2298	0.91	8/3131 (0.3%)
1	C	1.00	2/2303 (0.1%)	1.04	10/3138 (0.3%)
1	D	1.10	4/2301 (0.2%)	0.98	16/3135 (0.5%)
All	All	1.02	6/9198 (0.1%)	0.98	43/12533 (0.3%)

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	134	VAL	CB-CG2	-12.41	1.26	1.52
1	D	15	THR	CB-OG1	-12.06	1.19	1.43
1	D	134	VAL	CB-CG1	-11.99	1.27	1.52
1	D	99	GLU	CG-CD	5.44	1.60	1.51
1	C	106	GLN	CD-NE2	-5.42	1.19	1.32

The worst 5 of 43 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	C	73	ARG	NE-CZ-NH1	-19.66	110.47	120.30
1	A	73	ARG	NE-CZ-NH1	-19.13	110.73	120.30
1	C	73	ARG	NE-CZ-NH2	13.94	127.27	120.30
1	D	77	ASP	CB-CG-OD1	12.33	129.40	118.30
1	A	73	ARG	NE-CZ-NH2	12.14	126.37	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2263	0	2280	18	0
1	B	2265	0	2287	7	0
1	C	2269	0	2284	24	1
1	D	2268	0	2289	18	0
2	A	4	0	6	0	0
2	B	4	0	6	0	0
2	C	4	0	6	0	0
2	D	4	0	6	2	0
3	A	123	0	0	5	0
3	B	136	0	0	1	1
3	C	117	0	0	4	0
3	D	119	0	0	4	0
All	All	9576	0	9164	54	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 54 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:129:ARG:HD2	3:D:374:HOH:O	1.77	0.85
1:C:122:VAL:HG22	1:D:264:PRO:HD3	1.65	0.77
1:C:264:PRO:HD3	1:D:122:VAL:HG22	1.66	0.77
1:A:264:PRO:HD3	1:B:122:VAL:HG22	1.68	0.76
1:A:122:VAL:HG22	1:B:264:PRO:HD3	1.70	0.74

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:73:ARG:NH2	3:B:430:HOH:O[4_454]	2.18	0.02

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	295/298 (99%)	288 (98%)	7 (2%)	0	100	100
1	B	295/298 (99%)	290 (98%)	5 (2%)	0	100	100
1	C	296/298 (99%)	291 (98%)	5 (2%)	0	100	100
1	D	295/298 (99%)	289 (98%)	6 (2%)	0	100	100
All	All	1181/1192 (99%)	1158 (98%)	23 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	241/242 (100%)	227 (94%)	14 (6%)	20	23
1	B	241/242 (100%)	226 (94%)	15 (6%)	18	21
1	C	240/242 (99%)	227 (95%)	13 (5%)	22	26
1	D	242/242 (100%)	227 (94%)	15 (6%)	18	21
All	All	964/968 (100%)	907 (94%)	57 (6%)	19	23

5 of 57 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	308	LEU
1	D	275	LEU
1	C	183	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	252	LEU
1	D	145	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	102	GLN
1	D	62	GLN
1	B	254	GLN
1	C	188	HIS
1	B	102	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
1	KPI	A	174	1	11,13,14	1.09	0	10,15,17	2.57	3 (30%)
1	KPI	D	174	1	11,13,14	0.80	0	10,15,17	2.63	2 (20%)
1	KPI	C	174	1	11,13,14	1.06	1 (9%)	10,15,17	2.38	5 (50%)
1	KPI	B	174	1	11,13,14	1.48	2 (18%)	10,15,17	2.23	4 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KPI	A	174	1	-	5/13/14/16	-
1	KPI	D	174	1	-	3/13/14/16	-
1	KPI	C	174	1	-	4/13/14/16	-
1	KPI	B	174	1	-	3/13/14/16	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	174	KPI	O1-CX2	-2.33	1.23	1.30
1	C	174	KPI	O1-CX2	-2.28	1.23	1.30
1	B	174	KPI	CX2-CX1	2.11	1.52	1.49

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	D	174	KPI	C1-CX1-CX2	-7.33	111.04	118.17
1	A	174	KPI	C1-CX1-CX2	-6.84	111.52	118.17
1	B	174	KPI	C1-CX1-CX2	-4.23	114.05	118.17
1	C	174	KPI	C1-CX1-CX2	-4.02	114.25	118.17
1	C	174	KPI	O2-CX2-CX1	-3.96	116.32	121.38

There are no chirality outliers.

5 of 15 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	174	KPI	C1-CX1-NZ-CE
1	A	174	KPI	NZ-CX1-CX2-O1
1	A	174	KPI	C1-CX1-CX2-O2
1	B	174	KPI	C1-CX1-NZ-CE
1	B	174	KPI	NZ-CX1-CX2-O1

There are no ring outliers.

4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	174	KPI	2	0
1	D	174	KPI	2	0
1	C	174	KPI	3	0
1	B	174	KPI	1	0

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	EDO	A	1	-	3,3,3	0.37	0	2,2,2	0.72	0
2	EDO	D	4	-	3,3,3	0.48	0	2,2,2	0.58	0
2	EDO	C	3	-	3,3,3	0.61	0	2,2,2	0.67	0
2	EDO	B	2	-	3,3,3	0.33	0	2,2,2	1.92	1 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	1	-	-	0/1/1/1	-
2	EDO	D	4	-	-	1/1/1/1	-
2	EDO	C	3	-	-	0/1/1/1	-
2	EDO	B	2	-	-	0/1/1/1	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	B	2	EDO	O2-C2-C1	-2.52	93.81	111.91

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	4	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	4	EDO	2	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	297/298 (99%)	-0.60	0	100	100	7, 12, 23, 35	0
1	B	297/298 (99%)	-0.58	0	100	100	7, 12, 22, 34	0
1	C	297/298 (99%)	-0.71	1 (0%)	94	93	7, 12, 22, 33	0
1	D	297/298 (99%)	-0.72	0	100	100	7, 12, 23, 34	0
All	All	1188/1192 (99%)	-0.65	1 (0%)	95	95	7, 12, 23, 35	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	28	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
1	KPI	D	174	14/15	0.95	0.09	5,9,19,23	0
1	KPI	B	174	14/15	0.96	0.12	6,9,19,22	0
1	KPI	C	174	14/15	0.97	0.12	6,9,19,23	0
1	KPI	A	174	14/15	0.97	0.10	5,9,19,22	0

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	EDO	C	3	4/4	0.95	0.15	17,17,18,24	0
2	EDO	B	2	4/4	0.96	0.15	13,14,17,20	0
2	EDO	A	1	4/4	0.96	0.12	16,20,20,26	0
2	EDO	D	4	4/4	0.96	0.14	9,18,22,26	0

6.5 Other polymers (i)

There are no such residues in this entry.