



wwPDB EM Validation Summary Report ⓘ

Apr 15, 2026 – 10:15 PM EDT

PDB ID : 9N37 / pdb_00009n37
EMDB ID : EMD-48847
Title : Cryo-EM structure of Rubisco from Arabidopsis thaliana with the 1A small subunit isoform
Authors : Stavros, A.; Askey, B.; Ceminsky, M.; Gunn, L.H.
Deposited on : 2025-01-30
Resolution : 2.20 Å(reported)
Based on initial model : 5IU0

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	3184 (1.71 - 2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	479	 82% 5% 13%
1	B	479	 81% 6% 13%
1	C	479	 81% 7% 13%
1	D	479	 83% 13%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	479	 5% 83% 5% 13%
1	F	479	 5% 81% 7% 13%
1	G	479	 5% 83% 5% 13%
1	H	479	 5% 82% 6% 13%
2	I	131	 5% 83% 6% 11%
2	J	131	 5% 82% 8% 11%
2	K	131	 5% 82% 8% 11%
2	L	131	 5% 82% 7% 11%
2	M	131	 83% 6% 11%
2	N	131	 5% 82% 8% 11%
2	O	131	 85% 5% 11%
2	P	131	 5% 85% 5% 11%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 67150 atoms, of which 33158 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ribulose biphosphate carboxylase large chain.

Mol	Chain	Residues	Atoms						AltConf	Trace
1	A	419	Total	C	H	N	O	S	0	0
			6434	2062	3175	574	607	16		
1	B	419	Total	C	H	N	O	S	0	0
			6444	2062	3185	574	607	16		
1	C	419	Total	C	H	N	O	S	0	0
			6444	2062	3185	574	607	16		
1	D	419	Total	C	H	N	O	S	0	0
			6444	2062	3185	574	607	16		
1	E	419	Total	C	H	N	O	S	0	0
			6444	2062	3185	574	607	16		
1	F	419	Total	C	H	N	O	S	0	0
			6444	2062	3185	574	607	16		
1	G	419	Total	C	H	N	O	S	0	0
			6444	2062	3185	574	607	16		
1	H	419	Total	C	H	N	O	S	0	0
			6444	2062	3185	574	607	16		

- Molecule 2 is a protein called Ribulose biphosphate carboxylase small subunit 1A, chloroplastic.

Mol	Chain	Residues	Atoms						AltConf	Trace
2	I	117	Total	C	H	N	O	S	2	0
			1950	649	961	158	178	4		
2	J	117	Total	C	H	N	O	S	2	0
			1950	649	961	158	178	4		
2	K	117	Total	C	H	N	O	S	2	0
			1950	649	961	158	178	4		
2	L	117	Total	C	H	N	O	S	2	0
			1950	649	961	158	178	4		
2	M	117	Total	C	H	N	O	S	2	0
			1950	649	961	158	178	4		
2	N	117	Total	C	H	N	O	S	2	0
			1950	649	961	158	178	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms						AltConf	Trace
2	O	117	Total	C	H	N	O	S	2	0
			1950	649	961	158	178	4		
2	P	117	Total	C	H	N	O	S	2	0
			1950	649	961	158	178	4		

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
I	126	HIS	-	expression tag	UNP P10795
I	127	HIS	-	expression tag	UNP P10795
I	128	HIS	-	expression tag	UNP P10795
I	129	HIS	-	expression tag	UNP P10795
I	130	HIS	-	expression tag	UNP P10795
I	131	HIS	-	expression tag	UNP P10795
J	126	HIS	-	expression tag	UNP P10795
J	127	HIS	-	expression tag	UNP P10795
J	128	HIS	-	expression tag	UNP P10795
J	129	HIS	-	expression tag	UNP P10795
J	130	HIS	-	expression tag	UNP P10795
J	131	HIS	-	expression tag	UNP P10795
K	126	HIS	-	expression tag	UNP P10795
K	127	HIS	-	expression tag	UNP P10795
K	128	HIS	-	expression tag	UNP P10795
K	129	HIS	-	expression tag	UNP P10795
K	130	HIS	-	expression tag	UNP P10795
K	131	HIS	-	expression tag	UNP P10795
L	126	HIS	-	expression tag	UNP P10795
L	127	HIS	-	expression tag	UNP P10795
L	128	HIS	-	expression tag	UNP P10795
L	129	HIS	-	expression tag	UNP P10795
L	130	HIS	-	expression tag	UNP P10795
L	131	HIS	-	expression tag	UNP P10795
M	126	HIS	-	expression tag	UNP P10795
M	127	HIS	-	expression tag	UNP P10795
M	128	HIS	-	expression tag	UNP P10795
M	129	HIS	-	expression tag	UNP P10795
M	130	HIS	-	expression tag	UNP P10795
M	131	HIS	-	expression tag	UNP P10795
N	126	HIS	-	expression tag	UNP P10795
N	127	HIS	-	expression tag	UNP P10795
N	128	HIS	-	expression tag	UNP P10795
N	129	HIS	-	expression tag	UNP P10795

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
N	130	HIS	-	expression tag	UNP P10795
N	131	HIS	-	expression tag	UNP P10795
O	126	HIS	-	expression tag	UNP P10795
O	127	HIS	-	expression tag	UNP P10795
O	128	HIS	-	expression tag	UNP P10795
O	129	HIS	-	expression tag	UNP P10795
O	130	HIS	-	expression tag	UNP P10795
O	131	HIS	-	expression tag	UNP P10795
P	126	HIS	-	expression tag	UNP P10795
P	127	HIS	-	expression tag	UNP P10795
P	128	HIS	-	expression tag	UNP P10795
P	129	HIS	-	expression tag	UNP P10795
P	130	HIS	-	expression tag	UNP P10795
P	131	HIS	-	expression tag	UNP P10795

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms	AltConf
3	A	1	Total Mg 1 1	0
3	B	1	Total Mg 1 1	0
3	C	1	Total Mg 1 1	0
3	D	1	Total Mg 1 1	0
3	E	1	Total Mg 1 1	0
3	F	1	Total Mg 1 1	0
3	G	1	Total Mg 1 1	0
3	H	1	Total Mg 1 1	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ribulose biphosphate carboxylase large chain

Chain A:




- Molecule 1: Ribulose biphosphate carboxylase large chain

Chain B:



- Molecule 1: Ribulose biphosphate carboxylase large chain


Chain C:



GLU
VAL
TRP
LYS
GLU
ILE
THR
PHE
ASN
PHE
PHE
PRO
THR
THR
ILE
ASP
LYS
LEU
GLY
GLN
GLU


- Molecule 1: Ribulose biphosphate carboxylase large chain

Chain D:




- Molecule 1: Ribulose biphosphate carboxylase large chain

Chain E:




- Molecule 1: Ribulose biphosphate carboxylase large chain


Chain F:





- Molecule 1: Ribulose biphosphate carboxylase large chain

Chain G:








- Molecule 1: Ribulose biphosphate carboxylase large chain





- Molecule 2: Ribulose biphosphate carboxylase small subunit 1A, chloroplastic




- Molecule 2: Ribulose biphosphate carboxylase small subunit 1A, chloroplastic



- Molecule 2: Ribulose biphosphate carboxylase small subunit 1A, chloroplastic



- Molecule 2: Ribulose biphosphate carboxylase small subunit 1A, chloroplastic



- Molecule 2: Ribulose biphosphate carboxylase small subunit 1A, chloroplastic

Chain M:  83% 6% 11%




- Molecule 2: Ribulose biphosphate carboxylase small subunit 1A, chloroplastic

Chain N:  82% 8% 11%



- Molecule 2: Ribulose biphosphate carboxylase small subunit 1A, chloroplastic

Chain O:  85% 5% 11%



- Molecule 2: Ribulose biphosphate carboxylase small subunit 1A, chloroplastic

Chain P:  85% 0% 11%



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, D4	Depositor
Number of particles used	256188	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	40	Depositor
Minimum defocus (nm)	600	Depositor
Maximum defocus (nm)	2400	Depositor
Magnification	130000	Depositor
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.673	Depositor
Minimum map value	-0.423	Depositor
Average map value	0.002	Depositor
Map value standard deviation	0.055	Depositor
Recommended contour level	0.15	Depositor
Map size (Å)	212.48, 212.48, 212.48	wwPDB
Map dimensions	320, 320, 320	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.664, 0.664, 0.664	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, KCX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.17	0/3321	0.37	0/4498
1	B	0.17	0/3321	0.37	0/4498
1	C	0.17	0/3321	0.36	0/4498
1	D	0.18	0/3321	0.36	0/4498
1	E	0.17	0/3321	0.38	0/4498
1	F	0.17	0/3321	0.37	0/4498
1	G	0.17	0/3321	0.37	0/4498
1	H	0.17	0/3321	0.36	0/4498
2	I	0.16	0/1028	0.33	0/1395
2	J	0.17	0/1028	0.34	0/1395
2	K	0.17	0/1028	0.34	0/1395
2	L	0.16	0/1028	0.33	0/1395
2	M	0.18	0/1028	0.35	0/1395
2	N	0.16	0/1028	0.34	0/1395
2	O	0.17	0/1028	0.33	0/1395
2	P	0.16	0/1028	0.32	0/1395
All	All	0.17	0/34792	0.36	0/47144

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3259	3175	3179	15	0
1	B	3259	3185	3179	17	0
1	C	3259	3185	3179	20	0
1	D	3259	3185	3179	13	0
1	E	3259	3185	3179	13	0
1	F	3259	3185	3179	19	0
1	G	3259	3185	3179	13	0
1	H	3259	3185	3179	13	0
2	I	989	961	951	4	0
2	J	989	961	951	6	0
2	K	989	961	951	5	0
2	L	989	961	951	6	0
2	M	989	961	951	6	0
2	N	989	961	951	7	0
2	O	989	961	951	5	0
2	P	989	961	951	4	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0
3	F	1	0	0	0	0
3	G	1	0	0	0	0
3	H	1	0	0	0	0
All	All	33992	33158	33040	142	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 142 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:187:ARG:NH1	2:K:43:GLU:OE2	2.20	0.74
2:L:43:GLU:OE1	1:F:187:ARG:NH1	2.20	0.74
2:N:43:GLU:OE2	1:H:187:ARG:NH1	2.20	0.73
2:M:43:GLU:OE1	1:C:187:ARG:NH1	2.24	0.71
1:H:94:GLU:N	1:H:94:GLU:OE2	2.25	0.70

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	412/479 (86%)	397 (96%)	15 (4%)	0	100	100
1	B	412/479 (86%)	396 (96%)	16 (4%)	0	100	100
1	C	412/479 (86%)	394 (96%)	18 (4%)	0	100	100
1	D	412/479 (86%)	397 (96%)	15 (4%)	0	100	100
1	E	412/479 (86%)	398 (97%)	14 (3%)	0	100	100
1	F	412/479 (86%)	398 (97%)	14 (3%)	0	100	100
1	G	412/479 (86%)	398 (97%)	14 (3%)	0	100	100
1	H	412/479 (86%)	395 (96%)	17 (4%)	0	100	100
2	I	115/131 (88%)	112 (97%)	2 (2%)	1 (1%)	14	14
2	J	115/131 (88%)	111 (96%)	4 (4%)	0	100	100
2	K	115/131 (88%)	111 (96%)	4 (4%)	0	100	100
2	L	115/131 (88%)	111 (96%)	4 (4%)	0	100	100
2	M	115/131 (88%)	110 (96%)	5 (4%)	0	100	100
2	N	115/131 (88%)	109 (95%)	6 (5%)	0	100	100
2	O	115/131 (88%)	111 (96%)	4 (4%)	0	100	100
2	P	115/131 (88%)	111 (96%)	4 (4%)	0	100	100
All	All	4216/4880 (86%)	4059 (96%)	156 (4%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	I	13	GLU

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM

entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	337/388 (87%)	334 (99%)	3 (1%)	70	84
1	B	337/388 (87%)	335 (99%)	2 (1%)	78	89
1	C	337/388 (87%)	334 (99%)	3 (1%)	70	84
1	D	337/388 (87%)	334 (99%)	3 (1%)	70	84
1	E	337/388 (87%)	334 (99%)	3 (1%)	70	84
1	F	337/388 (87%)	333 (99%)	4 (1%)	63	78
1	G	337/388 (87%)	334 (99%)	3 (1%)	70	84
1	H	337/388 (87%)	332 (98%)	5 (2%)	57	73
2	I	109/119 (92%)	107 (98%)	2 (2%)	51	68
2	J	109/119 (92%)	106 (97%)	3 (3%)	38	52
2	K	109/119 (92%)	106 (97%)	3 (3%)	38	52
2	L	109/119 (92%)	106 (97%)	3 (3%)	38	52
2	M	109/119 (92%)	107 (98%)	2 (2%)	51	68
2	N	109/119 (92%)	106 (97%)	3 (3%)	38	52
2	O	109/119 (92%)	107 (98%)	2 (2%)	51	68
2	P	109/119 (92%)	107 (98%)	2 (2%)	51	68
All	All	3568/4056 (88%)	3522 (99%)	46 (1%)	59	76

5 of 46 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	134	ARG
1	F	134	ARG
1	D	382	ILE
1	E	382	ILE
1	F	459	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 33 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	229	GLN
1	G	294	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	H	294	HIS
2	O	48	HIS
2	N	82	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	KCX	H	201	1,3	10,11,12	1.40	1 (10%)	6,12,14	2.13	1 (16%)
1	KCX	A	201	1,3	10,11,12	1.39	1 (10%)	6,12,14	2.18	1 (16%)
1	KCX	F	201	1,3	10,11,12	1.40	1 (10%)	6,12,14	2.15	1 (16%)
1	KCX	D	201	1,3	10,11,12	1.40	1 (10%)	6,12,14	2.14	1 (16%)
1	KCX	E	201	1,3	10,11,12	1.40	1 (10%)	6,12,14	2.14	1 (16%)
1	KCX	B	201	1	10,11,12	1.41	1 (10%)	6,12,14	2.12	1 (16%)
1	KCX	G	201	1,3	10,11,12	1.39	2 (20%)	6,12,14	2.18	1 (16%)
1	KCX	C	201	1,3	10,11,12	1.40	1 (10%)	6,12,14	2.16	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	H	201	1,3	-	0/9/10/12	-
1	KCX	A	201	1,3	-	0/9/10/12	-
1	KCX	F	201	1,3	-	0/9/10/12	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	D	201	1,3	-	0/9/10/12	-
1	KCX	E	201	1,3	-	0/9/10/12	-
1	KCX	B	201	1	-	0/9/10/12	-
1	KCX	G	201	1,3	-	0/9/10/12	-
1	KCX	C	201	1,3	-	0/9/10/12	-

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	201	KCX	CX-NZ	3.57	1.41	1.35
1	B	201	KCX	CX-NZ	3.57	1.41	1.35
1	E	201	KCX	CX-NZ	3.54	1.41	1.35
1	F	201	KCX	CX-NZ	3.53	1.41	1.35
1	D	201	KCX	CX-NZ	3.52	1.41	1.35

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	201	KCX	OQ1-CX-NZ	-5.18	117.05	124.92
1	A	201	KCX	OQ1-CX-NZ	-5.16	117.08	124.92
1	C	201	KCX	OQ1-CX-NZ	-5.10	117.17	124.92
1	F	201	KCX	OQ1-CX-NZ	-5.08	117.20	124.92
1	D	201	KCX	OQ1-CX-NZ	-5.07	117.22	124.92

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-48847. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 160




Y Index: 160




Z Index: 160


6.2.2 Raw map



X Index: 160



Y Index: 160




Z Index: 160


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 136




Y Index: 184




Z Index: 147


6.3.2 Raw map



X Index: 136



Y Index: 136




Z Index: 148


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.15. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 157 nm³; this corresponds to an approximate mass of 142 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.455 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of $0.455 \text{ \AA}^{-1}$

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.20	-	-
Author-provided FSC curve	2.19	2.50	2.22
Unmasked-calculated*	2.64	2.96	2.66

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 2.64 differs from the reported value 2.2 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-48847 and PDB model 9N37. Per-residue inclusion information can be found in section [3](#) on page [7](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.15 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.15).

9.4 Atom inclusion [i](#)



At the recommended contour level, 85% of all backbone atoms, 85% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.15) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8530	0.6570
A	0.8590	0.6600
B	0.8570	0.6620
C	0.8590	0.6600
D	0.8580	0.6610
E	0.8620	0.6600
F	0.8620	0.6580
G	0.8630	0.6590
H	0.8630	0.6580
I	0.8430	0.6510
J	0.8480	0.6480
K	0.8470	0.6520
L	0.8440	0.6480
M	0.8410	0.6470
N	0.8400	0.6450
O	0.8450	0.6470
P	0.8370	0.6430

