wwPDB X-ray Structure Validation Summary Report (i)

Jun 16, 2024 – 08:11 PM EDT

PDB ID	:	5NGJ
Title	:	Crystal structure of pb6, major tail tube protein of bacteriophage T5
Authors	:	Arnaud, C.-A.; Effantin, G.; Vives, C.; Engilberge, S.; Bacia, M.; Boulanger, P.; Girard, E.; Schoehn, G.; Breyton, C.
Deposited on	:	2017-03-17
Resolution	:	2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	477	10% 89% • 8%
1	B	477	25% 84% 6% 9%

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 13513 atoms, of which 6596 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tail tube protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	440	6729	2128	3347	563	682	9	0	1	0
1	B	433	6551	2073	3249	552	668	9	0	2	1

There are 26 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	465	GLU	-	expression tag	UNP Q6QGE2
A	466	ASN	-	expression tag	UNP Q6QGE2
A	467	LEU	-	expression tag	UNP Q6QGE2
A	468	TYR	-	expression tag	UNP Q6QGE2
A	469	PHE	-	expression tag	UNP Q6QGE2
A	470	GLN	-	expression tag	UNP Q6QGE2
A	471	GLY	-	expression tag	UNP Q6QGE2
A	472	HIS	-	expression tag	UNP Q6QGE2
A	473	HIS	-	expression tag	UNP Q6QGE2
A	474	HIS	-	expression tag	UNP Q6QGE2
A	475	HIS	-	expression tag	UNP Q6QGE2
A	476	HIS	-	expression tag	UNP Q6QGE2
A	477	HIS	-	expression tag	UNP Q6QGE2
B	465	GLU	-	expression tag	UNP Q6QGE2
B	466	ASN	-	expression tag	UNP Q6QGE2
B	467	LEU	-	expression tag	UNP Q6QGE2
B	468	TYR	-	expression tag	UNP Q6QGE2
B	469	PHE	-	expression tag	UNP Q6QGE2
B	470	GLN	-	expression tag	UNP Q6QGE2
B	471	GLY	-	expression tag	UNP Q6QGE2
B	472	HIS	-	expression tag	UNP Q6QGE2
B	473	HIS	-	expression tag	UNP Q6QGE2
B	474	HIS	-	expression tag	UNP Q6QGE2
B	475	HIS	-	expression tag	UNP Q6QGE2
B	476	HIS	-	expression tag	UNP Q6QGE2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	477	HIS	-	expression tag	UNP Q6QGE2

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Cl		
2	A	5	5	5	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	145	145	145	0	0
			Total	O		
3	B	83	83	83	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tail tube protein

Chain A:



- Molecule 1: Tail tube protein

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	76.79Å 115.08Å 83.37Å 90.00° 111.88° 90.00°	Depositor
Resolution (Å)	44.00 – 2.20 57.32 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.7 (44.00-2.20) 99.7 (57.32-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.73 (at 2.20Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.205 , 0.235 0.209 , 0.235	Depositor DCC
R_{free} test set	3400 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å^2)	56.8	Xtriage
Anisotropy	0.495	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.39 , 71.1	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	13513	wwPDB-VP
Average B, all atoms (Å^2)	86.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.42% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.52	0/3442	0.71	2/4685 (0.0%)
1	B	0.46	0/3362	0.69	1/4575 (0.0%)
All	All	0.49	0/6804	0.70	3/9260 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	208	LEU	CD1-CG-CD2	5.57	127.21	110.50
1	A	98	LEU	CA-CB-CG	5.24	127.35	115.30
1	A	208	LEU	CD1-CG-CD2	5.03	125.59	110.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3382	3347	3347	3	0
1	B	3302	3249	3221	9	0
2	A	5	0	0	0	0
3	A	145	0	0	0	0
3	B	83	0	0	0	0
All	All	6917	6596	6568	12	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 12 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:376:VAL:HG12	1:B:452:VAL:HG11	1.67	0.76
1:B:213:ILE:HG22	1:B:295:ILE:HG22	1.82	0.61
1:B:376:VAL:CG1	1:B:452:VAL:HG11	2.32	0.60
1:B:159:VAL:O	1:B:159:VAL:HG12	2.11	0.51
1:B:450:SER:OG	1:B:452:VAL:HG12	2.12	0.49

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	437/477 (92%)	423 (97%)	14 (3%)	0	100	100
1	B	427/477 (90%)	406 (95%)	17 (4%)	4 (1%)	17	16
All	All	864/954 (91%)	829 (96%)	31 (4%)	4 (0%)	29	31

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	448	ASP
1	B	393	SER
1	B	329	VAL
1	B	47	VAL

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	374/406 (92%)	366 (98%)	8 (2%)	53	67
1	B	362/406 (89%)	348 (96%)	14 (4%)	32	41
All	All	736/812 (91%)	714 (97%)	22 (3%)	41	53

5 of 22 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	274	LEU
1	B	390	LEU
1	B	343	LEU
1	B	397	GLN
1	A	437	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 5 ligands modelled in this entry, 5 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	440/477 (92%)	0.84	47 (10%)	6	5	49, 70, 112, 149	0
1	B	433/477 (90%)	1.45	120 (27%)	0	0	57, 86, 178, 285	0
All	All	873/954 (91%)	1.15	167 (19%)	1	1	49, 76, 139, 285	0

The worst 5 of 167 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	389	THR	13.7
1	B	458	ILE	10.8
1	B	426	ALA	8.4
1	B	429	LEU	7.5
1	B	394	ALA	7.3

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	CL	A	503	1/1	0.29	0.18	107,107,107,107	0
2	CL	A	505	1/1	0.86	0.35	91,91,91,91	1
2	CL	A	504	1/1	0.90	0.25	92,92,92,92	0
2	CL	A	501	1/1	0.90	0.15	56,56,56,56	1
2	CL	A	502	1/1	0.96	0.29	50,50,50,50	1

6.5 Other polymers (i)

There are no such residues in this entry.