



Full wwPDB X-ray Structure Validation Report i

Jun 12, 2024 – 03:41 AM EDT

PDB ID : 6NGX
Title : Structure of rat neuronal nitric oxide synthase heme domain in complex with 6-(2,3-difluoro-5-(3-(methylamino)prop-1-yn-1-yl)phenethyl)-4-methylpyridin-2-amine
Authors : Li, H.; Poulos, T.L.
Deposited on : 2018-12-21
Resolution : 1.77 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriaage (Phenix) : 1.13
EDS : 2.36.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	9185 (1.80-1.76)
Clashscore	141614	10184 (1.80-1.76)
Ramachandran outliers	138981	10051 (1.80-1.76)
Sidechain outliers	138945	10050 (1.80-1.76)
RSRZ outliers	127900	9032 (1.80-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition


There are 8 unique types of molecules in this entry. The entry contains 7345 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitric oxide synthase, brain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	411	Total	C	N	O	S	0	2	0
			3348	2141	573	612	22			
1	B	411	Total	C	N	O	S	0	3	0
			3357	2147	575	613	22			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula: C₉H₁₅N₅O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O 17 9 5 3	0	0
3	B	1	Total C N O 17 9 5 3	0	0

- Molecule 4 is 6-(2,3-difluoro-5-[3-(methylamino)prop-1-yn-1-yl]phenyl)-4-methylpyridin-2-amine (three-letter code: KMV) (formula: C₁₈H₁₉F₂N₃).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C F N 23 18 2 3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	F	N	0	0
			23	18	2	3		

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	O		0	0
			4	2	2			
5	B	1	Total	C	O		0	0
			4	2	2			

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	Zn			0	0
			1	1				

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	B	1	Total C O 6 3 3	0	0
7	B	1	Total C O 6 3 3	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	188	Total O 188 188	0	0
8	B	265	Total O 265 265	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nitric oxide synthase, brain



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	52.02Å 111.01Å 164.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.00 – 1.77 39.00 – 1.77	Depositor EDS
% Data completeness (in resolution range)	99.0 (39.00-1.77) 99.4 (39.00-1.77)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	0.11	Depositor
$< I/\sigma(I) >$ ¹	1.00 (at 1.77Å)	Xtriage
Refinement program	PHENIX (1.11.1-2575_1496: ???)	Depositor
R , R_{free}	0.178 , 0.211 0.178 , 0.210	Depositor DCC
R_{free} test set	4615 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	32.3	Xtriage
Anisotropy	0.838	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 56.1	EDS
L-test for twinning ²	$< L > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7345	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.50% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, HEM, H4B, ACT, ZN, KMV

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/3444	0.51	0/4672
1	B	0.36	0/3456	0.52	0/4685
All	All	0.36	0/6900	0.51	0/9357

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3348	0	3257	23	0
1	B	3357	0	3273	24	0
2	A	43	0	30	8	0
2	B	43	0	30	7	0
3	A	17	0	15	0	0
3	B	17	0	15	0	0
4	A	23	0	0	5	0
4	B	23	0	0	4	0
5	A	4	0	3	0	0
5	B	4	0	3	0	0
6	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	B	12	0	16	1	0
8	A	188	0	0	4	0
8	B	265	0	0	3	0
All	All	7345	0	6642	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:801:HEM:O1D	8:B:901:HOH:O	2.03	0.77
2:A:801:HEM:O1D	8:A:901:HOH:O	2.04	0.76
1:A:523:LEU:HD22	1:A:531:PRO:HB2	1.68	0.74
1:A:363:ASP:OD1	8:A:902:HOH:O	2.05	0.73
2:B:801:HEM:HBB2	2:B:801:HEM:HHC	1.72	0.72
2:B:801:HEM:HMC2	2:B:801:HEM:HBC2	1.73	0.71
1:A:351:LYS:O	8:A:903:HOH:O	2.09	0.71
1:B:364:GLN:OE1	8:B:902:HOH:O	2.10	0.69
2:A:801:HEM:HBB2	2:A:801:HEM:HHC	1.76	0.68
2:A:801:HEM:HMC2	2:A:801:HEM:HBC2	1.75	0.67
1:B:706:TYR:OH	2:B:801:HEM:O2D	2.13	0.66
1:B:663:GLU:HB3	1:B:667:ARG:HH12	1.60	0.65
1:B:478:GLN:HB2	1:B:481:ARG:HG3	1.82	0.62
1:A:478:GLN:HB2	1:A:481:ARG:HG3	1.85	0.58
1:B:567:VAL:HG21	4:B:803:KMV:F12	1.94	0.58
2:A:801:HEM:HBA1	4:A:803:KMV:C09	2.35	0.57
1:A:678:TRP:HZ3	2:A:801:HEM:HBA2	1.70	0.56
1:B:663:GLU:HB3	1:B:667:ARG:NH1	2.21	0.55
1:B:570:MET:SD	4:B:803:KMV:F13	2.55	0.55
1:B:595:VAL:O	1:B:599:CYS:HB2	2.07	0.55
1:B:494:GLY:O	7:B:805:GOL:H12	2.08	0.54
1:B:525:GLN:HE21	1:B:528:GLY:HA2	1.73	0.54
1:B:678:TRP:HZ3	2:B:801:HEM:HBA2	1.71	0.53
2:A:801:HEM:HBD1	4:A:803:KMV:C13	2.37	0.53
1:A:504:ILE:O	1:A:508:GLN:HB2	2.08	0.53
1:A:567:VAL:HG21	4:A:803:KMV:F12	1.99	0.52
2:B:801:HEM:HBA1	4:B:803:KMV:C09	2.39	0.51
1:A:678:TRP:CZ3	2:A:801:HEM:HBA2	2.46	0.51
1:A:299:ARG:HG3	1:A:318:LEU:HD21	1.93	0.49
1:A:475:TRP:HB2	1:A:523:LEU:HB3	1.93	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:567:VAL:HG21	4:A:803:KMF:C12	2.43	0.48
1:B:567:VAL:HG21	4:B:803:KMF:C12	2.45	0.47
1:A:601:ASN:ND2	8:A:905:HOH:O	2.35	0.47
1:A:351:LYS:HE2	1:A:392:SER:HB3	1.98	0.45
1:A:525:GLN:HG3	1:A:529:ASN:O	2.16	0.45
1:B:325:GLY:O	1:B:332:MET:HG3	2.16	0.45
1:B:403:TYR:CE1	1:B:407:HIS:CE1	3.05	0.45
1:A:403:TYR:CE1	1:A:407:HIS:CE1	3.06	0.44
1:B:510:TRP:CE2	1:B:521:PRO:HD3	2.53	0.44
2:A:801:HEM:HBD1	4:A:803:KMF:C12	2.48	0.44
1:B:678:TRP:CZ3	2:B:801:HEM:HBA2	2.53	0.43
1:A:510:TRP:CE2	1:A:521:PRO:HD3	2.53	0.43
1:B:386:LYS:HB3	1:B:386:LYS:HE2	1.79	0.43
1:B:548:HIS:CG	1:B:549:PRO:HD2	2.54	0.43
1:B:366:TYR:HA	1:B:369:ILE:HG12	2.00	0.42
1:B:478:GLN:NE2	8:B:911:HOH:O	2.51	0.42
1:A:686:SER:HA	1:A:691:PHE:CG	2.55	0.42
1:A:645:LYS:HD3	1:A:645:LYS:HA	1.92	0.42
1:B:355:PHE:CE1	1:B:385:ASN:HB2	2.55	0.42
1:B:409:TRP:CE3	1:B:421:TRP:HA	2.55	0.41
1:A:370:LYS:HB2	1:A:370:LYS:HE3	1.77	0.41
1:B:355:PHE:N	1:B:356:PRO:HD2	2.35	0.41
1:A:480:ILE:HD13	1:A:541:VAL:HG13	2.03	0.41
1:A:676:TRP:CE2	1:A:680:VAL:HG21	2.56	0.41
1:B:386:LYS:O	1:B:390:SER:HB3	2.21	0.40
1:A:355:PHE:N	1:A:356:PRO:HD2	2.36	0.40
1:A:548:HIS:CE1	1:A:550:LYS:HB2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	408/422 (97%)	390 (96%)	16 (4%)	2 (0%)	29 14
1	B	410/422 (97%)	404 (98%)	6 (2%)	0	100 100
All	All	818/844 (97%)	794 (97%)	22 (3%)	2 (0%)	47 32

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	338	PRO
1	A	375	LYS

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	368/377 (98%)	356 (97%)	12 (3%)	38 21
1	B	369/377 (98%)	362 (98%)	7 (2%)	57 43
All	All	737/754 (98%)	718 (97%)	19 (3%)	46 29

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	321	THR
1	A	347	ASP
1	A	352	ASP
1	A	353	GLN
1	A	469	LYS
1	A	476	ASN
1	A	489	ASP
1	A	508	GLN
1	A	523	LEU
1	A	547	ARG
1	A	555	LYS
1	A	645	LYS
1	B	309	ASP
1	B	351	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	390	SER
1	B	392	SER
1	B	516	ARG
1	B	547	ARG
1	B	699	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

Of 11 ligands modelled in this entry, 1 is monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEM	B	801	1	41,50,50	1.51	5 (12%)	45,82,82	1.54	8 (17%)
4	KMV	B	803	-	23,24,24	1.38	1 (4%)	29,32,32	2.51	11 (37%)
3	H4B	B	802	-	16,18,18	0.80	0	11,26,26	2.68	5 (45%)
3	H4B	A	802	-	16,18,18	0.87	0	11,26,26	2.72	6 (54%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	KMV	A	803	-	23,24,24	1.38	1 (4%)	29,32,32	1.77	7 (24%)
2	HEM	A	801	1	41,50,50	1.52	6 (14%)	45,82,82	1.48	7 (15%)
5	ACT	A	804	-	3,3,3	0.77	0	3,3,3	0.73	0
5	ACT	B	804	-	3,3,3	0.79	0	3,3,3	0.60	0
7	GOL	B	806	-	5,5,5	0.36	0	5,5,5	0.39	0
7	GOL	B	805	-	5,5,5	0.35	0	5,5,5	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	B	801	1	-	2/12/54/54	-
4	KMV	B	803	-	-	0/8/10/10	0/2/2/2
3	H4B	B	802	-	-	0/8/17/17	0/2/2/2
3	H4B	A	802	-	-	0/8/17/17	0/2/2/2
4	KMV	A	803	-	-	1/8/10/10	0/2/2/2
2	HEM	A	801	1	-	4/12/54/54	-
7	GOL	B	806	-	-	2/4/4/4	-
7	GOL	B	805	-	-	4/4/4/4	-

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	803	KMV	C15-C17	-6.29	1.29	1.44
4	B	803	KMV	C15-C17	-6.18	1.29	1.44
2	B	801	HEM	C3C-C2C	-4.39	1.34	1.40
2	A	801	HEM	C3C-C2C	-4.07	1.34	1.40
2	A	801	HEM	C3C-CAC	3.69	1.55	1.47
2	B	801	HEM	C3C-CAC	3.49	1.55	1.47
2	A	801	HEM	CAB-C3B	2.96	1.55	1.47
2	B	801	HEM	CAB-C3B	2.87	1.55	1.47
2	A	801	HEM	CMD-C2D	2.30	1.55	1.50
2	A	801	HEM	FE-NB	2.28	2.08	1.96
2	B	801	HEM	CMD-C2D	2.22	1.55	1.50
2	B	801	HEM	CMB-C2B	2.06	1.55	1.50
2	A	801	HEM	C3B-C2B	-2.04	1.33	1.37

All (44) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	803	KMV	C02-N01-C06	6.27	122.86	118.10
4	B	803	KMV	C09-C11-C12	-5.57	116.25	120.73
3	A	802	H4B	C8A-C4A-C4	5.07	119.07	114.57
3	B	802	H4B	C8A-C4A-C4	4.36	118.44	114.57
4	A	803	KMV	C09-C11-C12	-4.31	117.27	120.73
4	A	803	KMV	C02-N01-C06	4.27	121.33	118.10
4	B	803	KMV	C05-C06-N01	-4.24	118.40	122.90
4	B	803	KMV	C08-C06-N01	4.21	122.21	115.95
2	B	801	HEM	CBA-CAA-C2A	-4.05	105.70	112.62
3	B	802	H4B	C4-C4A-N5	3.89	122.39	119.12
3	B	802	H4B	C2-N3-C4	3.83	122.02	115.93
3	A	802	H4B	N1-C2-N3	-3.52	119.90	125.42
3	A	802	H4B	C2-N3-C4	3.48	121.45	115.93
4	B	803	KMV	C14-C15-C17	-3.46	113.84	120.19
4	B	803	KMV	C16-C15-C17	3.43	126.48	120.19
3	B	802	H4B	N1-C2-N3	-3.38	120.12	125.42
2	A	801	HEM	CBA-CAA-C2A	-3.36	106.89	112.62
2	A	801	HEM	C4B-CHC-C1C	3.33	126.95	122.56
2	B	801	HEM	C4B-CHC-C1C	3.26	126.86	122.56
3	A	802	H4B	C4-C4A-N5	3.09	121.72	119.12
4	A	803	KMV	C05-C06-N01	-2.98	119.74	122.90
4	B	803	KMV	C08-C09-C11	2.91	117.53	112.81
3	A	802	H4B	C2-N1-C8A	2.91	121.06	114.54
2	B	801	HEM	CMA-C3A-C4A	-2.87	124.05	128.46
4	B	803	KMV	C16-C11-C12	2.84	119.36	116.76
4	A	803	KMV	C08-C06-N01	2.78	120.10	115.95
4	A	803	KMV	C09-C08-C06	-2.69	106.96	112.99
3	B	802	H4B	C2-N1-C8A	2.56	120.27	114.54
4	B	803	KMV	C09-C11-C16	2.46	124.36	119.41
4	A	803	KMV	C16-C11-C12	2.45	119.00	116.76
2	B	801	HEM	C3D-C4D-ND	-2.40	107.50	110.17
2	A	801	HEM	C4D-ND-C1D	2.37	107.52	105.07
3	A	802	H4B	N2-C2-N3	2.35	120.91	117.25
4	B	803	KMV	C15-C17-C18	-2.32	170.31	176.52
2	A	801	HEM	C3B-C2B-C1B	2.28	108.18	106.49
2	B	801	HEM	CHD-C1D-ND	2.28	126.90	124.43
2	B	801	HEM	CAD-C3D-C2D	-2.27	123.65	127.88
4	B	803	KMV	N02-C02-N01	2.20	119.97	116.49
2	A	801	HEM	CMC-C2C-C3C	2.16	128.73	124.68
2	A	801	HEM	C3D-C4D-ND	-2.15	107.78	110.17
4	A	803	KMV	C09-C11-C16	2.13	123.71	119.41
2	A	801	HEM	CHC-C4B-C3B	2.11	127.79	124.57
2	B	801	HEM	C4D-ND-C1D	2.09	107.23	105.07

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	801	HEM	CBD-CAD-C3D	-2.01	107.03	112.63

There are no chirality outliers.

All (13) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	801	HEM	C2A-CAA-CBA-CGA
4	A	803	KMV	C18-C19-N20-C21
7	B	806	GOL	O1-C1-C2-C3
2	B	801	HEM	C2A-CAA-CBA-CGA
7	B	806	GOL	O1-C1-C2-O2
7	B	805	GOL	O1-C1-C2-C3
7	B	805	GOL	C1-C2-C3-O3
7	B	805	GOL	O1-C1-C2-O2
2	A	801	HEM	C4B-C3B-CAB-CBB
2	B	801	HEM	C4B-C3B-CAB-CBB
7	B	805	GOL	O2-C2-C3-O3
2	A	801	HEM	CAD-CBD-CGD-O2D
2	A	801	HEM	CAD-CBD-CGD-O1D


There are no ring outliers.


5 monomers are involved in 21 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	801	HEM	7	0
4	B	803	KMV	4	0
4	A	803	KMV	5	0
2	A	801	HEM	8	0
7	B	805	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	411/422 (97%)	1.37	102 (24%) 0 0	32, 58, 110, 151	0
1	B	411/422 (97%)	0.64	45 (10%) 5 5	30, 46, 79, 118	0
All	All	822/844 (97%)	1.00	147 (17%) 1 1	30, 51, 101, 151	0

All (147) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	715	VAL	13.3
1	A	347	ASP	11.1
1	B	300	PHE	10.3
1	A	348	VAL	10.3
1	B	348	VAL	10.0
1	A	488	PRO	9.8
1	B	350	THR	8.3
1	A	355	PHE	7.5
1	A	388	ILE	7.2
1	A	391	THR	6.9
1	A	716	TRP	6.6
1	A	351	LYS	6.5
1	A	507	GLN	6.2
1	A	299	ARG	6.0
1	A	322	LEU	5.9
1	A	393	THR	5.8
1	A	489	ASP	5.8
1	A	713	THR	5.7
1	A	678	TRP	5.7
1	A	551	PHE	5.6
1	A	339	SER	5.6
1	A	352	ASP	5.4
1	A	350	THR	5.4
1	A	679	ILE	5.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	503	GLU	5.0
1	A	677	VAL	5.0
1	B	299	ARG	4.9
1	B	677	VAL	4.9
1	A	676	TRP	4.9
1	B	616	LEU	4.9
1	B	680	VAL	4.8
1	A	506	ILE	4.7
1	A	480	ILE	4.6
1	A	390	SER	4.6
1	B	352	ASP	4.6
1	A	349	ARG	4.5
1	A	479	LEU	4.3
1	A	392	SER	4.3
1	A	680	VAL	4.3
1	B	310	VAL	4.2
1	A	300	PHE	4.2
1	A	619	ARG	4.1
1	B	679	ILE	4.1
1	A	567	VAL	4.1
1	A	466	THR	4.1
1	A	491	SER	4.1
1	A	711	TRP	4.0
1	B	611	ALA	4.0
1	A	373	GLY	4.0
1	B	678	TRP	3.9
1	A	389	GLU	3.9
1	B	620	LYS	3.9
1	A	691	PHE	3.9
1	B	619	ARG	3.8
1	A	386	LYS	3.8
1	A	505	CYS	3.8
1	A	584	PHE	3.7
1	A	385	ASN	3.7
1	B	617	ASP	3.7
1	B	718	GLY	3.7
1	A	467	ASP	3.6
1	A	486	LYS	3.5
1	A	643	SER	3.5
1	B	612	LYS	3.5
1	B	309	ASP	3.5
1	A	714	HIS	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	553	TRP	3.5
1	B	567	VAL	3.5
1	A	508	GLN	3.5
1	B	691	PHE	3.5
1	A	593	ILE	3.4
1	A	611	ALA	3.3
1	A	667	ARG	3.3
1	A	681	PRO	3.3
1	A	382	GLU	3.3
1	A	566	ALA	3.3
1	A	321	THR	3.2
1	A	712	ASN	3.1
1	A	561	TRP	3.1
1	B	676	TRP	3.1
1	A	668[A]	CYS	3.1
1	A	353	GLN	3.1
1	B	615	ASP	3.0
1	A	490	GLY	3.0
1	B	562	TYR	3.0
1	A	487	GLN	3.0
1	A	588	TYR	3.0
1	B	584	PHE	2.9
1	A	469	LYS	2.9
1	A	470	HIS	2.9
1	A	354	LEU	2.8
1	A	485	TYR	2.8
1	A	499	VAL	2.8
1	A	511	LYS	2.8
1	A	478	GLN	2.8
1	A	415	CYS	2.8
1	A	565	PRO	2.8
1	A	504	ILE	2.7
1	A	591	THR	2.7
1	B	561	TRP	2.7
1	A	394	TYR	2.7
1	A	710	PRO	2.7
1	A	530	ASP	2.7
1	A	554	PHE	2.7
1	B	591	THR	2.7
1	B	681	PRO	2.7
1	B	302	LYS	2.7
1	A	500	GLN	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	566	ALA	2.6
1	A	552	ASP	2.6
1	A	416	VAL	2.6
1	A	517	PHE	2.6
1	A	609	GLU	2.5
1	A	370	LYS	2.5
1	B	715	VAL	2.5
1	B	351	LYS	2.5
1	A	465	ARG	2.5
1	B	682	PRO	2.5
1	A	446	VAL	2.4
1	A	564	LEU	2.4
1	B	353	GLN	2.4
1	A	492	THR	2.4
1	B	349	ARG	2.3
1	A	686	SER	2.3
1	B	588	TYR	2.3
1	B	608	GLU	2.3
1	A	615	ASP	2.3
1	B	311	VAL	2.3
1	A	682	PRO	2.2
1	A	685	GLY	2.2
1	B	595	VAL	2.2
1	A	620	LYS	2.2
1	A	512	ALA	2.2
1	B	318	LEU	2.2
1	B	389	GLU	2.2
1	A	590	GLY	2.2
1	B	416	VAL	2.1
1	B	593	ILE	2.1
1	A	639	TYR	2.1
1	A	658	PHE	2.1
1	B	329	HIS	2.1
1	A	514	ARG	2.0
1	A	399	THR	2.0
1	B	415	CYS	2.0
1	B	667	ARG	2.0
1	A	501	PHE	2.0
1	A	531	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)


There are no monosaccharides in this entry.


6.4 Ligands [\(i\)](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	GOL	B	805	6/6	0.50	0.27	76,80,82,83	0
7	GOL	B	806	6/6	0.69	0.29	82,83,84,85	0
4	KMV	B	803	23/23	0.88	0.35	33,95,119,122	0
4	KMV	A	803	23/23	0.90	0.33	30,87,107,108	0
3	H4B	A	802	17/17	0.95	0.18	32,36,43,43	0
3	H4B	B	802	17/17	0.96	0.20	33,36,41,41	0
5	ACT	B	804	4/4	0.96	0.11	63,67,71,76	0
2	HEM	A	801	43/43	0.97	0.20	30,38,62,81	0
2	HEM	B	801	43/43	0.97	0.18	29,36,56,61	0
5	ACT	A	804	4/4	0.97	0.11	66,70,70,71	0
6	ZN	A	805	1/1	0.99	0.06	37,37,37,37	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.