wwPDB X-ray Structure Validation Summary Report (i)

Jun 12, 2024 – 03:55 AM EDT

PDB ID	:	6NH2
Title	:	Structure of human endothelial nitric oxide synthase heme domain in complex with (R)-6-(3-fluoro-5-(2-(pyrrolidin-2-yl)ethyl)phenethyl)-4-methylpyridin-2-amine
Authors	:	Chreifi, G.; Li, H.; Poulos, T.L.
Deposited on	:	2018-12-21
Resolution	:	2.29 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36.2
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.29 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	440	4% 77% 14% • 8%
1	B	440	2% 82% 9% 9%
1	C	440	.% 80% 10% 9%
1	D	440	.% 82% 9% 9%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	BTB	C	501	-	-	X	-

2 Entry composition (i)

There are 10 unique types of molecules in this entry. The entry contains 14127 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Endothelial nitric oxide synthase splice variant eNOS13A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	404	3237	2062	570	589	16	0	2	0
1	B	402	3221	2051	566	587	17	0	3	0
1	C	401	3209	2044	563	586	16	0	2	0
1	D	402	3221	2051	566	587	17	0	3	0

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
2	A	1	43	34	1	4	4	0	0
2	B	1	43	34	1	4	4	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
2	C	1	43	34	1	4	4	0	0
			Total	C	Fe	N	O		
2	D	1	43	34	1	4	4	0	0

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula: $C_9H_{15}N_5O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	17	9	5	3	0	0
			Total	C	N	O		
3	C	1	17	9	5	3	0	0

- Molecule 4 is 6-[2-(3-fluoro-5-{2-[(2R)-pyrrolidin-2-yl]ethyl}phenyl)ethyl]-4-methylpyridin-2-amine (three-letter code: KMA) (formula: $C_{20}H_{26}FN_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	F	N		
4	A	1	24	20	1	3	0	0
			Total	C	F	N		
4	A	1	24	20	1	3	0	0
			Total	C	F	N		
4	B	1	24	20	1	3	0	0
			Total	C	F	N		
4	C	1	24	20	1	3	0	0
			Total	C	F	N		
4	C	1	24	20	1	3	0	0
			Total	C	F	N		
4	D	1	24	20	1	3	0	0

- Molecule 5 is 2-[BIS-(2-HYDROXY-ETHYL)-AMINO]-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: BTB) (formula: $C_8H_{19}NO_5$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	A	1	14	8	1	5	0	0
5	B	1	14	8	1	5	0	0
5	B	1	14	8	1	5	0	0
5	B	1	14	8	1	5	0	0
5	B	1	14	8	1	5	0	0
5	C	1	14	8	1	5	0	0
5	C	1	14	8	1	5	0	0
5	C	1	14	8	1	5	0	0
5	D	1	14	8	1	5	0	0
5	D	1	14	8	1	5	0	0
5	D	1	14	8	1	5	0	0

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
6	A	3	3	3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	C	2	Total Zn 2 2	0	0
6	D	1	Total Zn 1 1	0	0

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total C O 6 3 3	0	0
7	C	1	Total C O 6 3 3	0	0

- Molecule 8 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total Cl 1 1	0	0
8	B	1	Total Cl 1 1	0	0
8	C	1	Total Cl 1 1	0	0
8	D	1	Total Cl 1 1	0	0

- Molecule 9 is GADOLINIUM ATOM (three-letter code: GD) (formula: Gd).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Gd		
9	B	1	1	1	0	0
			Total	Gd		
9	D	1	1	1	0	0

- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
10	A	149	149	149	0	0
			Total	O		
10	B	192	192	192	0	0
			Total	O		
10	C	136	136	136	0	0
			Total	O		
10	D	234	234	234	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Endothelial nitric oxide synthase splice variant eNOS13A



• Molecule 1: Endothelial nitric oxide synthase splice variant eNOS13A



• Molecule 1: Endothelial nitric oxide synthase splice variant eNOS13A



• Molecule 1: Endothelial nitric oxide synthase splice variant eNOS13A

Chain D:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	60.33Å 153.77Å 108.94Å 90.00° 90.86° 90.00°	Depositor
Resolution (Å)	88.89 – 2.29 88.88 – 2.29	Depositor EDS
% Data completeness (in resolution range)	98.0 (88.89-2.29) 99.5 (88.88-2.29)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	0.14	Depositor
$< I/\sigma(I) >$ [1]	2.43 (at 2.29Å)	Xtriage
Refinement program	PHENIX 1.11.1-2575_1496	Depositor
R, R_{free}	0.173 , 0.222 0.166 , 0.216	Depositor DCC
R_{free} test set	4442 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å^2)	29.6	Xtriage
Anisotropy	0.560	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.33 , 51.6	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.095 for h,-k,-l	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	14127	wwPDB-VP
Average B, all atoms (Å^2)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.83% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GD, H4B, KMA, BTB, CL, ZN, HEM, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.37	0/3335	0.54	0/4543
1	B	0.41	0/3319	0.56	1/4523 (0.0%)
1	C	0.37	0/3307	0.52	0/4507
1	D	0.41	0/3319	0.56	1/4523 (0.0%)
All	All	0.39	0/13280	0.54	2/18096 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1
1	D	0	1
All	All	0	2

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	369	ASP	CB-CG-OD1	6.02	123.72	118.30
1	D	369	ASP	CB-CG-OD1	5.98	123.69	118.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	238	ARG	Peptide
1	D	89	GLN	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3237	0	3146	43	1
1	B	3221	0	3126	21	0
1	C	3209	0	3109	30	0
1	D	3221	0	3126	23	0
2	A	43	0	30	5	0
2	B	43	0	30	4	0
2	C	43	0	30	4	0
2	D	43	0	30	2	0
3	A	17	0	15	3	0
3	C	17	0	15	1	0
4	A	48	0	0	1	0
4	B	24	0	0	0	0
4	C	48	0	0	1	0
4	D	24	0	0	0	0
5	A	14	0	19	1	0
5	B	56	0	74	11	0
5	C	42	0	57	12	0
5	D	42	0	55	8	1
6	A	3	0	0	0	0
6	C	2	0	0	0	0
6	D	1	0	0	0	0
7	A	6	0	8	2	0
7	C	6	0	8	1	0
8	A	1	0	0	0	0
8	B	1	0	0	0	0
8	C	1	0	0	0	0
8	D	1	0	0	0	0
9	B	1	0	0	0	0
9	D	1	0	0	0	0
10	A	149	0	0	7	0
10	B	192	0	0	4	0
10	C	136	0	0	4	0
10	D	234	0	0	3	0
All	All	14127	0	12878	149	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 149 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:365:ARG:NH1	2:C:502:HEM:O2A	2.10	0.85
2:C:502:HEM:CGA	3:C:503:H4B:HN22	1.93	0.81
1:A:285:ARG:NH1	10:A:601:HOH:O	2.12	0.81
1:D:279:TRP:HB2	1:D:302:LEU:HD21	1.70	0.72
1:B:365:ARG:NH2	2:B:501:HEM:O1A	2.22	0.72

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:152:GLU:OE2	5:D:505:BTB:O4[2_851]	2.11	0.09

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	402/440 (91%)	382 (95%)	19 (5%)	1 (0%)	47	58
1	B	401/440 (91%)	392 (98%)	8 (2%)	1 (0%)	47	58
1	C	399/440 (91%)	385 (96%)	14 (4%)	0	100	100
1	D	401/440 (91%)	394 (98%)	7 (2%)	0	100	100
All	All	1603/1760 (91%)	1553 (97%)	48 (3%)	2 (0%)	51	64

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	283	ASN
1	B	258	ASP

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	345/373 (92%)	330 (96%)	15 (4%)	29	40
1	B	344/373 (92%)	338 (98%)	6 (2%)	60	76
1	C	342/373 (92%)	334 (98%)	8 (2%)	50	67
1	D	344/373 (92%)	340 (99%)	4 (1%)	71	84
All	All	1375/1492 (92%)	1342 (98%)	33 (2%)	49	66

5 of 33 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	470	SER
1	D	89	GLN
1	D	432	GLU
1	A	301	GLU
1	A	285	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 37 ligands modelled in this entry, 12 are monoatomic - leaving 25 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	KMA	D	503	-	25,26,26	0.41	0	32,35,35	1.46	6 (18%)
5	BTB	C	501	-	13,13,13	1.95	2 (15%)	7,16,16	0.79	0
5	BTB	D	505	-	13,13,13	0.72	0	7,16,16	1.08	1 (14%)
2	HEM	C	502	1	41,50,50	1.49	6 (14%)	45,82,82	1.64	12 (26%)
3	H4B	C	503	-	16,18,18	0.68	0	11,26,26	2.88	7 (63%)
5	BTB	D	504	9	13,13,13	0.39	0	7,16,16	0.68	0
5	BTB	B	503	9	13,13,13	0.40	0	7,16,16	0.47	0
3	H4B	A	502	-	16,18,18	0.72	0	11,26,26	2.59	6 (54%)
5	BTB	D	506	-	13,13,13	0.42	0	7,16,16	0.83	0
2	HEM	A	501	1	41,50,50	1.50	6 (14%)	45,82,82	1.89	13 (28%)
4	KMA	A	509	-	25,26,26	0.44	0	32,35,35	1.37	5 (15%)
5	BTB	B	508	-	13,13,13	0.94	2 (15%)	7,16,16	1.19	1 (14%)
4	KMA	C	504	-	25,26,26	0.41	0	32,35,35	1.52	6 (18%)
2	HEM	B	501	1	41,50,50	1.47	6 (14%)	45,82,82	1.69	10 (22%)
7	GOL	A	507	-	5,5,5	0.37	0	5,5,5	0.23	0
2	HEM	D	502	1	41,50,50	1.44	8 (19%)	45,82,82	1.68	9 (20%)
7	GOL	C	508	-	5,5,5	0.41	0	5,5,5	0.43	0
5	BTB	B	504	-	13,13,13	0.63	0	7,16,16	0.81	0
4	KMA	B	502	-	25,26,26	0.45	0	32,35,35	1.50	8 (25%)
5	BTB	B	505	-	13,13,13	0.36	0	7,16,16	0.48	0
4	KMA	C	510	-	25,26,26	0.41	0	32,35,35	1.79	6 (18%)
4	KMA	A	503	-	25,26,26	0.37	0	32,35,35	1.56	6 (18%)
5	BTB	A	504	-	13,13,13	0.65	0	7,16,16	0.92	0
5	BTB	C	506	-	13,13,13	0.36	0	7,16,16	0.35	0
5	BTB	C	505	-	13,13,13	1.56	2 (15%)	7,16,16	1.37	2 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	KMA	D	503	-	-	1/10/17/17	0/3/3/3
5	BTB	C	501	-	-	13/21/21/21	-
5	BTB	D	505	-	-	9/21/21/21	-
2	HEM	C	502	1	-	2/12/54/54	-
3	H4B	C	503	-	-	3/8/17/17	0/2/2/2
5	BTB	D	504	9	-	4/21/21/21	-
5	BTB	B	503	9	-	3/21/21/21	-
3	H4B	A	502	-	-	3/8/17/17	0/2/2/2
5	BTB	D	506	-	-	6/21/21/21	-
2	HEM	A	501	1	-	4/12/54/54	-
4	KMA	A	509	-	-	1/10/17/17	0/3/3/3
5	BTB	B	508	-	-	12/21/21/21	-
4	KMA	C	504	-	-	2/10/17/17	0/3/3/3
2	HEM	B	501	1	-	3/12/54/54	-
7	GOL	A	507	-	-	2/4/4/4	-
2	HEM	D	502	1	-	4/12/54/54	-
7	GOL	C	508	-	-	2/4/4/4	-
5	BTB	B	504	-	-	3/21/21/21	-
4	KMA	B	502	-	-	1/10/17/17	0/3/3/3
5	BTB	B	505	-	-	8/21/21/21	-
4	KMA	C	510	-	-	1/10/17/17	0/3/3/3
4	KMA	A	503	-	-	1/10/17/17	0/3/3/3
5	BTB	A	504	-	-	6/21/21/21	-
5	BTB	C	506	-	-	2/21/21/21	-
5	BTB	C	505	-	-	10/21/21/21	-

The worst 5 of 32 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	501	BTB	C5-N	-5.68	1.39	1.48
5	C	505	BTB	C5-N	-4.72	1.41	1.48
2	A	501	HEM	C3C-C2C	-3.94	1.34	1.40
2	C	502	HEM	C3C-C2C	-3.75	1.35	1.40
2	C	502	HEM	C3C-CAC	3.69	1.55	1.47

The worst 5 of 98 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	A	501	HEM	CMA-C3A-C4A	-5.35	120.25	128.46
3	C	503	H4B	C8A-C4A-C4	4.95	118.97	114.57
2	A	501	HEM	CBD-CAD-C3D	-4.82	99.22	112.63
3	C	503	H4B	C4-C4A-N5	4.57	122.96	119.12
3	A	502	H4B	C4-C4A-N5	4.50	122.90	119.12

There are no chirality outliers.

5 of 106 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	HEM	C1A-C2A-CAA-CBA
2	A	501	HEM	C3A-C2A-CAA-CBA
2	B	501	HEM	C1A-C2A-CAA-CBA
2	B	501	HEM	C3A-C2A-CAA-CBA
2	D	502	HEM	C1A-C2A-CAA-CBA

There are no ring outliers.

21 monomers are involved in 54 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	501	BTB	7	0
5	D	505	BTB	1	1
2	C	502	HEM	4	0
3	C	503	H4B	1	0
5	D	504	BTB	2	0
5	B	503	BTB	3	0
3	A	502	H4B	3	0
5	D	506	BTB	5	0
2	A	501	HEM	5	0
5	B	508	BTB	1	0
4	C	504	KMA	1	0
2	B	501	HEM	4	0
7	A	507	GOL	2	0
2	D	502	HEM	2	0
7	C	508	GOL	1	0
5	B	504	BTB	4	0
5	B	505	BTB	3	0
4	A	503	KMA	1	0
5	A	504	BTB	1	0
5	C	506	BTB	1	0
5	C	505	BTB	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand KMA D 503



Ligand HEM C 502



Bond lengths



Bond angles



Torsions



Rings

Ligand HEM A 501



Bond lengths



Bond angles



Torsions



Rings

Ligand KMA A 509



Ligand KMA C 504



Ligand HEM B 501



Bond lengths



Bond angles



Torsions



Rings

Ligand HEM D 502



Bond lengths



Bond angles



Torsions



Rings

Ligand KMA B 502



Ligand KMA C 510



Ligand KMA A 503



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	404/440 (91%)	-0.04	16 (3%)	38	45	20, 39, 84, 120	0
1	B	402/440 (91%)	-0.19	7 (1%)	70	76	16, 29, 62, 129	0
1	C	401/440 (91%)	-0.11	6 (1%)	73	79	20, 38, 78, 113	0
1	D	402/440 (91%)	-0.23	4 (0%)	82	86	15, 28, 57, 103	0
All	All	1609/1760 (91%)	-0.14	33 (2%)	63	70	15, 33, 75, 129	0

The worst 5 of 33 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	119	ALA	7.2
1	A	144	GLN	5.4
1	A	109	LEU	5.1
1	B	257	GLN	5.0
1	A	142	GLY	5.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
5	BTB	C	506	14/14	0.44	0.31	75,103,117,117	0
5	BTB	D	504	14/14	0.79	0.20	36,57,65,70	0
7	GOL	A	507	6/6	0.81	0.12	52,66,78,80	0
5	BTB	D	506	14/14	0.82	0.17	74,81,85,88	0
3	H4B	C	503	17/17	0.83	0.27	55,66,80,81	0
5	BTB	B	505	14/14	0.83	0.16	51,71,78,83	0
4	KMA	A	509	24/24	0.85	0.19	40,59,76,77	0
3	H4B	A	502	17/17	0.87	0.26	62,71,81,91	0
7	GOL	C	508	6/6	0.87	0.13	34,54,66,68	0
5	BTB	B	504	14/14	0.88	0.16	32,62,74,78	0
5	BTB	C	505	14/14	0.88	0.28	20,58,71,78	0
5	BTB	D	505	14/14	0.88	0.20	42,62,78,82	0
5	BTB	C	501	14/14	0.89	0.30	10,56,68,69	0
4	KMA	B	502	24/24	0.89	0.21	40,61,67,71	0
4	KMA	C	510	24/24	0.90	0.16	40,54,73,76	0
5	BTB	A	504	14/14	0.90	0.17	45,64,67,77	0
4	KMA	A	503	24/24	0.90	0.18	31,61,77,79	0
4	KMA	C	504	24/24	0.90	0.15	29,53,77,79	0
4	KMA	D	503	24/24	0.91	0.21	40,57,68,71	0
5	BTB	B	503	14/14	0.91	0.17	28,46,72,78	0
5	BTB	B	508	14/14	0.93	0.19	14,49,58,61	0
6	ZN	A	506	1/1	0.96	0.10	52,52,52,52	1
2	HEM	A	501	43/43	0.97	0.14	18,32,53,70	0
6	ZN	C	511	1/1	0.97	0.11	28,28,28,28	1
8	CL	C	509	1/1	0.97	0.06	51,51,51,51	0
2	HEM	D	502	43/43	0.98	0.12	8,18,52,78	0
2	HEM	B	501	43/43	0.98	0.13	7,18,46,75	0
2	HEM	C	502	43/43	0.98	0.14	19,32,66,80	0
8	CL	A	508	1/1	0.98	0.09	53,53,53,53	0
8	CL	B	506	1/1	0.98	0.12	31,31,31,31	0
6	ZN	C	507	1/1	0.98	0.08	42,42,42,42	1
8	CL	D	508	1/1	0.98	0.10	31,31,31,31	0
6	ZN	A	505	1/1	0.99	0.10	32,32,32,32	0
6	ZN	A	510	1/1	0.99	0.10	27,27,27,27	1
9	GD	B	507	1/1	0.99	0.14	35,35,35,35	0
9	GD	D	507	1/1	0.99	0.14	39,39,39,39	0
6	ZN	D	501	1/1	1.00	0.10	27,27,27,27	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around KMA A 509:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around KMA B 502:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around KMA C 510:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around KMA A 503:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around KMA C 504:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around KMA D 503:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM A 501:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around HEM D 502:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM B 501:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around HEM C 502:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







6.5 Other polymers (i)

There are no such residues in this entry.