Full wwPDB X-ray Structure Validation Report (i)

Sep 12, 2023 – 08:07 AM EDT

PDB ID	:	4O2P
Title	:	Kinase domain of cSrc in complex with a substituted pyrazolopyrimidine
Authors	:	Richters, A.; Rauh, D.
Deposited on	:	2013-12-17
Resolution	:	2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	286	3% 83% 9% • 7%
1	B	286	7% 87% 6% 7%

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 4561 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proto-oncogene tyrosine-protein kinase Src.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	265	2134	1371	356	391	16	0	2	0
1	B	265	2123	1363	353	391	16	0	1	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	248	GLY	-	expression tag	UNP P00523
A	249	HIS	-	expression tag	UNP P00523
A	250	MET	-	expression tag	UNP P00523
B	248	GLY	-	expression tag	UNP P00523
B	249	HIS	-	expression tag	UNP P00523
B	250	MET	-	expression tag	UNP P00523

- Molecule 2 is 1-[(2R)-2-chloro-2-phenylethyl]-6-{[2-(morpholin-4-yl)ethyl]sulfanyl}-N-phenyl-1H-pyrazolo[3,4-d]pyrimidin-4-amine (three-letter code: 11V) (formula: $C_{25}H_{27}ClN_6OS$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	Cl	N	O	S		
2	A	1	34	25	1	6	1	1	0	0
			Total	C	Cl	N	O	S		
2	B	1	34	25	1	6	1	1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	119	119	119	0	0
			Total	O		
3	B	117	117	117	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Proto-oncogene tyrosine-protein kinase Src

Chain A:



• Molecule 1: Proto-oncogene tyrosine-protein kinase Src

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α, β, γ	42.22Å 63.25Å 74.81Å 101.35° 90.44° 90.07°	Depositor
Resolution (Å)	43.34 – 2.10 43.34 – 2.10	Depositor EDS
% Data completeness (in resolution range)	96.1 (43.34-2.10) 96.1 (43.34-2.10)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.36 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.6.0093	Depositor
R, R_{free}	0.197 , 0.238 0.196 , 0.238	Depositor DCC
R_{free} test set	851 reflections (2.00%)	wwPDB-VP
Wilson B-factor (Å^2)	29.6	Xtriage
Anisotropy	0.082	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.34 , 44.8	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	0.026 for h,-k,-l	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	4561	wwPDB-VP
Average B, all atoms (Å^2)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.88% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: 11V

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.47	0/2191	0.54	0/2966
1	B	0.49	0/2176	0.57	0/2947
All	All	0.48	0/4367	0.55	0/5913

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2134	0	2108	13	0
1	B	2123	0	2103	7	0
2	A	34	0	26	0	0
2	B	34	0	26	1	0
3	A	119	0	0	0	0
3	B	117	0	0	2	0
All	All	4561	0	4263	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:257:LYS:HB3	1:B:261:GLU:HG3	1.52	0.91
1:B:493[B]:ASP:OD2	1:B:497:GLN:NE2	2.36	0.59
1:B:261:GLU:HG2	3:B:762:HOH:O	2.04	0.58
1:B:273:LEU:O	2:B:601:11V:H5	2.12	0.50
1:A:356:LYS:HD3	1:A:357:TYR:CE1	2.50	0.47
1:B:297:LEU:HD21	1:B:302:MET:HB3	1.96	0.47
1:A:281:VAL:HG22	1:A:295:LYS:HG2	1.99	0.45
1:A:310:GLU:HG3	1:A:410:LEU:HD22	1.98	0.45
1:A:378:GLU:HG3	1:A:441:ILE:HG12	1.99	0.45
1:A:323:VAL:HG12	1:A:403:ALA:HA	1.99	0.44
1:A:273:LEU:HD11	1:A:283:MET:HB3	2.00	0.44
1:A:260:TRP:CD1	1:A:315:LYS:HD3	2.53	0.44
1:A:261:GLU:OE1	1:A:330:SER:OG	2.35	0.43
1:A:317:LEU:HD21	1:A:382:TYR:CE1	2.54	0.42
1:A:298:LYS:HA	1:A:299:PRO:HD3	1.93	0.42
1:A:320[A]:GLU:HG3	1:A:321:LYS:HG2	2.02	0.42
1:B:319:HIS:HB3	1:B:322:LEU:HG	2.02	0.41
1:A:426:ILE:HD13	1:A:472:LEU:HG	2.03	0.41
1:B:411:ILE:HD13	3:B:792:HOH:O	2.21	0.40
1:A:424:PHE:HA	1:A:425:PRO:HD3	2.00	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	261/286 (91%)	253 (97%)	7 (3%)	1 (0%)	34	32
1	B	260/286 (91%)	251 (96%)	8 (3%)	1 (0%)	34	32
All	All	521/572 (91%)	504 (97%)	15 (3%)	2 (0%)	34	32

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	304	PRO
1	B	299	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	228/245 (93%)	223 (98%)	5 (2%)	52	57
1	B	227/245 (93%)	220 (97%)	7 (3%)	40	43
All	All	455/490 (93%)	443 (97%)	12 (3%)	44	50

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	283	MET
1	A	291	ARG
1	A	356	LYS
1	A	409	ARG
1	A	424	PHE
1	B	264	ARG
1	B	275	GLN
1	B	297	LEU
1	B	388	ARG
1	B	432	GLU
1	B	458	LYS
1	B	504	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	275	GLN
1	B	526	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	11V	B	601	-	35,38,38	2.23	5 (14%)	41,51,51	1.92	6 (14%)
2	11V	A	601	-	35,38,38	2.24	6 (17%)	41,51,51	1.75	7 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	11V	B	601	-	-	4/16/26/26	0/5/5/5
2	11V	A	601	-	-	6/16/26/26	0/5/5/5

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	11V	C6-C5	-8.04	1.34	1.44

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	11V	C6-C5	-7.95	1.35	1.44
2	B	601	11V	CBA-CBF	-7.81	1.41	1.51
2	A	601	11V	CBA-CBF	-7.78	1.41	1.51
2	B	601	11V	C5-C4	-3.84	1.33	1.43
2	B	601	11V	CAL-C5	-3.41	1.33	1.40
2	A	601	11V	C5-C4	-3.39	1.34	1.43
2	A	601	11V	CAL-C5	-3.28	1.34	1.40
2	B	601	11V	CAZ-NAW	-2.84	1.34	1.40
2	A	601	11V	C2-SAY	-2.84	1.70	1.75
2	A	601	11V	CAZ-NAW	-2.43	1.35	1.40

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	B	601	11V	CAS-NBH-NAT	7.05	130.31	117.08
2	A	601	11V	CAS-NBH-NAT	6.09	128.51	117.08
2	B	601	11V	CAL-NAT-NBH	4.60	109.02	104.23
2	B	601	11V	CAZ-NAW-C6	-4.41	116.78	128.26
2	A	601	11V	CAL-NAT-NBH	4.08	108.48	104.23
2	B	601	11V	N3-C2-N1	-4.04	119.55	126.98
2	A	601	11V	CAZ-NAW-C6	-3.99	117.87	128.26
2	A	601	11V	N3-C2-N1	-3.64	120.28	126.98
2	B	601	11V	C2-N3-C4	3.62	120.31	115.32
2	A	601	11V	C2-N3-C4	3.25	119.80	115.32
2	B	601	11V	CAQ-NBG-CAP	2.64	114.77	108.83
2	A	601	11V	CAO-SAY-C2	-2.26	99.82	102.90
2	A	601	11V	CAO-CAR-NBG	-2.04	107.80	112.76

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	601	11V	CAO-CAR-NBG-CAQ
2	A	601	11V	N1-C6-NAW-CAZ
2	A	601	11V	CAH-CAZ-NAW-C6
2	A	601	11V	CAI-CAZ-NAW-C6
2	B	601	11V	CAH-CAZ-NAW-C6
2	B	601	11V	CAI-CAZ-NAW-C6
2	A	601	11V	C5-C6-NAW-CAZ
2	A	601	11V	N1-C2-SAY-CAO
2	B	601	11V	CAO-CAR-NBG-CAP
2	A	601	11V	N3-C2-SAY-CAO

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	601	11V	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand 11V B 601





5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	265/286 (92%)	0.06	8 (3%)	50	56	16, 33, 74, 91	0
1	B	265/286 (92%)	0.12	19 (7%)	15	19	17, 32, 76, 93	0
All	All	530/572 (92%)	0.09	27 (5%)	28	33	16, 33, 75, 93	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	299	PRO	5.4
1	B	301	THR	4.9
1	B	300	GLY	4.3
1	A	313	VAL	3.6
1	B	306	ALA	3.5
1	B	297	LEU	3.1
1	A	424	PHE	3.1
1	A	312	GLN	3.1
1	A	277	CYS	3.0
1	B	303	SER	2.9
1	A	306	ALA	2.9
1	A	273	LEU	2.6
1	B	267	LEU	2.6
1	A	278	PHE	2.5
1	B	329	VAL	2.4
1	A	305	GLU	2.3
1	B	305	GLU	2.3
1	B	333	PRO	2.3
1	B	332	GLU	2.3
1	B	331	GLU	2.2
1	B	334	ILE	2.2
1	B	271	VAL	2.2
1	B	269	LEU	2.1
1	B	296	THR	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	335	TYR	2.1
1	B	282	TRP	2.0
1	B	302	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	11V	B	601	34/34	0.88	0.17	35,39,67,70	34
2	11V	A	601	34/34	0.90	0.16	27,33,55,56	34

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 11V B 601:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 11V A 601:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.