Full wwPDB X-ray Structure Validation Report (i)

Jun 17, 2024 – 08:56 AM EDT

PDB ID	:	3OBR
Title	:	Crystal structure of Botulinum neurotoxin serotype D binding domain
Authors	:	Zong, Y.; Lee, K.K.; Jin, R.
Deposited on	:	2010-08-09
Resolution	:	1.72 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.72 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5722 (1.74-1.70)
Clashscore	141614	6152 (1.74-1.70)
Ramachandran outliers	138981	6051 (1.74-1.70)
Sidechain outliers	138945	6051 (1.74-1.70)
RSRZ outliers	127900	5629 (1.74-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	434	6% 80% 16% •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	3	-	-	X	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	6	-	-	X	X
2	GOL	A	8	-	-	X	-
2	GOL	A	9	-	-	-	X

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 4058 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Botulinum neurotoxin type D.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	418	3483	2227	573	670	13	0	9	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	856	MET	-	EXPRESSION TAG	UNP P19321
A	857	ARG	-	EXPRESSION TAG	UNP P19321
A	858	GLY	-	EXPRESSION TAG	UNP P19321
A	859	SER	-	EXPRESSION TAG	UNP P19321
A	860	ALA	-	EXPRESSION TAG	UNP P19321
A	861	MET	-	EXPRESSION TAG	UNP P19321
A	862	ALA	-	EXPRESSION TAG	UNP P19321
A	1277	PRO	-	EXPRESSION TAG	UNP P19321
A	1278	PRO	-	EXPRESSION TAG	UNP P19321
A	1279	THR	-	EXPRESSION TAG	UNP P19321
A	1280	PRO	-	EXPRESSION TAG	UNP P19321
A	1281	GLY	-	EXPRESSION TAG	UNP P19321
A	1282	TRP	-	EXPRESSION TAG	UNP P19321
A	1283	SER	-	EXPRESSION TAG	UNP P19321
A	1284	HIS	-	EXPRESSION TAG	UNP P19321
A	1285	PRO	-	EXPRESSION TAG	UNP P19321
A	1286	GLN	-	EXPRESSION TAG	UNP P19321
A	1287	PHE	-	EXPRESSION TAG	UNP P19321
A	1288	GLU	-	EXPRESSION TAG	UNP P19321
A	1289	LYS	-	EXPRESSION TAG	UNP P19321

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	6	3	3	0	0
2	A	1	6	3	3	0	0
2	A	1	6	3	3	0	0
2	A	1	6	3	3	0	0
2	A	1	6	3	3	0	0
2	A	1	6	3	3	0	0
2	A	1	6	3	3	0	0
2	A	1	6	3	3	0	0
2	A	1	6	3	3	0	0
2	A	1	6	3	3	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	515	515	515	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Botulinum neurotoxin type D

Chain A:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	60.68Å 89.87Å 94.12Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.72 41.69 – 1.72	Depositor EDS
% Data completeness (in resolution range)	99.9 (50.00-1.72) 99.8 (41.69-1.72)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.45 (at 1.72Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.126 , 0.181 0.170 , 0.190	Depositor DCC
R_{free} test set	2795 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å^2)	13.8	Xtriage
Anisotropy	0.213	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.34 , 51.5	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	0.012 for -h,l,k	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	4058	wwPDB-VP
Average B, all atoms (Å^2)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.39% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.83	2/3563 (0.1%)	0.70	1/4828 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1047	GLU	CB-CG	5.30	1.62	1.52
1	A	1130	TYR	CE1-CZ	5.16	1.45	1.38

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	904	ASP	CB-CG-OD1	5.10	122.89	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3483	0	3427	70	1
2	A	60	0	79	21	0
3	A	515	0	0	21	3
All	All	4058	0	3506	74	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1178:THR:CB	2:A:6:GOL:H12	1.80	1.10
1:A:994:SER:OG	1:A:996:THR:HG22	1.49	1.10
1:A:1178:THR:HB	2:A:6:GOL:H12	1.08	1.06
1:A:1178:THR:HB	2:A:6:GOL:C1	1.89	1.02
1:A:1211:VAL:HG13	2:A:8:GOL:H12	1.49	0.95
1:A:965[B]:CYS:SG	1:A:972:GLU:HB3	2.16	0.85
1:A:1270:ARG:NH2	3:A:278:HOH:O	2.10	0.83
1:A:1183:CYS:HB2	3:A:456:HOH:O	1.85	0.76
2:A:8:GOL:H11	3:A:254:HOH:O	1.85	0.76
1:A:963:LYS:NZ	2:A:5:GOL:H11	2.01	0.76
1:A:862:ALA:HA	3:A:510:HOH:O	1.89	0.71
1:A:1251:VAL:HB	2:A:9:GOL:H32	1.72	0.71
1:A:1175:ILE:HG22	1:A:1176:TYR:CD1	2.28	0.69
1:A:966[A]:ILE:CG2	3:A:338:HOH:O	2.42	0.68
1:A:900:ILE:HD11	3:A:449:HOH:O	1.95	0.67
1:A:950:TYR:HB2	1:A:966[B]:ILE:CG1	2.25	0.66
1:A:1141[A]:SER:O	1:A:1142:ASP:HB2	1.95	0.66
1:A:1175:ILE:HG22	1:A:1176:TYR:CE1	2.31	0.66
2:A:8:GOL:H32	3:A:296:HOH:O	1.95	0.65
1:A:1009:ASN:ND2	1:A:1015:MET:HG3	2.13	0.63
1:A:963:LYS:HZ3	2:A:5:GOL:H11	1.61	0.62
1:A:1083[B]:ASN:OD1	3:A:323:HOH:O	2.16	0.61
1:A:1010:ASN:O	1:A:1013:GLY:N	2.34	0.60
1:A:1179:GLN:HB3	1:A:1182:GLU:N	2.17	0.59
1:A:1178:THR:HG22	1:A:1178:THR:O	2.03	0.58
1:A:936:TRP:HB2	1:A:1058:ARG:HG2	1.86	0.57
1:A:1178:THR:O	1:A:1179:GLN:HB2	2.05	0.56
1:A:1211:VAL:HG13	2:A:8:GOL:C1	2.30	0.56
1:A:998:TYR:OH	2:A:3:GOL:H12	2.06	0.56
1:A:950:TYR:HB2	1:A:966[B]:ILE:HG12	1.87	0.55
1:A:1208:LYS:HE2	3:A:223:HOH:O	2.06	0.55
1:A:1025:GLN:CG	3:A:14:HOH:O	2.55	0.55
1:A:1176:TYR:N	3:A:492:HOH:O	2.16	0.53
1:A:959:ASN:HD22	1:A:980:ARG:HE	1.58	0.52
1:A:962:TRP:HA	1:A:974:ILE:O	2.10	0.52
1:A:1178:THR:OG1	2:A:6:GOL:H12	2.10	0.52
1:A:900:ILE:CD1	3:A:449:HOH:O	2.56	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:927:ILE:HD12	1:A:1011:ILE:HG22	1.93	0.51
1:A:860:ALA:O	1:A:864:ILE:HG13	2.10	0.50
1:A:1126:ARG:HD3	3:A:317:HOH:O	2.11	0.50
1:A:959:ASN:ND2	1:A:980:ARG:HE	2.09	0.50
1:A:1020:ASN:HB2	2:A:3:GOL:H11	1.93	0.50
1:A:1024:LYS:HE3	2:A:3:GOL:O3	2.11	0.50
1:A:994:SER:OG	1:A:996:THR:CG2	2.42	0.49
1:A:1075:VAL:HG22	2:A:7:GOL:H32	1.94	0.48
1:A:1175:ILE:CG2	1:A:1176:TYR:CE1	2.96	0.47
1:A:1009:ASN:HD22	1:A:1015:MET:HG3	1.78	0.47
1:A:1025:GLN:HG3	3:A:14:HOH:O	2.13	0.47
1:A:971:ILE:O	1:A:971:ILE:HG13	2.15	0.47
2:A:6:GOL:H32	3:A:492:HOH:O	2.15	0.47
1:A:861:MET:HE3	2:A:7:GOL:O3	2.14	0.47
1:A:1252:THR:O	2:A:9:GOL:H12	2.15	0.46
1:A:1126:ARG:NH2	3:A:458:HOH:O	2.32	0.46
2:A:6:GOL:C3	3:A:216:HOH:O	2.64	0.46
1:A:927:ILE:HD12	1:A:1011:ILE:CG2	2.46	0.46
1:A:1117:VAL:HG11	1:A:1237:PRO:HB2	1.97	0.46
1:A:932:SER:HA	1:A:1007:ILE:O	2.16	0.45
1:A:938:LYS:HD2	1:A:1002:TRP:CD1	2.50	0.45
1:A:966[A]:ILE:HG22	3:A:338:HOH:O	2.12	0.45
1:A:1006:THR:HG21	1:A:1072:ILE:HG12	1.99	0.44
1:A:1030:GLU:HG2	3:A:419:HOH:O	2.19	0.43
1:A:943:LEU:HG	1:A:966[B]:ILE:HD12	2.01	0.43
1:A:1138[B]:LYS:HG2	1:A:1139[B]:SER:N	2.34	0.42
1:A:1020:ASN:HD22	2:A:3:GOL:C1	2.32	0.42
1:A:995[A]:HIS:NE2	1:A:1266:LYS:NZ	2.60	0.42
1:A:938:LYS:HD2	1:A:1002:TRP:NE1	2.34	0.42
1:A:1010:ASN:O	1:A:1012:MET:N	2.52	0.42
1:A:1172:THR:HG23	3:A:113:HOH:O	2.19	0.42
1:A:861:MET:HE3	1:A:865:ASN:HD21	1.85	0.41
1:A:966[A]:ILE:HG23	3:A:338:HOH:O	2.16	0.41
1:A:861:MET:CE	1:A:865:ASN:HD21	2.34	0.41
1:A:1119:VAL:HG21	2:A:1:GOL:H31	2.03	0.41
1:A:1234:ILE:HG23	1:A:1234:ILE:O	2.21	0.40
1:A:1099:TYR:CE1	1:A:1269:SER:HB3	2.56	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:153:HOH:O	3:A:497:HOH:O[2_565]	1.92	0.28
1:A:1030:GLU:OE1	3:A:369:HOH:O[3_645]	2.14	0.06
3:A:86:HOH:O	3:A:496:HOH:O[2_665]	2.15	0.05

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	423/434 (98%)	408 (96%)	14 (3%)	1 (0%)	47 30

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1011	ILE

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	399/402 (99%)	398 (100%)	1 (0%)	92 89

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1197	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	865	ASN
1	A	959	ASN
1	A	1009	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	GOL	A	9	-	5,5,5	0.31	0	5,5,5	0.35	0
2	GOL	A	3	-	5,5,5	1.03	0	5,5,5	1.57	0
2	GOL	A	6	-	5,5,5	0.41	0	5,5,5	0.70	0
2	GOL	A	1	-	5,5,5	0.44	0	5,5,5	0.55	0
2	GOL	A	8	-	5,5,5	0.35	0	5,5,5	0.88	0
2	GOL	A	4	-	5,5,5	1.07	0	5,5,5	1.12	0
2	GOL	A	10	-	5,5,5	0.85	0	5,5,5	1.30	0
2	GOL	A	7	-	5,5,5	0.54	0	5,5,5	1.25	1 (20%)
2	GOL	A	5	-	5,5,5	0.48	0	5,5,5	0.43	0
2	GOL	A	2	-	5,5,5	0.50	0	5,5,5	0.75	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	9	-	-	2/4/4/4	-
2	GOL	A	3	-	-	2/4/4/4	-
2	GOL	A	6	-	-	1/4/4/4	-
2	GOL	A	1	-	-	3/4/4/4	-
2	GOL	A	8	-	-	2/4/4/4	-
2	GOL	A	4	-	-	1/4/4/4	-
2	GOL	A	10	-	-	4/4/4/4	-
2	GOL	A	7	-	-	3/4/4/4	-
2	GOL	A	5	-	-	2/4/4/4	-
2	GOL	A	2	-	-	0/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	A	7	GOL	O2-C2-C3	2.23	118.40	109.18

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1	GOL	O1-C1-C2-O2
2	A	1	GOL	O1-C1-C2-C3
2	A	3	GOL	O1-C1-C2-C3
2	A	5	GOL	O1-C1-C2-C3
2	A	7	GOL	C1-C2-C3-O3
2	A	8	GOL	O1-C1-C2-C3
2	A	10	GOL	O1-C1-C2-C3
2	A	10	GOL	C1-C2-C3-O3
2	A	10	GOL	O2-C2-C3-O3
2	A	6	GOL	C1-C2-C3-O3
2	A	9	GOL	C1-C2-C3-O3
2	A	3	GOL	O1-C1-C2-O2
2	A	10	GOL	O1-C1-C2-O2
2	A	7	GOL	O2-C2-C3-O3
2	A	5	GOL	O1-C1-C2-O2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	9	GOL	O2-C2-C3-O3
2	A	7	GOL	O1-C1-C2-O2
2	A	8	GOL	O1-C1-C2-O2
2	A	1	GOL	O2-C2-C3-O3
2	A	4	GOL	O1-C1-C2-O2

There are no ring outliers.

7 monomers are involved in 21 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	9	GOL	2	0
2	A	3	GOL	4	0
2	A	6	GOL	6	0
2	A	1	GOL	1	0
2	A	8	GOL	4	0
2	A	7	GOL	2	0
2	A	5	GOL	2	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	418/434 (96%)	0.18	26 (6%) 20 23	6, 14, 35, 61	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1178	THR	11.0
1	A	1183	CYS	10.2
1	A	1179	GLN	9.3
1	A	1177	ALA	7.9
1	A	1184	SER	7.5
1	A	924	TYR	7.3
1	A	860	ALA	7.0
1	A	923	LEU	6.9
1	A	1176	TYR	6.8
1	A	922	ILE	6.7
1	A	1182	GLU	6.7
1	A	1278	PRO	5.6
1	A	925	SER	5.3
1	A	1242	PHE	5.3
1	A	859	SER	5.2
1	A	1185	GLN	3.7
1	A	1277	PRO	3.6
1	A	861	MET	3.0
1	A	1186	ASN	2.9
1	A	1175	ILE	2.9
1	A	921	ASN	2.7
1	A	958	GLN	2.6
1	A	928	TYR	2.5
1	A	926	ALA	2.2
1	A	1144	ASN	2.1
1	A	995[A]	HIS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	GOL	A	7	6/6	0.62	0.30	44,55,55,56	0
2	GOL	A	6	6/6	0.64	0.41	59,65,66,66	0
2	GOL	A	9	6/6	0.69	0.41	69,71,71,72	0
2	GOL	A	5	6/6	0.77	0.25	47,55,56,56	0
2	GOL	A	10	6/6	0.80	0.18	45,45,46,47	0
2	GOL	A	1	6/6	0.81	0.16	31,40,42,42	0
2	GOL	A	8	6/6	0.86	0.22	24,38,41,41	0
2	GOL	A	3	6/6	0.89	0.30	22,31,33,34	0
2	GOL	A	2	6/6	0.90	0.21	56,62,64,65	0
2	GOL	A	4	6/6	0.91	0.16	23,35,39,40	0

6.5 Other polymers (i)

There are no such residues in this entry.