wwPDB X-ray Structure Validation Summary Report (i)

Sep 12, 2023 – 08:03 PM EDT

PDB ID	:	4ODW
Title	:	Unliganded Fab structure of lipid A-specific antibody A6
Authors	:	Haji-Ghassemi, O.; Evans, S.V.
Deposited on	:	2014-01-10
Resolution	:	2.72 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.72 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3359 (2.74-2.70)
Clashscore	141614	3686 (2.74-2.70)
Ramachandran outliers	138981	3622 (2.74-2.70)
Sidechain outliers	138945	3623 (2.74-2.70)
RSRZ outliers	127900	3276 (2.74-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	215	7%, 70%, 25%, 5%
1	L	215	2%, 70%, 23%, 6%
2	A	222	5%, 72%, 20%, 6%
2	H	222	2%, 74%, 20%

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 6558 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called A6 Fab (IgG2b kappa) light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	L	214	1669	1039	282	343	5	0	0	0
1	B	214	1669	1039	282	343	5	0	0	0

- Molecule 2 is a protein called A6 Fab (IgG2b) heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	214	1618	1027	264	318	9	0	0	0
2	A	208	1578	1003	258	309	8	0	0	0

- Molecule 3 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: $C_4H_{12}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	8	4	1	3	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	L	4	4	4	0	0
			Total	O		
4	H	4	4	4	0	0
			Total	O		
4	B	5	5	5	0	0
			Total	O		
4	A	3	3	3	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: A6 Fab (IgG2b kappa) light chain

Chain L:



• Molecule 1: A6 Fab (IgG2b kappa) light chain

Chain B:



• Molecule 2: A6 Fab (IgG2b) heavy chain

Chain H:



• Molecule 2: A6 Fab (IgG2b) heavy chain

Chain A:

5%
72%
20%
6%


GLU V2 K3 V12 K13 L20 Y33 M34 S35 T40 P41 E42 K43 S52A Y58 Y59 P60 D61 T62 V63 K64 L80 S82A K83 S84 E85 L95 Y96 G97 N98 Y99 V100 R100A I100B H100C T100D M100E Q105 G106 T107 S112 Y122 P123 L124 A125 P126 GLY
CYS GLY ASP THR THR G132 L137 G138 C139 E147 S148 W153 N154 S155 GLY SER LEU SER S160 S161 L168 L169 Q170 V182 P183 S184 S185 T186 W187 E190 T191 V192 K208 L209 GLU PRO SER

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α, β, γ	42.13Å 69.23Å 69.07Å 70.99° 72.39° 88.29°	Depositor
Resolution (Å)	65.27 – 2.72 33.90 – 2.72	Depositor EDS
% Data completeness (in resolution range)	86.6 (65.27-2.72) 86.7 (33.90-2.72)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.53 (at 2.72Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.231 , 0.257 0.233 , 0.257	Depositor DCC
R_{free} test set	833 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å^2)	40.5	Xtriage
Anisotropy	0.137	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.32 , 43.7	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.035 for h,-k,h-l	Xtriage
F_o,F_c correlation	0.90	EDS
Total number of atoms	6558	wwPDB-VP
Average B, all atoms (Å^2)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.21% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: TRS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	B	0.55	0/1703	0.81	0/2312
1	L	0.58	0/1703	0.78	1/2312 (0.0%)
2	A	0.59	0/1615	0.73	0/2201
2	H	0.61	0/1657	0.78	1/2260 (0.0%)
All	All	0.58	0/6678	0.78	2/9085 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	3
1	L	0	1
2	H	0	1
All	All	0	5

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	L	108	ARG	NE-CZ-NH2	6.79	123.69	120.30
2	H	99	TYR	N-CA-C	-5.47	96.22	111.00

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	167	ASP	Peptide

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	B	170	ASP	Peptide
1	B	200	THR	Peptide
2	H	98	ASN	Peptide
1	L	201	SER	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1669	0	1613	31	2
1	L	1669	0	1613	42	1
2	A	1578	0	1557	33	1
2	H	1618	0	1597	33	1
3	A	8	0	12	1	0
4	A	3	0	0	0	0
4	B	5	0	0	0	0
4	H	4	0	0	1	0
4	L	4	0	0	0	0
All	All	6558	0	6392	129	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 129 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:27:GLN:NE2	1:L:28:ASP:O	1.95	1.00
1:L:81:GLU:N	1:L:81:GLU:OE2	1.96	0.98
2:A:97:GLY:CA	2:A:100(B):ILE:HD13	1.94	0.97
2:H:62:THR:CG2	2:H:66:LEU:HD13	1.95	0.96
2:A:97:GLY:HA2	2:A:100(B):ILE:HA	1.46	0.96

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:202:SER:OG	2:A:3:LYS:NZ[1_654]	1.85	0.35
1:B:27:GLN:NE2	1:B:213:GLU:CB[1_456]	1.89	0.31
1:L:202:THR:CG2	1:B:53:LYS:CG[1_644]	2.04	0.16

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	212/215 (99%)	196 (92%)	16 (8%)	0	100	100
1	L	212/215 (99%)	198 (93%)	14 (7%)	0	100	100
2	A	202/222 (91%)	191 (95%)	10 (5%)	1 (0%)	29	53
2	H	210/222 (95%)	199 (95%)	11 (5%)	0	100	100
All	All	836/874 (96%)	784 (94%)	51 (6%)	1 (0%)	51	77

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	184	SER

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	192/193 (100%)	168 (88%)	24 (12%)	4	10
1	L	192/193 (100%)	164 (85%)	28 (15%)	3	7

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	181/192 (94%)	164 (91%)	17 (9%)	8	19
2	H	186/192 (97%)	169 (91%)	17 (9%)	9	21
All	All	751/770 (98%)	665 (88%)	86 (12%)	5	12

5 of 86 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	137	ASN
2	A	35	SER
1	B	143	ASP
1	B	197	THR
2	A	63	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	189	HIS
2	A	81	GLN
2	A	170	GLN
2	A	98	ASN
2	H	100(C)	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	TRS	A	301	-	7,7,7	0.41	0	9,9,9	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TRS	A	301	-	-	5/9/9/9	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	301	TRS	N-C-C3-O3
3	A	301	TRS	C3-C-C2-O2
3	A	301	TRS	C2-C-C3-O3
3	A	301	TRS	N-C-C2-O2
3	A	301	TRS	C1-C-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	301	TRS	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	B	214/215 (99%)	0.38	15 (7%)	16	15	36, 60, 79, 89	0
1	L	214/215 (99%)	0.26	4 (1%)	66	69	34, 60, 84, 99	0
2	A	208/222 (93%)	0.09	11 (5%)	26	25	24, 41, 78, 96	0
2	H	214/222 (96%)	-0.03	5 (2%)	60	62	27, 46, 73, 90	0
All	All	850/874 (97%)	0.18	35 (4%)	37	36	24, 52, 79, 99	0

The worst 5 of 35 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	127	SER	5.0
2	A	160	SER	4.9
1	B	81	GLU	4.3
2	A	99	TYR	4.1
2	A	161	SER	4.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	TRS	A	301	8/8	0.89	0.16	38,39,41,41	0

6.5 Other polymers (i)

There are no such residues in this entry.