wwPDB X-ray Structure Validation Summary Report (i)

Feb 10, 2024 – 07:09 PM EST

PDB ID	:	2OSZ
Title	:	Structure of Nup58/45 suggests flexible nuclear pore diameter by intermolecular sliding
Authors	:	Melcak, I.; Hoelz, A.; Blobel, G.
Deposited on	:	2007-02-06
Resolution	:	2.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3168 (2.90-2.82)
Clashscore	141614	3438 (2.90-2.82)
Ramachandran outliers	138981	3348 (2.90-2.82)
Sidechain outliers	138945	3351 (2.90-2.82)
RSRZ outliers	127900	3103 (2.90-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	93	.% 49% 40% • 8%
1	B	93	2% 60% 30% • 8%
1	C	93	2% 57% 35% 8%
1	D	93	.% 53% 35% • 8%

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 2931 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nucleoporin p58/p45.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	86	713	455	123	132	3	0	0	0
1	B	86	713	455	123	132	3	3	0	0
1	C	86	713	455	123	132	3	0	0	0
1	D	86	713	455	123	132	3	2	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	323	GLY	-	cloning artifact	UNP P70581
A	324	SER	-	cloning artifact	UNP P70581
A	325	HIS	-	cloning artifact	UNP P70581
A	326	MET	-	cloning artifact	UNP P70581
B	323	GLY	-	cloning artifact	UNP P70581
B	324	SER	-	cloning artifact	UNP P70581
B	325	HIS	-	cloning artifact	UNP P70581
B	326	MET	-	cloning artifact	UNP P70581
C	323	GLY	-	cloning artifact	UNP P70581
C	324	SER	-	cloning artifact	UNP P70581
C	325	HIS	-	cloning artifact	UNP P70581
C	326	MET	-	cloning artifact	UNP P70581
D	323	GLY	-	cloning artifact	UNP P70581
D	324	SER	-	cloning artifact	UNP P70581
D	325	HIS	-	cloning artifact	UNP P70581
D	326	MET	-	cloning artifact	UNP P70581

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	25	25	25	0	0
			Total	O		
2	B	17	17	17	0	0
			Total	O		
2	C	17	17	17	0	0
			Total	O		
2	D	20	20	20	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Nucleoporin p58/p45

Chain A:



• Molecule 1: Nucleoporin p58/p45

Chain B:

• Molecule 1: Nucleoporin p58/p45

Chain C:



• Molecule 1: Nucleoporin p58/p45

Chain D:





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 43 2 2	Depositor
Cell constants a, b, c, α, β, γ	92.70Å 92.70Å 169.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.85 19.86 – 2.85	Depositor EDS
% Data completeness (in resolution range)	(Not available) (20.00-2.85) 90.5 (19.86-2.85)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.62 (at 2.83Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.250 , 0.288 0.250 , 0.282	Depositor DCC
R_{free} test set	1709 reflections (9.93%)	wwPDB-VP
Wilson B-factor (Å^2)	39.1	Xtriage
Anisotropy	0.254	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.35 , 58.0	EDS
L-test for twinning[2]	$< \|L\| > = 0.52$, $< L^2 > = 0.36$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.92	EDS
Total number of atoms	2931	wwPDB-VP
Average B, all atoms (Å^2)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 16.50% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.40	0/726	0.57	0/981
1	B	0.40	0/726	0.54	0/981
1	C	0.40	0/726	0.53	0/981
1	D	0.41	0/726	0.57	0/981
All	All	0.40	0/2904	0.55	0/3924

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	713	0	704	52	0
1	B	713	0	704	38	1
1	C	713	0	704	37	0
1	D	713	0	704	38	1
2	A	25	0	0	3	0
2	B	17	0	0	3	1
2	C	17	0	0	2	0
2	D	20	0	0	5	0
All	All	2931	0	2816	127	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 22.

The worst 5 of 127 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:342:GLN:HG2	1:A:388:GLN:HG3	1.48	0.92
1:A:342:GLN:CG	1:A:388:GLN:HG3	2.12	0.80
1:A:373:MET:HA	1:A:373:MET:HE2	1.62	0.80
1:C:339:PHE:HE2	1:D:357:LEU:HD11	1.50	0.77
1:B:367:THR:OG1	1:B:369:GLN:HG2	1.88	0.74

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:351:GLU:OE2	1:B:351:GLU:OE2[7_555]	2.16	0.04
2:B:14:HOH:O	2:B:14:HOH:O[7_555]	2.17	0.03
1:D:376:GLN:NE2	1:D:376:GLN:NE2[5_555]	2.18	0.02

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	84/93 (90%)	78 (93%)	6 (7%)	0	100	100
1	B	84/93 (90%)	80 (95%)	3 (4%)	1 (1%)	13	35
1	C	84/93 (90%)	79 (94%)	5 (6%)	0	100	100
1	D	84/93 (90%)	74 (88%)	8 (10%)	2 (2%)	6	19
All	All	336/372 (90%)	311 (93%)	22 (6%)	3 (1%)	17	43

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	365	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	363	ASN
1	D	363	ASN

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	77/81 (95%)	71 (92%)	6 (8%)	12	32
1	B	77/81 (95%)	74 (96%)	3 (4%)	32	63
1	C	77/81 (95%)	74 (96%)	3 (4%)	32	63
1	D	77/81 (95%)	71 (92%)	6 (8%)	12	32
All	All	308/324 (95%)	290 (94%)	18 (6%)	20	46

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	363	ASN
1	D	399	LEU
1	D	371	LEU
1	B	353	LEU
1	D	338	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 36 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	345	GLN
1	D	402	GLN
1	D	348	GLN
1	D	369	GLN
1	B	348	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	86/93 (92%)	-0.11	1 (1%)	79	78	15, 43, 63, 71	0
1	B	86/93 (92%)	-0.14	2 (2%)	60	57	17, 44, 63, 75	1 (1%)
1	C	86/93 (92%)	-0.16	2 (2%)	60	57	14, 40, 63, 70	0
1	D	86/93 (92%)	-0.22	1 (1%)	79	78	18, 39, 62, 74	1 (1%)
All	All	344/372 (92%)	-0.16	6 (1%)	70	68	14, 42, 63, 75	2 (0%)

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	363	ASN	3.0
1	B	363	ASN	2.6
1	C	362	ASN	2.6
1	C	363	ASN	2.5
1	B	362	ASN	2.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.