wwPDB X-ray Structure Validation Summary Report (i)

Jun 13, 2022 – 02:13 pm BST

PDB ID	:	7OSN
Title	:	IRED361 from Micromonospora sp. in complex with NADP+
Authors	:	Gilio, A.K.; Harawa, V.; Turner, N.; Grogan, G.J.
Deposited on	:	2021-06-09
Resolution	:	2.55 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.28.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0267
CCP4	:	7.1.010 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.28.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1284 (2.56-2.52)
Clashscore	141614	1332 (2.56-2.52)
Ramachandran outliers	138981	1315 (2.56-2.52)
Sidechain outliers	138945	1315 (2.56-2.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	297	71% 25% • •
1	B	297	72% 24% • •
1	C	297	73% 21% • •
1	D	297	76% 19% • •
1	E	297	77% 18% • •
1	F	297	76% 19% • •

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 12963 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 6-phosphogluconate dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	290	2124	1348	354	413	9	0	0	0
			Total	C	N	O	S			
1	B	291	2112	1346	353	404	9	0	0	0
			Total	C	N	O	S			
1	C	290	2125	1349	356	411	9	0	0	0
			Total	C	N	O	S			
1	D	290	2116	1343	355	409	9	0	0	0
			Total	C	N	O	S			
1	E	289	2002	1254	347	392	9	0	0	0
			Total	C	N	O	S			
1	F	289	2032	1278	350	396	8	0	0	0

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	48	21	7	17	3	0	0
2	B	1	48	21	7	17	3	0	0
2	C	1	48	21	7	17	3	0	0
2	D	1	48	21	7	17	3	0	0
2	E	1	48	21	7	17	3	0	0
2	F	1	48	21	7	17	3	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	29	29	29	0	0
3	B	41	41	41	0	0
3	C	43	43	43	0	0
3	D	30	30	30	0	0
3	E	10	10	10	0	0
3	F	11	11	11	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 6-phosphogluconate dehydrogenase

Chain A:



- Molecule 1: 6-phosphogluconate dehydrogenase

Chain B:

- Molecule 1: 6-phosphogluconate dehydrogenase

Chain C:



- Molecule 1: 6-phosphogluconate dehydrogenase

Chain D:



- Molecule 1: 6-phosphogluconate dehydrogenase

Chain E:



- Molecule 1: 6-phosphogluconate dehydrogenase

Chain F:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, α, β, γ	50.00Å 142.00Å 266.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.55 50.00 – 2.55	Depositor EDS
% Data completeness (in resolution range)	100.0 (50.00-2.55) 99.6 (50.00-2.55)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.27 (at 2.54Å)	Xtriage
Refinement program	REFMAC 5.0	Depositor
R, R_{free}	0.260 , 0.307 0.279 , 0.315	Depositor DCC
R_{free} test set	3255 reflections (5.18%)	wwPDB-VP
Wilson B-factor (Å^2)	51.4	Xtriage
Anisotropy	0.976	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	(Not available) , (Not available)	EDS
L-test for twinning[2]	$< \|L\| > = 0.51$, $< L^2 > = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.90	EDS
Total number of atoms	12963	wwPDB-VP
Average B, all atoms (Å^2)	78.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.34% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.69	0/2170	0.79	0/2973
1	B	0.69	0/2158	0.81	0/2959
1	C	0.68	0/2171	0.81	0/2975
1	D	0.70	0/2161	0.82	0/2961
1	E	0.70	0/2046	0.81	0/2811
1	F	0.71	0/2077	0.82	0/2854
All	All	0.70	0/12783	0.81	0/17533

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2124	0	2058	61	0
1	B	2112	0	2044	63	0
1	C	2125	0	2064	53	0
1	D	2116	0	2062	56	0
1	E	2002	0	1800	43	0
1	F	2032	0	1865	52	0
2	A	48	0	25	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	48	0	25	2	0
2	C	48	0	25	1	0
2	D	48	0	25	0	0
2	E	48	0	25	0	0
2	F	48	0	25	2	0
3	A	29	0	0	3	0
3	B	41	0	0	2	0
3	C	43	0	0	1	0
3	D	30	0	0	1	0
3	E	10	0	0	1	0
3	F	11	0	0	0	0
All	All	12963	0	12043	296	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 296 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:19:MET:O	1:F:23:LEU:HD12	1.69	0.93
1:F:69:ILE:HD13	1:F:95:VAL:HB	1.54	0.87
1:A:82:ILE:HG13	1:A:89:LEU:HD11	1.58	0.84
1:B:240:THR:O	1:B:243:VAL:HG22	1.78	0.83
1:B:248:MET:CE	1:B:269:LYS:HA	2.10	0.82

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	288/297 (97%)	272 (94%)	16 (6%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	289/297 (97%)	273 (94%)	16 (6%)	0	100	100
1	C	288/297 (97%)	274 (95%)	14 (5%)	0	100	100
1	D	288/297 (97%)	275 (96%)	13 (4%)	0	100	100
1	E	287/297 (97%)	271 (94%)	14 (5%)	2 (1%)	22	30
1	F	287/297 (97%)	270 (94%)	17 (6%)	0	100	100
All	All	1727/1782 (97%)	1635 (95%)	90 (5%)	2 (0%)	51	65

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	90	ALA
1	E	165	LEU

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	210/227 (92%)	192 (91%)	18 (9%)	10	13
1	B	205/227 (90%)	188 (92%)	17 (8%)	11	14
1	C	211/227 (93%)	189 (90%)	22 (10%)	7	7
1	D	210/227 (92%)	191 (91%)	19 (9%)	9	11
1	E	176/227 (78%)	161 (92%)	15 (8%)	10	13
1	F	184/227 (81%)	172 (94%)	12 (6%)	17	23
All	All	1196/1362 (88%)	1093 (91%)	103 (9%)	10	13

5 of 103 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	53	VAL
1	D	251	LEU
1	F	229	VAL
1	D	131	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	179	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 30 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	228	GLN
1	E	280	HIS
1	C	261	HIS
1	F	280	HIS
1	E	114	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	NAP	B	301	-	45,52,52	0.76	1 (2%)	56,80,80	0.98	1 (1%)
2	NAP	C	301	-	45,52,52	0.68	0	56,80,80	0.86	2 (3%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	NAP	E	301	-	45,52,52	0.87	2 (4%)	56,80,80	1.22	5 (8%)
2	NAP	D	301	-	45,52,52	0.66	1 (2%)	56,80,80	0.83	3 (5%)
2	NAP	A	301	-	45,52,52	0.67	1 (2%)	56,80,80	0.72	2 (3%)
2	NAP	F	301	-	45,52,52	0.85	2 (4%)	56,80,80	0.88	2 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	B	301	-	-	4/31/67/67	0/5/5/5
2	NAP	C	301	-	-	12/31/67/67	0/5/5/5
2	NAP	E	301	-	-	9/31/67/67	0/5/5/5
2	NAP	D	301	-	-	11/31/67/67	0/5/5/5
2	NAP	A	301	-	-	8/31/67/67	0/5/5/5
2	NAP	F	301	-	-	11/31/67/67	0/5/5/5

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	301	NAP	C2N-N1N	3.47	1.39	1.35
2	E	301	NAP	C2N-N1N	3.29	1.39	1.35
2	E	301	NAP	O4D-C1D	2.52	1.44	1.41
2	F	301	NAP	P2B-O2B	2.06	1.63	1.59
2	B	301	NAP	C8A-N7A	-2.03	1.31	1.34

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	E	301	NAP	C6N-N1N-C2N	-4.27	118.08	121.97
2	E	301	NAP	O4D-C1D-C2D	-3.92	101.20	106.93
2	B	301	NAP	C6N-N1N-C2N	-3.82	118.49	121.97
2	F	301	NAP	C6N-N1N-C2N	-3.25	119.02	121.97
2	E	301	NAP	C3B-C2B-C1B	-3.13	97.00	102.89

There are no chirality outliers.

5 of 55 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	NAP	C5D-O5D-PN-O3
2	A	301	NAP	C5D-O5D-PN-O1N
2	A	301	NAP	C5D-O5D-PN-O2N
2	A	301	NAP	O4D-C4D-C5D-O5D
2	A	301	NAP	C3D-C4D-C5D-O5D

There are no ring outliers.

4 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	301	NAP	2	0
2	C	301	NAP	1	0
2	A	301	NAP	5	0
2	F	301	NAP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand NAP B 301	
 Bond lengths	 Bond angles
 Torsions	 Rings

Ligand NAP C 301



Bond lengths



Bond angles



Torsions



Rings

Ligand NAP E 301	
 Bond lengths	 Bond angles
 Torsions	 Rings

Ligand NAP D 301	
 Bond lengths	 Bond angles
 Torsions	 Rings

Ligand NAP A 301	
 Bond lengths	 Bond angles
 Torsions	 Rings

Ligand NAP F 301



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around NAP A 301: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)	
	

Electron density around NAP B 301:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around NAP C 301:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around NAP D 301:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around NAP E 301:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around NAP F 301: $2mF_o$-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)	
	

6.5 Other polymers (i)

Unable to reproduce the depositors R factor - this section is therefore empty.