Full wwPDB X-ray Structure Validation Report (i)

Jun 17, 2024 – 07:28 PM EDT

PDB ID	:	5OTC
Title	:	Structure of the periplasmic binding protein (PBP) NocT from Agrobacterium tumefaciens C58 in complex with noroctopinic acid.
Authors	:	Vigouroux, A.; Morera, S.
Deposited on	:	2017-08-21
Resolution	:	2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	265	87% 9% •
1	B	265	8% 89% 8% •

2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 4100 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nopaline-binding periplasmic protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	256	1967	1259	322	374	12	0	0	0
1	B	256	1966	1258	321	375	12	0	0	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	25	MET	-	initiating methionine	UNP P35120
A	284	HIS	-	expression tag	UNP P35120
A	285	HIS	-	expression tag	UNP P35120
A	286	HIS	-	expression tag	UNP P35120
A	287	HIS	-	expression tag	UNP P35120
A	288	HIS	-	expression tag	UNP P35120
A	289	HIS	-	expression tag	UNP P35120
B	25	MET	-	initiating methionine	UNP P35120
B	284	HIS	-	expression tag	UNP P35120
B	285	HIS	-	expression tag	UNP P35120
B	286	HIS	-	expression tag	UNP P35120
B	287	HIS	-	expression tag	UNP P35120
B	288	HIS	-	expression tag	UNP P35120
B	289	HIS	-	expression tag	UNP P35120

- Molecule 2 is (2 {S})-5-azanyl-2-(2-hydroxy-2-oxoethylamino)pentanoic acid (three-letter code: AQK) (formula: $C_7H_{14}N_2O_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	13	7	2	4	0	0
2	B	1	13	7	2	4	0	0

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O			
3	A	1	4	2	2		0	0
3	A	1	4	2	2		0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	10	Total Cl 10 10	0	0
4	B	1	Total Cl 1 1	0	0

- Molecule 5 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 7 4 3	0	0

- Molecule 6 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: $C_6H_{14}O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
6	A	1	10	6	4	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
7	A	73	73	73	0	0
			Total	O		
7	B	16	16	16	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Nopaline-binding periplasmic protein

Chain A:



• Molecule 1: Nopaline-binding periplasmic protein

Chain B:





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, α, β, γ	113.99Å 113.99Å 37.80Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	37.31 – 2.20 37.31 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.9 (37.31-2.20) 99.6 (37.31-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$< I/\sigma(I) >$ [1]	1.27 (at 2.20Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, R_{free}	0.190 , 0.208 0.193 , 0.214	Depositor DCC
R_{free} test set	1398 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å^2)	46.3	Xtriage
Anisotropy	0.121	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.33 , 40.8	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.027 for -h,-k,l 0.034 for h,-h-k,-l 0.020 for -k,-h,-l	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	4100	wwPDB-VP
Average B, all atoms (Å^2)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.94% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, EDO, AQK, CL, PGE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.52	0/2005	0.63	0/2706
1	B	0.46	0/2004	0.63	0/2705
All	All	0.49	0/4009	0.63	0/5411

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1967	0	1987	12	0
1	B	1966	0	1983	8	0
2	A	13	0	0	0	0
2	B	13	0	0	0	0
3	A	24	0	36	2	0
4	A	10	0	0	0	0
4	B	1	0	0	0	0
5	A	7	0	10	1	0
6	A	10	0	14	0	0
7	A	73	0	0	0	0
7	B	16	0	0	0	0
All	All	4100	0	4030	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:66:MET:HB3	1:B:68:ILE:HD12	1.85	0.59
1:A:142:ILE:HA	1:A:146:GLN:HE21	1.69	0.57
1:B:197:LYS:HE3	1:B:220:ASN:OD1	2.11	0.50
1:B:160:VAL:HG13	1:B:202:ASP:HB2	1.95	0.49
1:A:247:ASP:HA	3:A:302:EDO:H22	1.95	0.49
1:A:174:MET:HE2	1:A:205:LEU:HD22	1.95	0.48
1:B:162:PHE:HB3	1:B:183:ILE:HG22	1.95	0.48
1:B:174:MET:HE2	1:B:205:LEU:HD22	1.95	0.47
1:A:155:LYS:HE2	3:A:304:EDO:H21	1.95	0.47
1:A:160:VAL:HG13	1:A:202:ASP:HB2	1.96	0.47
1:B:110:ARG:HD2	1:B:258:ILE:HG22	1.96	0.47
1:A:162:PHE:HB3	1:A:183:ILE:HG22	1.97	0.46
1:A:66:MET:HB3	1:A:68:ILE:HD12	1.98	0.44
1:A:172:ALA:HB2	5:A:318:PEG:H41	2.01	0.42
1:B:173:PHE:HA	1:B:234:LEU:HD21	2.02	0.42
1:A:47:ALA:O	1:B:194:MET:HG3	2.18	0.42
1:A:163:GLY:HA3	1:A:201:ILE:HD13	2.01	0.41
1:A:134:ILE:O	1:A:228:PRO:HB3	2.21	0.41
1:A:177:MET:HB3	1:A:177:MET:HE2	1.99	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	254/265 (96%)	247 (97%)	7 (3%)	0	100	100
1	B	254/265 (96%)	247 (97%)	7 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	508/530 (96%)	494 (97%)	14 (3%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	210/219 (96%)	203 (97%)	7 (3%)	38 49
1	B	210/219 (96%)	203 (97%)	7 (3%)	38 49
All	All	420/438 (96%)	406 (97%)	14 (3%)	38 49

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	88	ARG
1	A	110	ARG
1	A	158	GLU
1	A	162	PHE
1	A	219	ASP
1	A	237	LYS
1	A	282	LYS
1	B	64	LYS
1	B	110	ARG
1	B	145	GLU
1	B	158	GLU
1	B	219	ASP
1	B	237	LYS
1	B	282	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	146	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	273	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 21 ligands modelled in this entry, 11 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	EDO	A	302	-	3,3,3	0.70	0	2,2,2	0.13	0
3	EDO	A	307	-	3,3,3	0.60	0	2,2,2	0.09	0
2	AQK	A	301	-	12,12,12	1.40	3 (25%)	11,14,14	1.49	4 (36%)
3	EDO	A	304	-	3,3,3	0.52	0	2,2,2	0.40	0
2	AQK	B	301	-	12,12,12	1.41	2 (16%)	11,14,14	1.43	2 (18%)
3	EDO	A	306	-	3,3,3	0.61	0	2,2,2	0.20	0
5	PEG	A	318	-	6,6,6	0.27	0	5,5,5	0.21	0
3	EDO	A	303	-	3,3,3	0.67	0	2,2,2	0.26	0
3	EDO	A	305	-	3,3,3	0.56	0	2,2,2	0.30	0
6	PGE	A	319	-	9,9,9	0.14	0	8,8,8	0.11	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	302	-	-	1/1/1/1	-
3	EDO	A	307	-	-	1/1/1/1	-
2	AQK	A	301	-	-	7/13/13/13	-
3	EDO	A	304	-	-	0/1/1/1	-
2	AQK	B	301	-	-	6/13/13/13	-
3	EDO	A	306	-	-	1/1/1/1	-
5	PEG	A	318	-	-	0/4/4/4	-
3	EDO	A	303	-	-	0/1/1/1	-
3	EDO	A	305	-	-	1/1/1/1	-
6	PGE	A	319	-	-	2/7/7/7	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	301	AQK	CB-CA	-2.63	1.47	1.53
2	B	301	AQK	CA-C	-2.36	1.46	1.52
2	A	301	AQK	CB-CA	-2.32	1.48	1.53
2	A	301	AQK	CAI-CAK	-2.25	1.46	1.51
2	A	301	AQK	CA-C	-2.21	1.46	1.52

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	A	301	AQK	CAI-N-CA	-2.76	108.21	114.01
2	B	301	AQK	OXT-C-CA	2.34	121.17	113.40
2	A	301	AQK	OXT-C-O	-2.27	118.93	124.09
2	A	301	AQK	OXT-C-CA	2.10	120.37	113.40
2	B	301	AQK	OXT-C-O	-2.06	119.41	124.09
2	A	301	AQK	CAG-CB-CA	2.04	120.22	113.92

There are no chirality outliers.

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	AQK	CB-CA-N-CAI
2	A	301	AQK	N-CA-CB-CAG
2	B	301	AQK	N-CA-CB-CAG
2	A	301	AQK	C-CA-CB-CAG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	B	301	AQK	C-CA-CB-CAG
2	A	301	AQK	N-CAI-CAK-OAB
2	B	301	AQK	N-CAI-CAK-OAB
2	B	301	AQK	N-CAI-CAK-OAD
2	A	301	AQK	CAK-CAI-N-CA
2	A	301	AQK	N-CAI-CAK-OAD
2	B	301	AQK	CB-CA-N-CAI
2	A	301	AQK	C-CA-N-CAI
2	B	301	AQK	C-CA-N-CAI
6	A	319	PGE	C1-C2-O2-C3
6	A	319	PGE	C3-C4-O3-C5
3	A	302	EDO	O1-C1-C2-O2
3	A	307	EDO	O1-C1-C2-O2
3	A	305	EDO	O1-C1-C2-O2
3	A	306	EDO	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	302	EDO	1	0
3	A	304	EDO	1	0
5	A	318	PEG	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	256/265 (96%)	-0.31	0 100 100	31, 43, 65, 115	0
1	B	256/265 (96%)	0.18	21 (8%) 11 10	51, 71, 104, 127	0
All	All	512/530 (96%)	-0.06	21 (4%) 37 35	31, 58, 94, 127	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	159	GLY	6.2
1	B	27	ASP	4.6
1	B	29	LYS	3.7
1	B	131	GLN	3.1
1	B	263	ALA	3.0
1	B	129	LYS	2.9
1	B	130	THR	2.9
1	B	158	GLU	2.9
1	B	128	LEU	2.8
1	B	132	VAL	2.7
1	B	64	LYS	2.6
1	B	30	SER	2.5
1	B	91	ALA	2.5
1	B	156	ILE	2.4
1	B	260	ALA	2.3
1	B	253	LEU	2.2
1	B	259	ASP	2.2
1	B	88	ARG	2.1
1	B	155	LYS	2.1
1	B	227	GLY	2.1
1	B	89	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
5	PEG	A	318	7/7	0.53	0.29	74,74,76,77	0
4	CL	A	315	1/1	0.59	0.28	87,87,87,87	0
3	EDO	A	302	4/4	0.74	0.16	72,74,74,74	0
3	EDO	A	307	4/4	0.77	0.25	60,60,62,62	0
3	EDO	A	303	4/4	0.79	0.15	49,51,51,52	0
4	CL	A	308	1/1	0.82	0.13	85,85,85,85	0
3	EDO	A	305	4/4	0.84	0.10	78,79,79,79	0
4	CL	A	316	1/1	0.86	0.20	97,97,97,97	0
4	CL	A	314	1/1	0.88	0.07	78,78,78,78	0
3	EDO	A	306	4/4	0.88	0.17	65,66,67,68	0
2	AQK	A	301	13/13	0.89	0.19	40,45,60,65	0
6	PGE	A	319	10/10	0.89	0.20	68,70,72,72	0
4	CL	B	302	1/1	0.90	0.17	84,84,84,84	0
4	CL	A	310	1/1	0.91	0.10	73,73,73,73	0
4	CL	A	311	1/1	0.91	0.14	93,93,93,93	0
4	CL	A	317	1/1	0.93	0.12	73,73,73,73	0
2	AQK	B	301	13/13	0.93	0.15	57,69,82,84	0
3	EDO	A	304	4/4	0.93	0.16	60,61,61,62	0
4	CL	A	312	1/1	0.93	0.08	78,78,78,78	0
4	CL	A	309	1/1	0.95	0.09	68,68,68,68	0
4	CL	A	313	1/1	0.99	0.09	55,55,55,55	0

6.5 Other polymers (i)

There are no such residues in this entry.