



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 22, 2023 – 07:30 am BST

PDB ID : 8OW9  
Title : Crystal structure of Tannerella forsythia MurNAc kinase MurK in complex with N-acetylmuramic acid (MurNAc)  
Authors : Stasiak, A.C.; Gogler, K.; Fink, P.; Stehle, T.; Zocher, G.  
Deposited on : 2023-04-27  
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	292	
1	B	292	
1	C	292	
1	D	292	
1	E	292	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	292	 <p>21% 39% 48% 7% • 5%</p>

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 12864 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative novel MurNAc kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	277	2147	1373	365	396	13	0	0	0
1	A	277	2134	1366	359	395	14	0	1	0
1	D	277	2138	1369	359	396	14	0	1	0
1	E	277	2146	1374	362	396	14	0	1	0
1	C	274	2097	1343	352	389	13	0	0	0
1	F	276	2122	1357	360	391	14	0	1	0

There are 96 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	1	ALA	-	insertion	UNP A0A1D3UBL2
B	19	GLN	ARG	variant	UNP A0A1D3UBL2
B	21	ARG	GLN	variant	UNP A0A1D3UBL2
B	23	ILE	VAL	variant	UNP A0A1D3UBL2
B	129	ALA	THR	variant	UNP A0A1D3UBL2
B	281	LYS	-	expression tag	UNP A0A1D3UBL2
B	282	THR	-	expression tag	UNP A0A1D3UBL2
B	283	VAL	-	expression tag	UNP A0A1D3UBL2
B	284	LEU	-	expression tag	UNP A0A1D3UBL2
B	285	GLU	-	expression tag	UNP A0A1D3UBL2
B	286	HIS	-	expression tag	UNP A0A1D3UBL2
B	287	HIS	-	expression tag	UNP A0A1D3UBL2
B	288	HIS	-	expression tag	UNP A0A1D3UBL2
B	289	HIS	-	expression tag	UNP A0A1D3UBL2
B	290	HIS	-	expression tag	UNP A0A1D3UBL2
B	291	HIS	-	expression tag	UNP A0A1D3UBL2
A	1	ALA	-	insertion	UNP A0A1D3UBL2

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
A	19	GLN	ARG	variant	UNP A0A1D3UBL2
A	21	ARG	GLN	variant	UNP A0A1D3UBL2
A	23	ILE	VAL	variant	UNP A0A1D3UBL2
A	129	ALA	THR	variant	UNP A0A1D3UBL2
A	281	LYS	-	expression tag	UNP A0A1D3UBL2
A	282	THR	-	expression tag	UNP A0A1D3UBL2
A	283	VAL	-	expression tag	UNP A0A1D3UBL2
A	284	LEU	-	expression tag	UNP A0A1D3UBL2
A	285	GLU	-	expression tag	UNP A0A1D3UBL2
A	286	HIS	-	expression tag	UNP A0A1D3UBL2
A	287	HIS	-	expression tag	UNP A0A1D3UBL2
A	288	HIS	-	expression tag	UNP A0A1D3UBL2
A	289	HIS	-	expression tag	UNP A0A1D3UBL2
A	290	HIS	-	expression tag	UNP A0A1D3UBL2
A	291	HIS	-	expression tag	UNP A0A1D3UBL2
D	1	ALA	-	insertion	UNP A0A1D3UBL2
D	19	GLN	ARG	variant	UNP A0A1D3UBL2
D	21	ARG	GLN	variant	UNP A0A1D3UBL2
D	23	ILE	VAL	variant	UNP A0A1D3UBL2
D	129	ALA	THR	variant	UNP A0A1D3UBL2
D	281	LYS	-	expression tag	UNP A0A1D3UBL2
D	282	THR	-	expression tag	UNP A0A1D3UBL2
D	283	VAL	-	expression tag	UNP A0A1D3UBL2
D	284	LEU	-	expression tag	UNP A0A1D3UBL2
D	285	GLU	-	expression tag	UNP A0A1D3UBL2
D	286	HIS	-	expression tag	UNP A0A1D3UBL2
D	287	HIS	-	expression tag	UNP A0A1D3UBL2
D	288	HIS	-	expression tag	UNP A0A1D3UBL2
D	289	HIS	-	expression tag	UNP A0A1D3UBL2
D	290	HIS	-	expression tag	UNP A0A1D3UBL2
D	291	HIS	-	expression tag	UNP A0A1D3UBL2
E	1	ALA	-	insertion	UNP A0A1D3UBL2
E	19	GLN	ARG	variant	UNP A0A1D3UBL2
E	21	ARG	GLN	variant	UNP A0A1D3UBL2
E	23	ILE	VAL	variant	UNP A0A1D3UBL2
E	129	ALA	THR	variant	UNP A0A1D3UBL2
E	281	LYS	-	expression tag	UNP A0A1D3UBL2
E	282	THR	-	expression tag	UNP A0A1D3UBL2
E	283	VAL	-	expression tag	UNP A0A1D3UBL2
E	284	LEU	-	expression tag	UNP A0A1D3UBL2
E	285	GLU	-	expression tag	UNP A0A1D3UBL2
E	286	HIS	-	expression tag	UNP A0A1D3UBL2

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
E	287	HIS	-	expression tag	UNP A0A1D3UBL2
E	288	HIS	-	expression tag	UNP A0A1D3UBL2
E	289	HIS	-	expression tag	UNP A0A1D3UBL2
E	290	HIS	-	expression tag	UNP A0A1D3UBL2
E	291	HIS	-	expression tag	UNP A0A1D3UBL2
C	1	ALA	-	insertion	UNP A0A1D3UBL2
C	19	GLN	ARG	variant	UNP A0A1D3UBL2
C	21	ARG	GLN	variant	UNP A0A1D3UBL2
C	23	ILE	VAL	variant	UNP A0A1D3UBL2
C	129	ALA	THR	variant	UNP A0A1D3UBL2
C	281	LYS	-	expression tag	UNP A0A1D3UBL2
C	282	THR	-	expression tag	UNP A0A1D3UBL2
C	283	VAL	-	expression tag	UNP A0A1D3UBL2
C	284	LEU	-	expression tag	UNP A0A1D3UBL2
C	285	GLU	-	expression tag	UNP A0A1D3UBL2
C	286	HIS	-	expression tag	UNP A0A1D3UBL2
C	287	HIS	-	expression tag	UNP A0A1D3UBL2
C	288	HIS	-	expression tag	UNP A0A1D3UBL2
C	289	HIS	-	expression tag	UNP A0A1D3UBL2
C	290	HIS	-	expression tag	UNP A0A1D3UBL2
C	291	HIS	-	expression tag	UNP A0A1D3UBL2
F	1	ALA	-	insertion	UNP A0A1D3UBL2
F	19	GLN	ARG	variant	UNP A0A1D3UBL2
F	21	ARG	GLN	variant	UNP A0A1D3UBL2
F	23	ILE	VAL	variant	UNP A0A1D3UBL2
F	129	ALA	THR	variant	UNP A0A1D3UBL2
F	281	LYS	-	expression tag	UNP A0A1D3UBL2
F	282	THR	-	expression tag	UNP A0A1D3UBL2
F	283	VAL	-	expression tag	UNP A0A1D3UBL2
F	284	LEU	-	expression tag	UNP A0A1D3UBL2
F	285	GLU	-	expression tag	UNP A0A1D3UBL2
F	286	HIS	-	expression tag	UNP A0A1D3UBL2
F	287	HIS	-	expression tag	UNP A0A1D3UBL2
F	288	HIS	-	expression tag	UNP A0A1D3UBL2
F	289	HIS	-	expression tag	UNP A0A1D3UBL2
F	290	HIS	-	expression tag	UNP A0A1D3UBL2
F	291	HIS	-	expression tag	UNP A0A1D3UBL2

- Molecule 2 is N-acetyl-beta-muramic acid (three-letter code: AMU) (formula: C<sub>11</sub>H<sub>19</sub>NO<sub>8</sub>) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	N			O
2	B	1	Total	C	N	O	0	0
			20	11	1	8		
2	A	1	Total	C	N	O	0	0
			20	11	1	8		
2	D	1	Total	C	N	O	0	0
			20	11	1	8		
2	E	1	Total	C	N	O	0	0
			20	11	1	8		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Putative novel MurNAc kinase

Chain B: 



- Molecule 1: Putative novel MurNAc kinase


Chain A: 




- Molecule 1: Putative novel MurNAc kinase

Chain D: 






• Molecule 1: Putative novel MurNAc kinase



• Molecule 1: Putative novel MurNAc kinase



• Molecule 1: Putative novel MurNAc kinase





## 4 Data and refinement statistics i

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	113.42Å 113.42Å 294.76Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.13 – 2.70 49.13 – 2.70	Depositor EDS
% Data completeness (in resolution range)	98.5 (49.13-2.70) 90.2 (49.13-2.70)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.31 (at 2.69Å)	Xtrriage
Refinement program	PHENIX 1.16_3549	Depositor
R, $R_{free}$	0.257 , 0.276 0.258 , 0.273	Depositor DCC
$R_{free}$ test set	1495 reflections (3.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	49.8	Xtrriage
Anisotropy	0.308	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 24.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.468 for -h,k,-l	Xtrriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	12864	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	68.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.08% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AMU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.46	1/2182 (0.0%)	1.00	14/2954 (0.5%)
1	B	0.29	0/2192	0.57	3/2964 (0.1%)
1	C	0.39	0/2141	0.78	6/2901 (0.2%)
1	D	0.34	0/2186	0.69	3/2958 (0.1%)
1	E	0.42	0/2194	0.96	13/2967 (0.4%)
1	F	0.49	1/2170 (0.0%)	1.15	19/2937 (0.6%)
All	All	0.40	2/13065 (0.0%)	0.88	58/17681 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	F	0	3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	44	GLU	CB-CG	7.49	1.66	1.52
1	F	111	VAL	CB-CG2	5.02	1.63	1.52

The worst 5 of 58 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	89	LEU	CB-CG-CD2	11.05	129.78	111.00
1	F	49	LEU	CA-CB-CG	11.00	140.59	115.30
1	A	50	LEU	CA-CB-CG	9.68	137.57	115.30
1	F	240	GLY	N-CA-C	-9.59	89.12	113.10
1	E	84	LEU	CA-CB-CG	9.14	136.32	115.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	F	107	SER	Peptide
1	F	203	GLU	Sidechain
1	F	241	SER	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2134	0	2118	145	0
1	B	2147	0	2148	61	0
1	C	2097	0	2063	98	1
1	D	2138	0	2131	75	0
1	E	2146	0	2146	121	0
1	F	2122	0	2099	196	2
2	A	20	0	17	1	0
2	B	20	0	17	1	0
2	D	20	0	17	0	0
2	E	20	0	17	0	0
All	All	12864	0	12773	665	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 26.

The worst 5 of 665 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:242:VAL:HA	1:F:266:ARG:NH2	1.16	1.44
1:F:242:VAL:CA	1:F:266:ARG:NH2	2.05	1.19
1:D:41:MET:O	1:D:45:ILE:HD12	1.48	1.09
1:E:49:LEU:HG	1:E:50:LEU:N	1.67	1.09
1:E:49:LEU:HG	1:E:50:LEU:H	0.91	1.04

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:29:LYS:NZ	1:C:46:GLU:OE2[4_555]	1.96	0.24
1:F:46:GLU:OE1	1:F:77:ARG:NH1[3_555]	2.01	0.19
1:F:47:ARG:NH2	1:F:70:PRO:O[3_555]	2.10	0.10

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	276/292 (94%)	260 (94%)	15 (5%)	1 (0%)	34 60
1	B	275/292 (94%)	266 (97%)	9 (3%)	0	100 100
1	C	270/292 (92%)	255 (94%)	14 (5%)	1 (0%)	34 60
1	D	276/292 (94%)	265 (96%)	11 (4%)	0	100 100
1	E	276/292 (94%)	259 (94%)	16 (6%)	1 (0%)	34 60
1	F	275/292 (94%)	247 (90%)	23 (8%)	5 (2%)	8 21
All	All	1648/1752 (94%)	1552 (94%)	88 (5%)	8 (0%)	29 54

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	87	SER
1	F	90	ILE
1	F	113	ILE
1	F	242	VAL
1	E	49	LEU

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	226/245 (92%)	218 (96%)	8 (4%)	36	65
1	B	229/245 (94%)	223 (97%)	6 (3%)	46	75
1	C	220/245 (90%)	214 (97%)	6 (3%)	44	74
1	D	228/245 (93%)	221 (97%)	7 (3%)	40	69
1	E	229/245 (94%)	219 (96%)	10 (4%)	28	56
1	F	224/245 (91%)	213 (95%)	11 (5%)	25	52
All	All	1356/1470 (92%)	1308 (96%)	48 (4%)	35	65

5 of 48 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	238	PHE
1	C	230	ASP
1	E	270	GLU
1	C	9	SER
1	F	7	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	216	ASN
1	F	119	ASN
1	F	207	HIS
1	A	187	GLN
1	A	179	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	AMU	D	301	-	20,20,20	1.84	6 (30%)	24,28,28	1.28	2 (8%)
2	AMU	E	301	-	20,20,20	1.84	6 (30%)	24,28,28	1.19	3 (12%)
2	AMU	A	301	-	20,20,20	1.83	6 (30%)	24,28,28	1.19	2 (8%)
2	AMU	B	301	-	20,20,20	1.84	6 (30%)	24,28,28	1.10	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AMU	D	301	-	-	0/14/34/34	0/1/1/1
2	AMU	E	301	-	-	0/14/34/34	0/1/1/1
2	AMU	A	301	-	-	0/14/34/34	0/1/1/1
2	AMU	B	301	-	-	1/14/34/34	0/1/1/1

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	301	AMU	C7-N2	3.85	1.47	1.34
2	D	301	AMU	C7-N2	3.85	1.47	1.34
2	A	301	AMU	C7-N2	3.83	1.47	1.34
2	B	301	AMU	C7-N2	3.78	1.47	1.34
2	D	301	AMU	C2-N2	3.72	1.51	1.45

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	301	AMU	O5-C1-C2	3.06	112.59	109.52
2	D	301	AMU	C8-C7-N2	2.34	120.06	116.10
2	A	301	AMU	C8-C7-N2	2.24	119.89	116.10
2	E	301	AMU	C8-C7-N2	2.24	119.89	116.10
2	B	301	AMU	C8-C7-N2	2.20	119.82	116.10

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	301	AMU	O11-C10-C9-O3


There are no ring outliers.


2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	AMU	1	0
2	B	301	AMU	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	277/292 (94%)	0.39	19 (6%) 16 15	33, 71, 101, 115	0
1	B	277/292 (94%)	0.03	1 (0%) 92 93	31, 47, 64, 77	0
1	C	274/292 (93%)	0.37	14 (5%) 28 26	47, 72, 87, 96	0
1	D	277/292 (94%)	0.08	3 (1%) 80 82	34, 49, 62, 89	0
1	E	277/292 (94%)	0.33	5 (1%) 68 70	37, 70, 105, 116	0
1	F	276/292 (94%)	1.18	60 (21%) 0 0	62, 101, 120, 130	0
All	All	1658/1752 (94%)	0.40	102 (6%) 20 19	31, 65, 109, 130	0

The worst 5 of 102 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	58	PHE	8.0
1	F	130	ALA	6.9
1	F	17	LEU	6.6
1	F	63	PHE	6.5
1	F	239	ILE	5.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	AMU	A	301	20/20	0.91	0.24	57,66,77,82	0
2	AMU	B	301	20/20	0.94	0.18	30,39,47,49	0
2	AMU	E	301	20/20	0.94	0.16	53,60,66,68	0
2	AMU	D	301	20/20	0.96	0.17	33,45,58,64	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around AMU A 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around AMU B 301:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around AMU E 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.